

## ORIGINAL ARTICLE

# Aerobic denitrification in permeable Wadden Sea sediments

Hang Gao<sup>1,2,3</sup>, Frank Schreiber<sup>1</sup>, Gavin Collins<sup>1,4</sup>, Marlene M Jensen<sup>1,5</sup>, Joel E Kostka<sup>1,6</sup>, Gaute Lavik<sup>1</sup>, Dirk de Beer<sup>1</sup>, Huai-yang Zhou<sup>2,7</sup> and Marcel MM Kuypers<sup>1</sup>

<sup>1</sup>Nutrient Group, Max Planck Institute for Marine Microbiology, Bremen, Germany; <sup>2</sup>Guangzhou Institute of Geochemistry, Chinese Academy Sciences, Guangzhou, PR China and <sup>3</sup>Graduate School of Chinese Academy Sciences, Beijing, PR China

**Permeable or sandy sediments cover the majority of the seafloor on continental shelves worldwide, but little is known about their role in the coastal nitrogen cycle. We investigated the rates and controls of nitrogen loss at a sand flat (Janssand) in the central German Wadden Sea using multiple experimental approaches, including the nitrogen isotope pairing technique in intact core incubations, slurry incubations, a flow-through stirred retention reactor and microsensor measurements. Results indicate that permeable Janssand sediments are characterized by some of the highest potential denitrification rates ( $\geq 0.19 \text{ mmol N m}^{-2} \text{ h}^{-1}$ ) in the marine environment. Moreover, several lines of evidence showed that denitrification occurred under oxic conditions. In intact cores, microsensor measurements showed that the zones of nitrate/nitrite and  $\text{O}_2$  consumption overlapped. In slurry incubations conducted with  $^{15}\text{NO}_3^-$  enrichment in gas-impermeable bags, denitrification assays revealed that  $\text{N}_2$  production occurred at initial  $\text{O}_2$  concentrations of up to  $\sim 90 \mu\text{M}$ . Initial denitrification rates were not substantially affected by  $\text{O}_2$  in surficial (0–4 cm) sediments, whereas rates increased by twofold with  $\text{O}_2$  depletion in the 4–6 cm depth interval. In a well mixed, flow-through stirred retention reactor (FTSRR),  $^{29}\text{N}_2$  and  $^{30}\text{N}_2$  were produced and  $\text{O}_2$  was consumed simultaneously, as measured online using membrane inlet mass spectrometry. We hypothesize that the observed high denitrification rates in the presence of  $\text{O}_2$  may result from the adaptation of denitrifying bacteria to recurrent tidally induced redox oscillations in permeable sediments at Janssand.**

*The ISME Journal* (2010) 4, 417–426; doi:10.1038/ismej.2009.127; published online 10 December 2009

**Subject Category:** geomicrobiology and microbial contributions to geochemical cycles

**Keywords:** aerobic denitrification; nitrogen loss; permeable sediments; simultaneously  $\text{NO}_x^-$  and  $\text{O}_2$  respiration

## Introduction

Nitrogen (N) is primarily removed from coastal ecosystems by microbially mediated denitrification that occurs on the seafloor (Hulth *et al.*, 2005). Continental shelf sediments are important sites of N removal, which may account for 50–70% of oceanic

N loss (Codispoti *et al.*, 2001). Although the majority of continental margins is covered by coarse-grained relict sediments (Emery, 1968; Johnson and Baldwin, 1986), most previous biogeochemical research has focused on muddy or fine-grained sediments. Pore-water advection, driven mainly by pressure gradients from wave action and bottom currents interacting with surface topography, causes rapid solute exchange and allows direct transfer of suspended particles into permeable sediment strata. Recent studies indicate that advective transport leads to an acceleration of organic matter mineralization and a stimulation of biogeochemical cycling proportional to the extent of pore-water exchange (Huettel and Rusch, 2000; de Beer *et al.*, 2005; Werner *et al.*, 2006). Up- and downward flow of pore-water associated with migrating sandy sediment ripples generates vertical oscillations in oxic and anoxic conditions as redox zones move horizontally through the surface layer of the bed. The dynamic redox conditions found in permeable marine sediments resemble those found in wastewater treatment plants (Gray, 1990). In other words, high

Correspondence: MMM Kuypers, Nutrient Group, Max Planck Institute for Marine Microbiology, Celsiusstr. 1, Bremen D-28359, Germany.

E-mail: mkuypers@mpi-bremen.de

<sup>4</sup>Current address: Department of Microbiology and Environmental Change Institute, National University of Ireland, University Road, Galway, Ireland.

<sup>5</sup>Current address: Institute of Biology and Nordic Center of Earth Evolution, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark.

<sup>6</sup>Current address: Department of Oceanography, Florida State University, Leroy Collins Research Lab, Tallahassee, FL, USA.

<sup>7</sup>Current address: School of Ocean and Earth Science, Tongji University, Shanghai, PR China.

Received 12 August 2009; revised 27 October 2009; accepted 27 October 2009; published online 10 December 2009

transport rates of organic matter and electron acceptors from the water column into the seafloor and the presence of oscillating oxic/anoxic conditions allow marine sands to act as an efficient nutrient filter that may facilitate N removal. However, few studies have investigated N-loss by denitrification in coastal permeable sediments; of these studies, fewer still have considered the effects of advective pore water flows on the rates of denitrification (Cook *et al.*, 2006; Hunter *et al.*, 2006; Rao *et al.*, 2007, 2008; Gihring *et al.*, 2010). Further research is needed to determine the rates and controls of N removal from permeable marine sediments.

The current paradigm is that denitrification is an anaerobic process in marine sediments, and oxygen is believed to act as a major control of the process (Brandes *et al.*, 2007). Denitrification is considered to require completely anoxic conditions due to the fact that  $O_2$  acts as a competing electron acceptor for  $NO_3^-$  respiration and key enzymes of the denitrification pathways are inhibited by relatively small amounts of  $O_2$  (Tiedje *et al.*, 1982; Zumft, 1997; Shapleigh, 2006). However, in contrast to the observations made in natural environments, a large number of laboratory studies have reported that denitrification occurs under aerobic conditions in pure cultures of bacteria (Robertson and Kuenen, 1984; Ronner and Sorensson, 1985; Trevors and Starodub, 1987; Robertson *et al.*, 1995). Such findings suggest that denitrification may not always be so effectively inhibited by  $O_2$ . Microbiologists have defined aerobic denitrification as the co-respiration or co-metabolism of  $O_2$  and  $NO_3^-$ . Physiological studies show that microorganisms are able to use branches of their electron transport chain to direct electron flow simultaneously to denitrifying enzymes as well as to  $O_2$  (Robertson and Kuenen, 1988; Huang and Tseng, 2001; Chen *et al.*, 2003). Although some environmental studies suggest that denitrification can occur in the presence of  $O_2$  (Carter *et al.*, 1995; Bateman and Baggs, 2005; Rao *et al.*, 2007), substantial rates of aerobic denitrification have not yet been verified in the natural marine environment. Through a combination of new techniques using stable N isotopes for the direct determination of denitrification rates as well as the rapid quantification of aqueous gases over short time scales, the study of aerobic denitrification becomes more feasible.

The main objective of this study was to investigate the impact of  $O_2$  dynamics on N loss by denitrification in permeable marine sediments of the Wadden Sea. Taking advective transport into account, we investigated denitrification rates in permeable sediments under near *in situ* conditions using a variety of experimental approaches. Surprisingly, multiple lines of evidence indicated that denitrification was not inhibited in the presence of substantial oxygen concentrations but rather the co-respiration of  $O_2$  and  $NO_3^-$  occurred. Therefore, we hypothesized that where  $NO_3^-$  and  $O_2$  co-occur,  $O_2$  may not act as the

primary or exclusive control of  $N_2$  production in permeable sediment environments.

## Materials and methods

### Site description

The Janssand sand flat (13 km<sup>2</sup>) is in the back barrier area of Spiekeroog Island in the German Wadden Sea. The western edge of the flat faces the 17-m-deep tidal channel separating the barrier islands Spiekeroog and Langeoog. The entire Janssand flat is inundated with ~2 m of seawater for 6–8 h during each semi-diurnal tidal cycle, becoming exposed to air for 6–8 h during low tide, depending on the tidal range.

The central region of Janssand comprises the main area of the sand flat and is termed the upper flat due to the sloping margin downwards to the low water line. The upper flat is itself almost level and the physical appearance is homogeneous, mainly consisting of well-sorted silica sand with a mean grain size of 176  $\mu\text{m}$  (Billerbeck *et al.*, 2006a). The mean permeability of  $7.2\text{--}9.5 \times 10^{-12} \text{ m}^2$  (upper 15 cm of sediment; Billerbeck *et al.*, 2006a) permits advective pore water flows (Huettel and Gust, 1992). Detailed descriptions of the Janssand flat are available in Billerbeck *et al.* (2006a,b), Røy *et al.* (2008) and Jansen *et al.* (2009).

The sampling site (53°44.11'N, 7°41.95'E) is situated on the northeastern margin of the upper flat, about 80 m upslope from the mean low water line. Flat-bottom ships, the Spes Mea and Doris von Ochtum, were used to investigate the site in March 2007 and April 2008, respectively. All sediment core and seawater sampling was conducted at the upper Janssand tidal flat in March 2007, unless otherwise indicated.

### Dissolved inorganic nitrogen in sediment pore water

Rhizon samplers (Seeberg-Elverfeldt *et al.*, 2005) were used to extract pore water directly from sediment cores on the deck of the ship. Cores were sampled by pushing Plexiglas core liners (inner diameter (ID), 9.5 cm) with side ports into the sediment, and Rhizons were then inserted horizontally into the ports at 1 cm intervals to 10 cm depth. Site seawater was also collected during low tide and filtered through a 0.2- $\mu\text{m}$  syringe filter. All samples were immediately frozen onboard ship at  $-20^\circ\text{C}$  for later analysis. Dissolved ammonium ( $NH_4^+$ ) concentrations were determined using a flow injection analyzer (Hall and Aller, 1992). Nitrate + nitrite ( $NO_x^-$ ) was determined by chemiluminescence after reduction to NO with acidic vanadium(II) chloride (Braman and Hendrix, 1989).

### Intact core incubations

Denitrification rates were determined by the isotope pairing technique (Nielsen, 1992) in intact core

incubations modified according to De Beer *et al.* (2005) to simulate *in situ* pore water advection in the permeable sediments. A set of 15 sediment cores was collected in parallel to obtain 10 cm of sediment and 15 cm of overlying water each from a 1 m<sup>2</sup> area using Plexiglas push-cores (ID, 3.5 cm; height, 28 cm). Seawater was also collected in parallel from the site. Water overlying the sediment was removed and replaced with <sup>15</sup>NO<sub>3</sub><sup>-</sup>-amended site seawater (final concentration of 50 μM). Rubber bottom stoppers were fitted with valves to allow for pore water perfusion over the upper 5 cm depth, and each core was percolated by 20 ml <sup>15</sup>NO<sub>3</sub><sup>-</sup>-amended aerated seawater at the perfusion speed of 12 ml min<sup>-1</sup>. Cores were immediately sealed without any headspace by rubber stoppers after percolation, incubated at *in situ* temperature (8–10 °C) and were destructively sampled in triplicate at regular intervals between 0 and 6 h. The overlying water of all cores was mixed continuously at approximately 60 r.p.m. during the incubations by externally driven magnetic stirring bars. Cores were killed in reverse order of percolation. Briefly, 1 ml of zinc chloride (50% w/v) was added to the sediment surface. The cores were resealed with no headspace before mixing by inversion. After allowing the sediment particles to settle, an aliquot of water was removed from each core and transferred to a 12 ml Exetainer (Labco, High Wycombe, UK) pre-filled with 200 μl saturated HgCl<sub>2</sub> for <sup>29</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub> determinations. The concentrations of excess <sup>29</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub> were calculated from <sup>29</sup>N<sub>2</sub>:<sup>28</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub>:<sup>28</sup>N<sub>2</sub> ratios of He-equilibrated headspace in Exetainers determined by gas chromatography-isotope ratio mass spectrometry (GC-IRMS; VG Optima, Manchester, UK). Denitrification rates were calculated based on the linear production of excess <sup>29</sup>N<sub>2</sub> and <sup>30</sup>N<sub>2</sub> according to Nielsen (1992).

Another set of intact core incubations was conducted to provide direct evidence for the co-respiration of O<sub>2</sub> and NO<sub>x</sub><sup>-</sup> using multiple microsensors. An O<sub>2</sub> microsensor and a NO<sub>x</sub><sup>-</sup> biosensor were simultaneously applied to freshly sampled sediment cores. Advective transport in sediments cores was simulated by percolation as described above.

Oxygen microsensors were constructed as described previously (Revsbech, 1989). A two-point calibration method for the O<sub>2</sub> sensor was performed using the signal in saturated overlying site seawater and anoxic sediments, and the O<sub>2</sub> solubility was corrected for the ambient water temperature (18 °C) and salinity (32‰) using the spreadsheet supplied by Unisense ([www.unisense.dk](http://www.unisense.dk)). The NO<sub>x</sub> biosensor was constructed, according to Larsen *et al.* (1997), with a tip diameter of 100 μm and was calibrated in seawater with additions of increasing amounts of NO<sub>3</sub><sup>-</sup> to confirm linearity of the response (0–500 μM). For the calculation of pore water concentrations, the slope and offset were corrected for NO<sub>3</sub><sup>-</sup> concentrations in the overlying water determined as described above.

For simultaneous measurements of vertical concentration profiles, the O<sub>2</sub> and NO<sub>x</sub> sensors were mounted on a three-axis micromanipulator (MM 33; Märzhäuser, Wetzlar, Germany). The vertical axis was motorized for μ-positioning and controlled by μ-Profiler software described in Polerecky *et al.* (2005). The microsensor tips were aligned carefully to the same horizontal axis. The even sediment topography allowed alignment of both sensor profiles to the sediment surface with a precision of 1 mm, using the initial decrease of O<sub>2</sub> in the diffuse boundary layer. Microsensor measurements were made at 0.5 and 1 cm below the sediment surface during each percolation. Data were recorded over a time series to determine rates of potential O<sub>2</sub> uptake and NO<sub>x</sub><sup>-</sup> consumption under oxic and anoxic conditions.

#### *Slurry incubations in gas-tight bags*

The depth-specific response of denitrification to O<sub>2</sub> was initially examined in bag incubations using the <sup>15</sup>N tracer isotope pairing technique according to Thamdrup and Dalsgaard (2002). Sediments were sampled using Plexiglas push-cores (ID, 9.5 cm; height, 60 cm) and sectioned into 2-cm-depth intervals to a depth of 6 cm. Afterwards, sediment and air-saturated site seawater were mixed at a ratio of ~1:1 in the gas-tight bag while expelling all air bubbles. The slurries were amended with <sup>15</sup>NO<sub>3</sub><sup>-</sup> to a final concentration of 200 μM, and the bags were incubated at *in situ* temperature (same as for intact core incubations). During the incubation, the bags were periodically shaken to ensure that the labeled N species were homogeneously distributed. Subsamples of the interstitial water were withdrawn from the bags at regular intervals and preserved in 6 ml Exetainer vials (Labco) pre-filled with 100 μl saturated HgCl<sub>2</sub>. An initial subsample was taken immediately after the addition of the tracer and at regular intervals to 16 h.

The aqueous O<sub>2</sub> concentration of the subsamples was determined using an O<sub>2</sub> microelectrode (MPI, Bremen, Germany) as described above. The 6 ml Exetainer vials were opened only briefly during the measurement and were afterwards stored with no headspace for further analysis of dissolved N<sub>2</sub> by GC-IRMS as described above.

#### *FTSRR incubation experiment*

To provide further corroboration for the co-respiration of O<sub>2</sub> and NO<sub>3</sub><sup>-</sup> in a sediment slurry, we directly determined aqueous gases (O<sub>2</sub> and N<sub>2</sub>) in line by membrane inlet mass spectrometry (MIMS; GAM200, IPI) in a FTSRR system. Surface sediments (0–3 cm) and site seawater were collected from the upper flat in April 2008, and stored at 4 °C during transport to the laboratory. Sediments and site seawater were mixed at a ratio of 1:3 in the gas-tight FTSRR without headspace. The slurry

was vigorously mixed at 200 r.p.m. by a magnetic stir bar, and the incubation was carried out in the dark at room temperature. The FTSRR system consisted of a sealed cylinder chamber (Plexiglas, inner diameter 9 cm, height 6 cm) fitted with three ports for the input and output of water. The effluent pumped through a filter by one port from the chamber was injected directly into the membrane inlet using a peristaltic pump with the pumping speed of  $0.5 \text{ ml min}^{-1}$ . Gastight syringes connected to the chamber by the other two ports, each filled with 50 ml of air-saturated site seawater, provided replacement water during pumping.

Simultaneous online measurements of mass 28 ( $^{14}\text{N}^{14}\text{N}$ ), 29 ( $^{14}\text{N}^{15}\text{N}$ ), 30 ( $^{15}\text{N}^{15}\text{N}$ ), 40 (Ar), 31 ( $^{15}\text{NO}$ ), 32 ( $\text{O}_2$ ), 44 ( $^{14+14}\text{N}_2\text{O}/\text{CO}_2$ ), 45 ( $^{14}\text{N}^{15}\text{NO}$ ) and 46 ( $^{15}\text{N}^{15}\text{NO}$ ) were obtained by MIMS. A standard calibration curve was constructed, based on measurements obtained under both air-saturated and anoxic conditions using a two-point calibration for each. The slurry in the incubator was amended with  $^{15}\text{NO}_3^-$  to a final concentration of  $\sim 150 \mu\text{M}$ . Mass abundance signals were recorded by MIMS at 1 s time intervals and the flow-through samples were collected in 2 ml vials and stored at  $-20^\circ\text{C}$  for dissolved inorganic nitrogen (DIN) analysis as described above (see section Dissolved inorganic nitrogen in sediment pore water).

## Results


### DIN and $\text{O}_2$ penetration in permeable sediments

Zones of  $\text{O}_2$  penetration and  $\text{NO}_x^-$  depletion largely overlapped in the upper 2–3 cm of tidal flat sediments. During the winter/spring,  $\text{NO}_x^-$  concentrations with a mean value of  $\sim 67 \mu\text{M}$  were observed in the overlying seawater, whereas  $\text{NH}_4^+$  concentra-


tions were comparably 10 times lower ( $< 7 \mu\text{M}$ ) (Figure 1a). In surficial sediments, pore water  $\text{NO}_x^-$  decreased rapidly with depth to  $\sim 40 \mu\text{M}$  at 1 cm depth, and a minimum concentration was observed below 3 cm depth. Concomitantly, pore water  $\text{NH}_4^+$  increased to  $\sim 70 \mu\text{M}$  from the surface to 3 cm depth and remained consistently high ( $70\text{--}105 \mu\text{M}$ ) below that depth (Figure 1a). *In situ*  $\text{O}_2$  measurements in the upper flat from March 2006 showed that  $\text{O}_2$  penetrated to  $\sim 3 \text{ cm}$  during tidal inundation (Figure 1b) and  $\text{O}_2$  penetration depths of up to 5 cm were observed at other locations on Janssand tidal flat (Billerbeck *et al.*, 2006b; Jansen *et al.*, 2009). The decrease in  $\text{NO}_x^-$  was equivalent to approximately half of the observed increase of  $\text{NH}_4^+$  with depth (Figure 1a).

### Denitrification potential in intact cores and gastight bag incubations

Following with the overlap in  $\text{O}_2$  penetration and  $\text{NO}_x^-$  depletion, we observed the immediate and rapid production of  $^{15}\text{N}$ -labeled  $\text{N}_2$  in both incubations amended with  $^{15}\text{NO}_3^-$  throughout the first 4 h of incubation under oxic conditions (Figure 2). Our study of the intact core incubations was motivated by a previous study of  $\text{O}_2$  consumption using the same pore water percolation method that observed substantial  $\text{O}_2$  was present during the first 1–2 h of intact core incubations in March (Polerecky *et al.*, 2005; Billerbeck *et al.*, 2006b). In this study,  $^{29}\text{N}_2$  and  $^{30}\text{N}_2$  were produced linearly ( $r_{29\text{N}_2}^2 = 0.93$  and  $r_{30\text{N}_2}^2 = 0.91$ , respectively) without any lag during the first 2 h of incubation in the presence of  $\text{O}_2$  (Figure 2a). In bag incubation experiments conducted in parallel, high potential denitrification rates were observed in sediment slurries from the 0 to 2, 2 to 4 and 4 to 6 cm depth intervals in which



**Figure 1** Distribution of dissolved inorganic nitrogen species ( $\text{NH}_4^+$  and  $\text{NO}_x^-$  ( $\text{NO}_3^- + \text{NO}_2^-$ )) and  $\text{O}_2$  penetration in sediments from the upper flat at Janssand. (a)  $\text{NO}_x^-$  (closed circles) and  $\text{NH}_4^+$  (open circles) concentration profiles in permeable sediments in March 2007 during exposure. The daily mean values of  $\text{NO}_x^-$  and  $\text{NH}_4^+$  concentrations in overlying seawater are depicted as closed and open squares, respectively. (b)  $\text{O}_2$  penetration depth during a tidal cycle measured by two oxygen sensors in March 2006 (modified from Jansen *et al.*, 2009).



**Figure 2**  $^{15}\text{N}$ -labeled  $\text{N}_2$  production and oxygen consumption in whole-core and bag incubations. (a)  $^{29}\text{N}_2$  (black circles) and  $^{30}\text{N}_2$  (open circles) production from  $^{15}\text{NO}_3^-$  amendments in percolated whole-core incubations.  $\text{O}_2$  consumption and  $^{15}\text{N}$ -labeled  $\text{N}_2$  production versus time in  $^{15}\text{NO}_3^-$ -amended, oxic, gastight bag incubations with sediment from (b) 0 to 2 cm, (c) 2 to 4 cm and (d) 4 to 6 cm depth intervals.

initial  $\text{O}_2$  concentrations of  $\sim 95$ , 30 and 35  $\mu\text{M}$  were observed, respectively. Higher  $^{29}\text{N}_2$  and  $^{30}\text{N}_2$  production rates were observed in incubations from the 0 to

4 cm depth intervals where higher  $\text{NO}_x^-$  concentrations are observed in sediment pore waters (Figures 1a and 2b–d). No significant change in the denitrification rates was observed in the incubations under oxic conditions (during the first 4 h) in comparison to anoxic conditions (during the last 12 h) (Table 1). In the incubation from the deepest depth interval (4–6 cm) where  $\text{NO}_x^-$  was depleted, the lowest denitrification rates were observed whereas  $\text{O}_2$  depleted earliest ( $\sim 2$  h) (Table 1). Moreover at 4–6 cm depth, the rate under anoxic conditions was 2.3 times higher as that under oxic conditions (Figure 2d and Table 1). When extrapolated over the 0–5 cm depth interval, potential denitrification rates measured in percolated intact cores and bag incubations were in the same range (Table 1).

#### Microelectrode and biosensor measurements

Similar to the observations in the bag incubations, time series measurements by microsensors upon percolation of air-saturated and  $\text{NO}_3^-$ -rich seawater showed that  $\text{NO}_x^-$  and  $\text{O}_2$  were consumed simultaneously at 0.5–1 cm depth in intact sediment cores (Figure 3; Table 1).  $\text{O}_2$ - and  $\text{NO}_x^-$ -rich seawater was transported downwards into the sediment by percolation, which increased concentrations at 0.5 cm to  $\sim 240 \mu\text{M}$   $\text{O}_2$  and  $50 \mu\text{M}$   $\text{NO}_x^-$ , and at 1 cm to  $\sim 230 \mu\text{M}$   $\text{O}_2$  and  $40 \mu\text{M}$   $\text{NO}_x^-$ . Under these high initial  $\text{O}_2$  concentrations, a slight accumulation of 3–6  $\mu\text{M}$   $\text{NO}_x^-$  was detected after 2–3 min, followed by a substantial linear decrease in  $\text{NO}_x^-$  over the next 0.5–1.0 h of incubation in the presence of  $\text{O}_2$ .  $\text{NO}_x^-$  was consumed at a higher rate at 1 cm than at 0.5 cm under oxic conditions (Table 1). After  $\text{O}_2$  was completely consumed, the  $\text{NO}_x^-$  reduction rate increased slightly at 0.5 cm depth; however,  $\text{NO}_x^-$  consumption rates showed no significant difference under oxic and anoxic conditions at 1 cm depth (Table 1), which corresponded to the results observed in the bag incubations with sediments from 0 to 4 cm depth.

#### Aerobic denitrification in an FTSRR


To provide further evidence for the simultaneous consumption of  $\text{NO}_x^-$  and  $\text{O}_2$  in permeable sediments, we conducted an incubation in a stirred retention reactor, in which the slurry was vigorously and continuously mixed. Under constant mixing, substantial  $^{30}\text{N}_2$  production was again observed by real-time MIMS measurements in the presence of 32  $\mu\text{M}$   $\text{O}_2$  (Figure 4).  $^{15}\text{NO}_3^-$  was amended to the continuously stirred chamber 20 min after the start of the incubation in the presence of 128  $\mu\text{M}$   $\text{O}_2$ . Online MIMS analyses indicated that after an initial lag period of 1.1 h, significant  $^{30}\text{N}_2$  production occurred in the presence of 40  $\mu\text{M}$   $\text{O}_2$ . Concomitantly,  $\text{O}_2$  consumption slowed below that concentration. Simultaneously, there was a slight accumulation of  $\text{NO}_x^-$  (data not shown) during  $^{30}\text{N}_2$

**Table 1** Summary of denitrification rates measured in all incubations

Measurement	Investigated depth (cm)	Denitrification/NO <sub>x</sub> consumption (mmol N m <sub>sediment</sub> <sup>-3</sup> h <sup>-1</sup> )	
		Oxic	Anoxic
Intact core incubation	5	4.60 ± 0.46	
	Integrated to 5	0.23 ± 0.02 <sup>a</sup>	
Intact core by multi-microsensors	0.5	15.5 ± 0.04	21.9 ± 0.05
	1	22.0 ± 0.04	21.5 ± 0.04
Slurry incubation	0–2	6.40 ± 0.37	10.57 ± 3.20
	2–4	8.27 ± 0.32	9.63 ± 1.15
	4–6	2.72 ± 0.16	6.28 ± 0.84
Constant mixing, flow-through retention reactor incubation	Extrapolated to 5	0.32 ± 0.01 <sup>a</sup>	0.47 ± 0.07 <sup>a</sup>
	0–3	6.23 ± 0.07	
	Extrapolated to 3	0.187 ± 0.002 <sup>a</sup>	

The mean porosity of sediments in upper flat is 35% (Billerbeck *et al.*, 2006b).

<sup>a</sup>The unit is mmol N m<sub>sediment</sub><sup>-3</sup> h<sup>-1</sup>.




**Figure 3** Time series of O<sub>2</sub> (gray) and NO<sub>x</sub><sup>-</sup> (black) concentrations in an intact sediment core after percolation (indicated by arrows) with air-saturated overlying seawater containing ~60 μM NO<sub>x</sub><sup>-</sup>. O<sub>2</sub> and NO<sub>x</sub><sup>-</sup> microsensors were adjusted on a horizontal axis and measurements were carried out simultaneously. The first percolation treatment started at 0 h when sensors were positioned at 0.5 cm depth. The sensors were moved from 0.5 to 1 cm after 0.9 h and the second percolation began at 1.1 h when sensors were positioned at 1 cm depth.

production. However, during that period, <sup>29</sup>N<sub>2</sub> production was not concurrent with <sup>30</sup>N<sub>2</sub> production and the increase of NO<sub>x</sub><sup>-</sup>. In contrast, <sup>29</sup>N<sub>2</sub> began to accumulate only when NO<sub>x</sub><sup>-</sup> decreased at 1.5 h of incubation in parallel with a sevenfold higher rate of <sup>30</sup>N<sub>2</sub> production (Figure 4).

## Discussion

In permeable marine sediments of the Wadden Sea, zones of NO<sub>x</sub><sup>-</sup> and O<sub>2</sub> penetration often overlap to several centimeters depth due to pore water advection (Figure 1) (Werner *et al.*, 2006; Billerbeck *et al.*, 2006a,b; Jansen *et al.*, 2009). Further, previous O<sub>2</sub> percolation experiments that incorporated pore water advection, showed that during the spring season when NO<sub>x</sub><sup>-</sup> is at high concentration in the



**Figure 4** <sup>15</sup>N-labeled N<sub>2</sub> production and oxygen concentration versus time during the incubation of permeable sediments in the flow-through stirred retention reactor (FTSRR). Sediments were sampled from the 0 to 3 cm depth interval of the upper flat during April 2008. <sup>29</sup>N<sub>2</sub>, <sup>30</sup>N<sub>2</sub> and O<sub>2</sub> concentrations are depicted as open circles, black squares and open triangles, respectively.

overlying seawater, O<sub>2</sub> persisted in the bulk pore water over the first 1–2 h of incubation in intact cores of Wadden Sea sediments (Polerecky *et al.*, 2005; Billerbeck *et al.*, 2006b). From these observations, it could be inferred that where NO<sub>x</sub><sup>-</sup> and O<sub>2</sub> co-occur, O<sub>2</sub> may not act as the primary or exclusive control of N<sub>2</sub> production in permeable sediment environments. To test this assumption, we investigated N loss by denitrification in relation to O<sub>2</sub> dynamics. Several lines of independent evidence collected with multiple experimental approaches under near *in situ* conditions showed that denitrification occurs in the presence of oxygen. We observed the immediate and rapid consumption of NO<sub>x</sub><sup>-</sup> under air-saturated pore water in the intact core, and the directly determined production of <sup>15</sup>N-labeled N<sub>2</sub> in the presence of up to 90 μM O<sub>2</sub> in slurry incubations. Further, the rapid production of labeled N<sub>2</sub> was not diminished in a vigorously stirred system—FTSRR. Thus, our results strongly

suggest that aerobic denitrification makes a substantial contribution to N loss in permeable marine sediments.

The rates and mechanisms of N removal in permeable marine sediments remain in question. Few studies have quantified  $N_2$  production in coastal permeable sediments, and the rate measurements in this small but growing database vary widely, ranging from 0.1 to  $3\text{ mmol m}^{-2}$  per day (Laursen and Seitzinger, 2002; Vance-Harris and Ingall, 2005; Cook *et al.*, 2006; Rao *et al.*, 2007, 2008). However, researchers have now become aware of the fact that in experiments where pore water advection is absent or impeded, a realistic determination of diagenetic processes is not achieved (Jahnke *et al.*, 2000; Cook *et al.*, 2006). At present, at least two mechanisms have been proposed to explain denitrification in the presence of oxygen: (1) co-respiration of  $\text{NO}_x^-$  and  $\text{O}_2$  (Bateman and Baggs, 2005), and (2) closely coupled nitrification–denitrification in microenvironments isolated from bulk sediment pore water (Rao *et al.*, 2007). Bateman and Baggs (2005) provided one of the few observations of the contribution of aerobic denitrifying bacteria to denitrification potential in the environment. Using a combined stable isotope and acetylene inhibition approach, they were able to distinguish the relative contribution of nitrification and denitrification to  $\text{N}_2\text{O}$  production in arable soil. The results suggested that aerobic denitrification occurred at 20% water-filled pore space.

Although biogeochemical evidence exists for denitrification in the presence of oxygen in the marine environment (Hulth *et al.*, 2005; Hunter *et al.*, 2006; Brandes *et al.*, 2007; Rao *et al.*, 2007), significant rates of aerobic denitrification have not been verified until now. New techniques such as  $\text{NO}_x$  biosensors and stable N isotope tracers applied in conjunction with MIMS allowed for the further confirmation of aerobic denitrification. Rao *et al.* (2007, 2008) incorporated the effects of pore water advection, and in corroboration with our results, observed high rates of  $\text{N}_2$  production in flow-through columns of oxic permeable sediments. In continental shelf sediments of the South Atlantic Bight, Rao *et al.* (2007) observed that pore water nitrate was only above detection in the oxic zone. N released as  $\text{N}_2$  accounted from 80% to 100% of remineralized N, and the C:N ratio of regeneration supported the interpretation of  $\text{N}_2$  produced by denitrification. In the study by Rao *et al.* (2007), the addition of  $^{15}\text{N}$ -nitrate caused only a small and gradual rise in  $^{29}\text{N}_2$  and  $^{30}\text{N}_2$  production in sediment columns over up to 12 days of incubation. Only columns with anoxic outflow showed substantial  $^{29}\text{N}_2$  or  $^{30}\text{N}_2$  production. Thus, Rao *et al.* (2007) concluded that their evidence for aerobic denitrification was equivocal, and  $\text{N}_2$  production more likely occurred from coupled nitrification–denitrification in microenvironments.

In this study, we observed the rapid and immediate production of  $^{15}\text{N}$ -labeled  $\text{N}_2$  in the presence of

$\text{O}_2$  under a variety of experimental conditions. Oxygen and  $\text{NO}_x^-$  dynamics were directly determined in real time under well-mixed conditions in sediment slurries and intact core incubations. Microsensor measurements showed that  $\text{NO}_x^-$  and  $\text{O}_2$  consumption occurred simultaneously in intact cores (Figure 3). Further direct evidence for the co-respiration of  $\text{O}_2$  and  $\text{NO}_x^-$  was provided using  $^{15}\text{N}$  tracer experiments in slurries that were constructed with sediments from different depths. Successive incubation experiments showed the reliability and uniformity of aerobic denitrification rates, despite the fact that the experimental setup differed (including the amount of sediments, volume of associated water and starting concentration of labeled nitrate; Supplementary Table 1). Although the concentrations of  $^{29}\text{N}_2$  and  $^{30}\text{N}_2$  in the associated water varied, the denitrification rates normalized to sediment volume were in the same range, with the exception of the higher rate measured by the microsensor, which incorporated  $\text{NO}_3^-$  assimilation as well as denitrification (Table 1). Under the experimental conditions used,  $^{29}\text{N}_2$  could be attributed to coupled nitrification–denitrification or anammox in the slurry (Thamdrup and Dalsgaard, 2002). In contrast,  $^{30}\text{N}_2$  would only be produced by complete denitrification. Anammox was shown to comprise only a small percentage of  $\text{N}_2$  production in parallel slurry experiments conducted in gastight bags (Gao *et al.*, in preparation). Therefore, we conclude that the  $^{15}\text{N}$ -labeled  $\text{N}_2$  production is mainly contributed by denitrification, and the occurrence in the presence of  $\text{O}_2$  provided evidence for aerobic denitrification.

At each depth examined in slurry incubation, the potential denitrification rate under aerobic conditions was similar to that measured under anaerobic conditions. Moreover, the maximum denitrification rate was not observed in the deepest depth interval with the lowest initial  $\text{O}_2$  concentration, but rather in the surface 0–4 cm depth (Figures 2b–d). This suggests that the overlapping  $\text{NO}_x^-$  concentration may act together with  $\text{O}_2$  to control the denitrification rate. On the Janssand tidal flat during winter/spring, rapid denitrification is likely to be supported by the constant supply of  $\text{NO}_x^-$  advected from the overlying seawater (Gao *et al.*, in preparation). In short,  $\text{O}_2$  dynamics did not strongly affect N loss by denitrification in the presence of abundant  $\text{NO}_x^-$ , but rather denitrification coexisted with  $\text{O}_2$  respiration in permeable Wadden Sea sediments affected by advection.

To further exclude the possibility of anoxic microniches forming in our sediment incubations, we conducted an experiment in a vigorously mixed FTSRR. The initial production of  $^{30}\text{N}_2$  in the presence of  $40\ \mu\text{M}$   $\text{O}_2$  (Figure 4) provided evidence for the process of aerobic denitrification. The concomitantly suppressed  $\text{O}_2$  consumption may indicate that nitrate acts as a competitive electron acceptor to facultatively aerobic denitrifying bacteria.

Whereas at lower  $O_2$  concentrations later in the incubation (where an increased ratio of unlabeled  $NO_3^-$  was observed), the production of  $^{29}N_2$  indicated denitrification coupled to nitrification. Due to the variability in the mass abundance signals, we cannot exclude the possibility that some  $^{29}N_2$  was also produced in the early stages of the FTSRR incubation. Thus, we observed the mechanism for rapid denitrification under oxic conditions depended on two pathways— aerobic denitrification and denitrification coupled to nitrification.

During the FTSRR incubation, the bulk pore water was vigorously flushed by aerated seawater and the labeled isotope ratio was kept constant. Thus, the possibility that denitrification occurred in anoxic microzones can be completely excluded. In corroboration with our results, previous studies on the formation of anoxic microzones in particles and aggregates showed that the respiration capacity is simply not sufficient to create anoxia under high ambient  $O_2$  concentration, and anoxic microzones more likely form at around 10% of air saturation (under  $\sim 25 \mu M O_2$  in the bulk phase; Schramm *et al.*, 1999; Ploug, 2001). In our study, at  $O_2$  concentrations of  $\sim 20\%$  air saturation and above, the establishment of anoxic microzones would be unlikely. Given the larger grain sizes present in marine sands,  $O_2$  transport is enhanced by advection/interstitial fluid flow, which produces less steep  $O_2$  gradients at the sediment–water interface and within particles/aggregates compared with those that develop under pure diffusion conditions (Ploug, 2001). The above-mentioned experiments were conducted in a vertical flow system under nonturbulent uniform flow conditions. Thus, for the coarse-grained sediments in our well-mixed retention reactor experiments where the sediment slurry is exposed to turbulent aerated flow, anoxic microzones would not form. Therefore, we conclude that substantial N loss occurs by aerobic denitrification in the permeable Wadden Sea sediments.

Denitrification has long been considered as an anoxic biogeochemical process in marine and aquatic environments, and oxygen has been shown to inhibit denitrifying enzyme activity (Tiedje *et al.*, 1982; Hulth *et al.*, 2005; Brandes *et al.*, 2007). However, a phylogenetically and physiologically diverse group of microorganisms has been shown to denitrify in the presence of oxygen in laboratory studies (Zehr and Ward, 2002; Hayatsu *et al.*, 2008). Bacteria capable of aerobic nitrate respiration were cultured in abundance from freshwater soils and sediments (Carter *et al.*, 1995). Aerobic denitrifiers were further isolated from a variety of managed and natural ecosystems (Patureau *et al.*, 2000). Thus, the influence of oxygen on nitrate respiration activity appears to vary between microorganisms, with some strains able to respire nitrate at or above air saturation (Lloyd *et al.*, 1987; Hayatsu *et al.*, 2008). Microbiological studies have gone so far as to suggest that the capacity for denitrification under

aerobic conditions is the rule rather than the exception amongst ecologically important denitrifying microbial communities (Lloyd *et al.*, 1987).

Previous studies indicate that the diversity, as well as the metabolic activity, of bacterial communities is high in permeable sediment environments, likely due to increased transport of growth substrates and the removal of metabolites by advective exchange with the overlying water column (Hunter *et al.*, 2006; Mills *et al.*, 2008; Boer *et al.*, 2009). Denitrification in the marine environment is believed to be mediated by a group of facultative anaerobes that display a wide range in phylogenetic affiliation and metabolic capabilities (Zehr and Ward, 2002). In pristine ecosystems, nitrate concentrations are typically too low to select for large populations of denitrifying organisms, and denitrifiers are thought to rely on aerobic heterotrophy in conjunction with their denitrification capacity (Tiedje, 1988). In permeable marine sediments, up- and downwelling of pore water associated with sandy sediment ripples generates redox oscillations that may promote the microbially mediated oxidation and reduction of N species.

Although the consensus is that low or no  $O_2$  is required for the initiation of denitrification, most information on the  $O_2$  level at which denitrification starts comes from pure cultures. Denitrification has been observed in the laboratory at  $O_2$  concentrations approaching air saturation (Zehr and Ward, 2002), but previous environmental studies are equivocal with regard to the impact of  $O_2$  dynamics on denitrification. Large differences are observed in the expression and regulation of denitrification genes between species studied in pure culture (Shapleigh, 2006). The expression of denitrification genes was shown to require  $O_2$  in some cases, and the presence of denitrification intermediates may impact the denitrification rate in the presence of  $O_2$ . A possible explanation is that the accumulation of intermediates slows  $O_2$  respiration, particularly at low  $O_2$  levels, thereby slowing down the aerobic–anaerobic transition and allowing the expression of  $O_2$ -requiring denitrification genes (Bergaust *et al.*, 2008).

We hypothesize that the co-respiration of nitrate and  $O_2$  represents an adaptation of denitrifiers to recurrent tidally induced redox oscillations in permeable sediments of the Wadden Sea. Some evidence from pure cultures of denitrifying bacteria supports this hypothesis. For example, when the selective pressure of environmental redox changes was removed, the aerobic denitrification ability of *Paracoccus denitrificans* decayed (Dalsgaard *et al.*, 1995; Robertson *et al.*, 1995). Further, Bergaust *et al.* (2008) proposed that denitrifiers adapt to recurrent oscillations in oxygen concentrations through a protection mechanism, which consists of the coordinated expression and activity of the denitrification enzymes for survival during the rapid transition from oxic to anoxic conditions. A ‘bottle-neck effect’ was also proposed, whereby nitrifying

and denitrifying bacteria react to oxygen and nitrate in the environment by coordinating their respective activities. Schmidt *et al.* (2003) observed that the onset of the aerobic denitrification did not depend on oxygen sensitivity of the corresponding enzymes, but rather on regulation of redox-sensing factors at the transcriptional level. Our biogeochemical evidence corroborates microbiological studies to indicate a clear need to elucidate the significance and the controls of aerobic denitrification in permeable marine sediments.

In contrast to the paradigm that denitrification is an exclusively anaerobic process, our experiments point to aerobic denitrification and indicate that O<sub>2</sub> may not act as a primary or exclusive control of N<sub>2</sub> production in permeable marine sediments. We propose that the availability of NO<sub>x</sub><sup>-</sup> as well as O<sub>2</sub> limit the denitrification rate at depths of marine sands that are impacted by pore water advection. We can only speculate on the mechanism of aerobic denitrification at this time. Co-metabolism would imply that both NO<sub>x</sub><sup>-</sup> and O<sub>2</sub> are used simultaneously as electron acceptor in a single organism. Alternatively, separated denitrifying and oxygen respiring populations may be active within the community. In the first case, one would expect a competition for electrons within the electron transport chain, thus an enhanced denitrification upon oxygen depletion. In the second case, denitrification would be uncoupled entirely from the presence of oxygen, as denitrification is not kinetically inhibited by oxygen, nor can oxygen compete for electrons. In the FTSRR, we observed a pronounced effect of oxygen on denitrification rate whereas in other incubations less of an effect was found, indicating that both mechanisms may be present. Further research is needed to elucidate the true mechanisms of aerobic denitrification in permeable marine sediments.

## Acknowledgements

We thank the Captains Ronald Monas, Ole Pfeiler and colleagues Hans Roy, Stefan Jansen and Ingrid Dohrmann for their cheerful support on the ship and shipping time; Phyllis Lam for her constructive comments; Gabriele Klockgether and Daniela Franzke for technical supports. This research was supported by German Academic Exchange Center (Deutscher Akademischer Austausch Dienst, DAAD), Max-Planck-Society (MPG) and German Research Foundation (DFG). JEK was partially supported by the Hanse-Wissenschaftskolleg and by grants from the US National Science Foundation (OCE-0424967 and OCE-0726754).

## References

Bateman EJ, Baggs EM. (2005). Contributions of nitrification and denitrification to N<sub>2</sub>O emissions from soils at different water-filled pore space. *Biol Fert Soils* **41**: 379–388.

- Bergaust L, Shapleigh J, Frostegård Å, Bakken L. (2008). Transcription and activities of NO<sub>x</sub> reductases in *Agrobacterium tumefaciens*: the influence of nitrate, nitrite and oxygen availability. *Environ Microbiol* **10**: 3070–3081.
- Billerbeck M, Werner U, Bosselmann K, Walpersdorf E, Huettel M. (2006a). Nutrient release from an exposed intertidal sand flat. *Mar Ecol-Prog Ser* **316**: 35–51.
- Billerbeck M, Werner U, Polerecky L, Walpersdorf E, de Beer D, Huettel M. (2006b). Surficial and deep pore water circulation governs spatial and temporal scales of nutrient recycling in intertidal sand flat sediment. *Mar Ecol-Prog Ser* **326**: 61–76.
- Boer SI, Hedtkamp SIC, Beusekom JEE, Fuhrman JA, Boetius A, Ramette A. (2009). Time-, sediment depth-related variations in bacterial diversity, community structure in subtidal sands. *ISME J* **3**: 780–791.
- Braman RS, Hendrix SA. (1989). Nanogram nitrite and nitrate determination in environmental and biological-materials by vanadium(III) reduction with chemiluminescence detection. *Anal Chem* **61**: 2715–2718.
- Brandes JA, Devol AH, Deutsch C. (2007). New developments in the marine nitrogen cycle. *Chem Rev* **107**: 577–589.
- Carter J, Hsaio Y, Spiro S, Richardson D. (1995). Soil and sediment bacteria capable of aerobic nitrate respiration. *Appl Environ Microb* **61**: 2852–2858.
- Chen F, Xia Q, Ju LK. (2003). Aerobic denitrification of *Pseudomonas aeruginosa* monitored by online NAD(P)H fluorescence. *Appl Environ Microb* **69**: 6715–6722.
- Codispoti LA, Brandes JA, Christensen JP, Devol AH, Naqvi SWA, Paerl HW *et al.* (2001). The oceanic fixed nitrogen and nitrous oxide budgets: moving targets as we enter the anthropocene? *Sci Mar* **65**: 85–105.
- Cook PLM, Wenzhofer F, Rysgaard S, Galaktionov OS, Meysman FJR, Eyre BD *et al.* (2006). Quantification of denitrification in permeable sediments: insights from a two-dimensional simulation analysis and experimental data. *Limnol Oceanogr-Meth* **4**: 294–307.
- Dalsgaard T, Zwart Jd, Robertson LA, Kuenen JG, Revsbech NP. (1995). Nitrification, denitrification and growth in artificial *Thiosphaera pantotropha* biofilms as measured with a combined microsensor for oxygen and nitrous oxide. *FEMS Microbiol Ecol* **17**: 137–148.
- de Beer D, Wenzhofer F, Ferdelman TG, Boehme SE, Huettel M, van Beusekom JEE *et al.* (2005). Transport and mineralization rates in North Sea sandy intertidal sediments, Sylt-Rømø Basin, Wadden Sea. *Limnol Oceanogr* **50**: 113–127.
- Emery KO. (1968). Relict sands on continental shelves of the world. *Am Assoc Petrol Geo Bull* **52**: 445–464.
- Gihring TM, Canion A, Riggs A, Huettel M, Kostka JE. (2010). Denitrification in shallow, sublittoral Gulf of Mexico permeable sediments. *Limnol Oceanogr* **54**: 43–54.
- Gray NF. (1990). *Activated Sludge: Theory and Practice*. Oxford University Press: Oxford, United Kingdom.
- Hall PJ, Aller RC. (1992). Rapid, small-volume, flow injection analysis for ΣCO<sub>2</sub>, and NH<sub>4</sub><sup>+</sup> in marine and freshwaters. *Limnol Oceanogr* **37**: 1113–1119.
- Hayatsu M, Tago K, Saito M. (2008). Various players in the nitrogen cycle: diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Sci Plant Nutr* **54**: 33–45.
- Huang HK, Tseng SK. (2001). Nitrate reduction by *Citrobacter diversus* under aerobic environment. *Appl Microbiol Biot* **55**: 90–94.

- Huettel M, Rusch A. (2000). Transport and degradation of phytoplankton in permeable sediment. *Limnol Oceanogr* **45**: 534–549.
- Huettel M, Gust G. (1992). Impact of bioroughness on interfacial solute exchange in permeable sediments. *Mar Ecol-Prog Ser* **89**: 253–267.
- Hulth S, Aller RC, Canfield DE, Dalsgaard T, Engstrom P, Gilbert F *et al.* (2005). Nitrogen removal in marine environments: recent findings and future research challenges. *Mar Chem* **94**: 125–145.
- Hunter EM, Mills HJ, Kostka JE. (2006). Microbial community diversity associated with carbon and nitrogen cycling in permeable shelf sediments. *Appl Environ Microb* **72**: 5689–5701.
- Jahnke RA, Nelson JR, Marinelli RL, Eckman JE. (2000). Benthic flux of biogenic elements on the Southeastern US continental shelf: influence of pore water advective transport and benthic microalgae. *Cont Shelf Res* **20**: 109–127.
- Jansen S, Walpersdorf E, Werner U, Billerbeck M, Böttcher M, de Beer D. (2009). Functioning of intertidal flats inferred from temporal and spatial dynamics of O<sub>2</sub>, H<sub>2</sub>S and pH in their surface sediment. *Ocean Dynam* **59**: 317–332.
- Johnson HD, Baldwin CT. (1986). Shallow siliciclastic seas. In: Reading HG (ed). *Sedimentary Environments and Facies*, 2nd edn. Blackwell Scientific Publications: Oxford. pp 229–282.
- Larsen LH, Kjaer T, Revsbech NP. (1997). A microscale NO<sub>3</sub> biosensor for environmental applications. *Anal Chem* **69**: 3527–3531.
- Laursen AE, Seitzinger SP. (2002). The role of denitrification in nitrogen removal and carbon mineralization in Mid-Atlantic Bight sediments. *Cont Shelf Res* **22**: 1397–1416.
- Lloyd D, Boddy L, Davies KJP. (1987). Persistence of bacterial denitrification capacity under aerobic conditions—the rule rather than the exception. *FEMS Microbiol Ecol* **45**: 185–190.
- Mills HJ, Hunter E, Humphrys M, Kerkhof L, McGuinness L, Huettel M *et al.* (2008). Characterization of nitrifying, denitrifying, and overall bacterial communities in permeable marine sediments of the northeastern Gulf of Mexico. *Appl Environ Microb* **74**: 4440–4453.
- Nielsen LP. (1992). Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiol Ecol* **86**: 357–362.
- Patureau D, Zumstein E, Delgenes JP, Moletta R. (2000). Aerobic denitrifiers isolated from diverse natural and managed ecosystems. *Microbiol Ecol* **39**: 145–152.
- Ploug H. (2001). Small-scale oxygen fluxes and remineralization in sinking aggregates. *Limnol Oceanogr* **46**: 1624–1631.
- Polerecky L, Franke U, Werner U, Grunwald B, de Beer D. (2005). High spatial resolution measurement of oxygen consumption rates in permeable sediments. *Limnol Oceanogr-Meth* **3**: 75–85.
- Rao AMF, McCarthy MJ, Gardner WS, Jahnke RA. (2007). Respiration and denitrification in permeable continental shelf deposits on the South Atlantic Bight: rates of carbon and nitrogen cycling from sediment column experiments. *Cont Shelf Res* **27**: 1801–1819.
- Rao AMF, McCarthy MJ, Gardner WS, Jahnke RA. (2008). Respiration and denitrification in permeable continental shelf deposits on the South Atlantic Bight: N<sub>2</sub>:Ar and isotope pairing measurements in sediment column experiments. *Cont Shelf Res* **28**: 602–613.
- Revsbech NP. (1989). An oxygen microsensor with a guard cathode. *Limnol Oceanogr* **34**: 474–478.
- Robertson LA, Dalsgaard T, Revsbech NP, Kuenen JG. (1995). Confirmation of ‘aerobic denitrification’ in batch cultures, using gas chromatography and 15N mass spectrometry. *FEMS Microbiol Ecol* **18**: 113–119.
- Robertson LA, Kuenen JG. (1984). Aerobic denitrification—a controversy revived. *Arch Microbiol* **139**: 351–354.
- Robertson LA, Kuenen JG. (1988). Heterotrophic nitrification in thiosphaera-pantotropha—oxygen-uptake and enzyme studies. *J Gen Microbiol* **134**: 857–863.
- Ronner U, Sorensson F. (1985). Denitrification rates in the low-oxygen waters of the stratified baltic proper. *Appl Environ Microbiol* **50**: 801–806.
- Røy H, Lee JS, Jansen S, de Beer D. (2008). Tide-driven deep pore-water flow in intertidal sand flats. *Limnol Oceanogr* **53**: 1521–1530.
- Schmidt I, Sliemers O, Schmid M, Bock E, Fuerst J, Kuenen JG *et al.* (2003). New concepts of microbial treatment processes for the nitrogen removal in wastewater. *FEMS Microbiol Rev* **27**: 481–492.
- Schramm A, Santegoeds CM, Nielsen HK, Ploug H, Wagner M, Pribyl M *et al.* (1999). On the occurrence of anoxic microniches, denitrification, and sulfate reduction in aerated activated sludge. *Appl Environ Microb* **65**: 4189–4196.
- Seeberg-Elverfeldt J, Schlüter M, Feseker T, Kölling M. (2005). Rhizon sampling of pore waters near the sediment-water interface of aquatic systems. *Limnol Oceanogr-Meth* **3**: 361–371.
- Shapleigh J. (2006). The Denitrifying Prokaryotes. In: M Dworkin (ed). *The Prokaryotes: A 740 Handbook on the Biology of Bacteria: Volume 2: Ecophysiology and Biochemistry*. Springer-Verlag: New York, NY, pp 769–792.
- Thamdrup B, Dalsgaard T. (2002). Production of N<sub>2</sub> through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. *Appl Environ Microbiol* **68**: 1312–1318.
- Tiedje JM. (1988). Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: Zehnder AJB (ed). *Environmental Microbiology of Anaerobes*. John Wiley and Sons: New York. pp 179–244.
- Tiedje JM, Sextstone AJ, Myrold DD, Robinson JA. (1982). Denitrification: ecological niches, competition and survival. *Antonie Van Leeuwenhoek* **48**: 569–583.
- Trevors JT, Starodub ME. (1987). Effect of oxygen concentration on denitrification in freshwater sediment. *J Basic Microb* **27**: 387–391.
- Vance-Harris C, Ingall E. (2005). Denitrification pathways and rates in the sandy sediments of the Georgia continental shelf. *Geochim Cosmochim Acta* **69**: A578.
- Werner U, Billerbeck M, Polerecky L, Franke U, Huettel M. (2006). Spatial and temporal patterns of mineralization rates and oxygen distribution in a permeable intertidal sand flat (Sylt, Germany). *Limnol Oceanogr* **51**: 2549–2563.
- Zehr JP, Ward BB. (2002). Nitrogen cycling in the ocean: new perspectives on processes and paradigms. *Appl Environ Microb* **68**: 1015–1024.
- Zumft W. (1997). Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* **61**: 533–616.

Supplementary Information accompanies the paper on The ISME Journal website (<http://www.nature.com/ismej>)