

## Corrigendum

# Plasmablastic lymphomas and plasmablastic plasma cell myelomas have nearly identical immunophenotypic profiles

Francisco Vega<sup>1</sup>, Chung-Che Chang<sup>2</sup>, L Jeffrey Medeiros<sup>3</sup>, Mark M Udden<sup>4</sup>, Jeong Hee Cho-Vega<sup>5</sup>, Ching-Ching Lau<sup>6</sup>, Chris J Finch<sup>1</sup>, Regis A Vilchez<sup>4,7</sup>, David McGregor<sup>1</sup> and Jeffrey L Jorgensen<sup>3</sup>

<sup>1</sup>Department of Pathology, Baylor College of Medicine, Houston, TX, USA; <sup>2</sup>Department of Pathology, The Methodist Hospital, Weil Medical College of Cornell University, Houston, TX, USA; <sup>3</sup>Department of Hematopathology, University of Texas MD Anderson Cancer Center, Houston, TX, USA; <sup>4</sup>Department of Medicine, Baylor College of Medicine, Houston, TX, USA; <sup>5</sup>Department of Molecular Pathology, University of Texas MD Anderson Cancer Center, Houston, TX, USA; <sup>6</sup>Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA and <sup>7</sup>Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, TX, USA

*Modern Pathology* (2005) **18**, 873. doi:10.1038/modpathol.3800430

**Correction to:** *Modern Pathology* (2005) **18**, 806–815. doi:10.1038/modpathol.3800355

authors and in the affiliations. The correct authors' names and affiliations are above.

Following the online publication of the above paper, errors were identified in the name of one of the