## **GAMMA-RAY BURSTS**

## Huge explosion in the early Universe

Long gamma-ray bursts (GRBs) are bright flashes of high-energy photons that can last for tens of minutes; they are generally associated with galaxies that have a high rate of star formation and probably arise from the collapsing cores of massive stars, which produce highly relativistic jets (collapsar model1). Here we describe y- and X-ray observations of the most distant GRB ever observed (GRB 050904): its redshift2,3 (z) of 6.29 means that this explosion happened 12.8 billion years ago, corresponding to a time when the Universe was just 890 million years old, close to the reionization era4. This means that not only did stars form in this short period of time after the Big Bang, but also that enough time had elapsed for them to evolve and collapse into black holes.

GRB 050904 triggered the Burst Alert Telescope (BAT) on board the Swift<sup>5</sup> satellite on 4 September 2005 at 1:51:44 GMT. The spacecraft quickly slewed to allow observations by




Figure 1 | Light curve and spectral evolution of GRB 050904 as observed by the Burst Aler t Telescope (BAT) and the X-ray Telescope (XRT).

WT, windowed timing mode data; PC, photon counting data. Top, evolution of the gamma-ray burst (GRB) K-corrected 0.2-10 keV luminosity (the K-correction accounts for the redshift dependence of the luminosity in a given band of wavelengths). Error bars show 90% confidence. Times are referred to the BAT trigger. Rest-frame time is obtained by applying correction factor  $(1+z)^{-1}$  to the observer frame time. Gaps in XRT-PC data correspond to the part of the orbit when the satellite was not observing this GRB. Bottom, change in photon index ( $\Gamma$ , defined by the power law  $F(E) = E^{F+1}$ , where F(E) is the observed flux of energy E) of GRB 050904 during the observation. Spectra were modelled using a power law with two absorbing components (galactic and intrinsic).

the X-ray Telescope (XRT)<sup>67</sup>, which measured the burst for ten days after its onset. Figure 1 (top panel) shows the history of the burst. We shall present and discuss the GRB phenomenology from the point of view of the rest frame of its source.

The BAT light curve shows three main peaks: two of about 2 s at T+3.8 s and T+7.7 s, and a long-lasting one at about T+13.7 s, where T is the time of the burst onset. It also shows a weak peak at about T+64 s. The early XRT light curve shows a steep power-law decay with an index of  $-2.07 \pm 0.03$ ; two flares are superimposed at T+64 s (coincident with the last peak of the BAT light curve) and T+170 s. Although interrupted by the constraints of low-Earthorbit observation, the X-ray light curve reveals highly irregular intensity variations, probably due to the presence of flares for up to T+1.5 hours. At later times, flaring activity is not detected, leaving only a residual emission that is 106 times lower than the initial intensity.

The flares in the XRT light curve can be interpreted as late internal shocks related to central engine activity. In this scenario, they would have the same origin as the first  $\gamma$ -ray emission<sup>8-10</sup>, which would require the central engine to remain active for at least 5,000 seconds, consistent with the collapsar model<sup>1</sup>.

Spectral analysis was performed by selecting time intervals corresponding to characteristic phases of the light curve evolution. All spectra were well modelled by a single power law, with both galactic and intrinsic absorption components in the case of the XRT spectra. Figure 1 (bottom) shows the evolution with time of the photon index  $\Gamma$ . The BAT spectra have  $\Gamma$ = -1.2. If we exclude the spectrum of the first XRT flare at T+64 s, the XRT photon indices show a clear, decreasing trend from about -1.2 to about -1.8 in the first T+200 s. No further spectral evolution is present in later XRT data.

The overall phenomenology of GRB 050904 is not peculiar with respect to other GRBs at lower redshift. This suggests that the mechanisms of GRB explosions in the early Universe and today are similar.

Based on the likely existence of population I/II stars in galaxies that were already metalenriched at these high redshifts<sup>11</sup>, we expect about 10% of all bursts detected by Swift to be located at z≥5. A higher percentage would require an additional contribution to the high-redshift GRB population by metal-free population III stars, which are viable GRB progenitors for long-duration GRBs<sup>11</sup>. A more systematic search for GRB optical counterparts will increase the sample of these high-redshift GRBs, allowing us to probe the existence of metal-free massive stars of population III.

G. Cusumano¹, V. Mangano¹, G. Chincarini²³,

A. Panaitescu⁴, D. N. Burrows⁵, V. La Parola¹,

T. Sakamoto<sup>6,7</sup>, S. Campana<sup>2</sup>, T. Mineo<sup>1</sup>, G. Tagliaferri<sup>2</sup>, L. Angelini<sup>6</sup>, S. D. Barthelemy<sup>6</sup>, A. P. Beardmore<sup>8</sup>, P. T. Boyd<sup>6</sup>, L. R. Cominsky<sup>9</sup>, C. Gronwall<sup>5</sup>, E. E. Fenimore<sup>4</sup>, N. Gehrels<sup>6</sup>, P. Giommi<sup>10</sup>, M. Goad<sup>8</sup>, K. Hurley<sup>11</sup>, J. A. Kennea<sup>5</sup>, K. O. Mason<sup>12</sup>, F. Marshall<sup>6</sup>, P. Mészáros<sup>5,13</sup>, J. A. Nousek<sup>5</sup>, J. P. Osborne<sup>8</sup>, D. M. Palmer<sup>4</sup>, P. W. A. Roming<sup>5</sup>, A. Wells<sup>8</sup>, N. E. White<sup>6</sup>, B. Zhang<sup>14</sup>

<sup>1</sup>INAF-Istituto di Astrofisica Spaziale e Fisica Cosmica di Palermo, 90146 Palermo, Italy e-mail: cusumano@ifc.inafit

<sup>2</sup>INAF Osservatorio Astronomico di Brera, 23807 Merate, Italy; <sup>3</sup>Università degli studi di Milano-Bicocca, Dipartimento di Fisica, 20126 Milan, Italy; 4Los Alamos National Laboratory, PO Box 1663, Los Alamos, New Mexico 87545, USA; 5Department of Astronomy & Astrophysics and Department of Physics, Pennsylvania State University, Philadelphia, Pennsylvania 16802, USA; 6NA SA/Goddard Space Flight Center, Greenbelt, Maryland 20771, USA; National Research Council, TJ2114, Washington DC 20418, USA; Department of Physics and Astronomy, University of Lei cester, Leicester LE1 7RH, UK; Department of Physics and Astronomy, Sonoma State University, Rohnert Park, Cali fornia 9 4928-36 09, USA; \*\*A SI Science Data Center, 00044 Frascati, Italy; "Space Sciences Laboratory, University of California, Berkeley, California 94720-7450, USA; ™Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK; MDepartment of Physics, University of Nevada, Box 454002, Las Vegas, Nevada 89154-4002, USA

- MacFadyen, A. I., Woosley, S. E. & Heger, A. Astrophys. J. 550, 410-425 (2001).
- Taglia fer ri, G. et al. Astron. Astrophys. 443, L1-L5 (2005).
- 3. Kawai, N. et al. GCNCirc. 3937 (2005)
- Becker, R. H. et al. Astron. J. 122, 2850-2857 (2001).
- 5. Gehrels, N. et al. Astrophys. J. 611, 1005-1020
- Sakamoto T. et al. GCN Circ 3938 (2005).
- Mineo, T. et al. GCN Circ. 3920 (2005)
- 8. Burrows, D.N. et al. Science 309, 1833-1835 (2005).
- Zhang, B. et al. Astrophys. J. (in the press); preprint at <a href="http://arXiv.org/astro-ph/0508321">http://arXiv.org/astro-ph/0508321</a> (2005).
- Nousek, J. A. et al. Astrophys. J. (in the press); preprint at <a href="http://arXiv.org/astro-ph/0508332">http://arXiv.org/astro-ph/0508332</a> (2005).
- Bromm, V. & Loeb, A. Astrophys. J. (in the press); preprint at <a href="http://arxiv.org/astro-ph/0509303">http://arxiv.org/astro-ph/0509303</a> (2005).

Competing financial interests: declared none. Received 11 November 2005; accepted 23 November 2005. doi:10.1038/440164a

## CORRIGENDUM

## Developmental technology: Dogs cloned from adult somatic cells

Byeong Chun Lee, Min Kyu Kim, Goo Jang, Hyun Ju Oh, Fibrianto Yuda, Hye Jin Kim, M. Hossein Shamim, Jung Ju Kim, Sung Keun Kang, Gerald Schatten, Woo Suk Hwang Nature 436, 641 (2005)

Supplementary Table 1 of this communication has been replaced (9 March 2006; corrections shown in red) as the original peak values reported for two of the canine microsatellite markers (PEZO2, REN105LO3) were in error; also, those for PEZO8 have been removed (further details are available from B.C.L. at belee@snu.ac.kr). In the Table legend, the URL giving details of the markers has been updated. The patent application mentioned in the legend should originally have been declared as a competing financial interest. doi:10.1038/440164b

BRIEF COMMUNICATIONS ARISING online

www.nature.com/bca see Nature contents.