

Activity probe for *in vivo* profiling of the specificity of proteasome inhibitor bortezomib

Celia R Berkers^{1,2,6}, Martijn Verdoes^{1,5,6}, Eben Lichtman¹, Edda Fiebiger¹, Benedikt M Kessler¹, Kenneth C Anderson³, Hidde L Ploegh¹, Huib Ovaa^{1,2} & Paul J Galardy^{1,4}

Proteasome inhibitors, such as the dipeptide boronic acid bortezomib, are emerging as important tools in the treatment of the fatal hematologic malignancy multiple myeloma. Despite the recent US Food and Drug Administration approval of bortezomib (PS341, Velcade) for the treatment of refractory multiple myeloma, many of the basic pharmacologic parameters of bortezomib and its mode of action on myeloma cells remain to be determined. We describe the synthesis and use of a cell-permeant active site-directed probe, which allows profiling of proteasomal activities in living cells. When we compared proteasome activity patterns in cultured cells and crude cell extracts with this probe, we observed substantial differences, stressing the importance for bioassays compatible with live cells to ensure accuracy of such measurements. Using this probe, we investigated the *in vivo* subunit specificities of bortezomib and another inhibitor, MG132.

The proteasome inhibitor bortezomib (2) (PS341, Velcade; **Fig. 1**) is among the most promising pharmaceuticals developed for the treatment of multiple myeloma¹. The ability to target the proteasome pharmacologically without prohibitive toxicity is in itself remarkable: the proteasome is a large proteolytic complex, which is abundant in the cytosol and the nucleus, and it is considered central to many different aspects of cellular physiology, as polypeptides are continuously synthesized and destroyed to maintain steady-state protein levels². Protein turnover by the proteasome has many critical regulatory roles, including quality control by removing abnormal proteins that arise by mutation, metabolic damage or misfolding.

Eukaryotic proteasomes consist of a 20S core, which is most often the central part of higher-order complexes, of which the 26S proteasome is the major active species (**Fig. 2**). In 26S proteasomes, the 20S core particle is complexed at one or both ends with a 19S regulatory complex that modulates the entrance of substrates to the core and is required for the recognition of ubiquitin-tagged substrates and the unfolding of substrates prior to their threading into the proteolytic chamber³. Hybrid proteasomes, in which

different regulatory caps are attached to the 20S core, also exist. The 20S core is a hollow, barrel-shaped protein complex⁴ composed of 14 α and 14 β subunits arranged in four stacked rings with an overall architecture of $\alpha(1-7)\beta(1-7)\beta(1-7)\alpha(1-7)$. The two outer α rings form gated channels through which substrates enter and products exit the catalytic core. In addition, the outer α rings provide binding sites for regulatory complexes such as the 19S caps. The proteolytic activity of the 20S particle resides in the two inner β rings. The proteasome has three distinct catalytic activities, caspase-like, tryptic-like and chymotryptic-like, each of which can be roughly correlated to the action of an individual catalytic β subunit. These are termed $\beta 1$, $\beta 2$ and $\beta 5$, respectively, and can be replaced by three interferon-inducible subunits, termed $\beta 1i$, $\beta 2i$ and $\beta 5i$ to form immunoproteasomes—proteasomes with altered catalytic specificity favoring production of antigenic peptides. The secondary alcohol of the N-terminal threonine of the active β subunits acts as the nucleophilic species.

The dipeptide boronic acid bortezomib is a potent inhibitor of the proteasome's chymotryptic-like activity⁵. But variations in proteasomal composition alter the proteolytic specificities of the proteasome⁶⁻⁸ and may therefore affect the sensitivity of the proteasome to the action of bortezomib. Little is known about the composition of active proteasomes in living multiple myeloma cells. Furthermore, most investigators measure proteasomal activity in cell extracts using fluorogenic substrates, and as a result, it has been difficult to assess the extent of inhibition of individual catalytic activities *in vivo* after treatment with bortezomib. This underscores the need for improved tools that allow the accurate assessment of proteasomal activity *in vivo*.

To facilitate proteomic research, activity-based chemical probes have been developed for the study of serine, cysteine and threonine proteases⁹⁻¹¹. These probes form covalent adducts with active proteases of the class that is targeted, leading to a convenient readout of enzyme activity. Most of these probes, however, can not be used in intact living cells^{12,13}. Tags that are incorporated to provide an enzymatic activity readout, such as [¹²⁵I]iodotyrosine, often render probes impermeant to cell membranes, thereby

¹Department of Pathology, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, Massachusetts 02115, USA. ²Division of Cellular Biochemistry, Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX Amsterdam, the Netherlands. ³Department of Medical Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, Massachusetts 02115, USA. ⁴Program in Hematology/Oncology, Children's Hospital Boston and Dana-Farber Cancer Institute, 44 Binney Street, Boston, Massachusetts 02115, USA. ⁵Present address: Leiden Institute of Chemistry, Leiden University, P.O. Box 9502, 2300 RA Leiden, the Netherlands. ⁶These authors contributed equally to this work. Correspondence should be addressed to H.O. (h.ovaa@nki.nl).




Figure 1 | Representative structures of bortezomib and vinyl sulfone-based proteasome probes.

restricting their use to cell extracts. Furthermore, they are non-specific and target many proteases at the same time. Thus, such probes do not provide information on *in vivo* activities of specific proteases.

Peptide vinyl sulfone-based proteasome inhibitors^{9,14} such as ZL₃VS (1) (Fig. 1) modify catalytically active threonine residues through a Michael addition reaction, resulting in the formation of a covalent β -sulfonyl ether linkage. When a detectable tag is attached to such inhibitors, they can be used to label and detect proteasome active sites. Recently we set out to profile proteasome activity in multiple biological samples. A previously developed peptide vinyl sulfone-based inhibitor proved to be cell-permeant¹⁵, but the need for secondary chemical labeling procedures made this method unsuitable as a rapid screen of proteasome inhibition in multiple samples. In need for fast, reproducible activity readouts we reasoned that the use of small haptens, which can be recognized by antibodies, may allow retention of cell permeability of probes and allow detection of labeled enzymes directly by SDS-PAGE and Western blot analysis. Therefore we synthesized an inhibitor with a dansyl-sulfonamido-hexanoyl hapten as a fast and easy method to monitor proteasome activity in living cells. Using this probe, we could profile active proteasome subunits in cell extracts and cultured cells. Furthermore, we investigated the *in vivo* subunit specificities of the proteasome inhibitors bortezomib and MG132. We found that the β 1 and β 1i subunits are additional targets for bortezomib and that MG132 inhibits all subunits.

RESULTS

Synthesis and characterization of proteasome probes

We synthesized compound 3 (Fig. 1) containing a dansyl-sulfonamido-hexanoyl hapten, as well as control 4 in which the reactive vinyl sulfone moiety is reduced by hydrogenation, rendering the overall probe unreactive. Using a cell-based assay¹⁶, we tested proteasome inhibition by compounds 3 and 4, and compared the inhibition to that of other known inhibitors. We exposed cells

expressing a substrate fusion protein comprising ubiquitin, the amino acid arginine and green fluorescent protein (ubiquitin-Arg-GFP) to proteasome inhibitors. Efficient proteasome inhibition results in the accumulation of GFP, which can be quantitated by fluorescence activated cell sorting (FACS; Fig. 3a,b). Whereas treatment with increasing amounts of 3 or AdaAhx₃L₃VS, the parent compound, resulted in a substantial increase in fluorescence, treatment with the control inhibitor 4 was ineffective. The approximate half-maximal inhibitory concentration (IC₅₀) values for AdaAhx₃L₃VS and 3 were 3 μ M and 8 μ M, respectively (Fig. 3b). Thus, although the potency and/or bioavailability of compound 3 is somewhat lower than that of the parent compound, it is by far sufficient as a probe for activity measurements. By observing the relatively weak fluorescence emitted by the dansyl moiety of inhibitor 3, we confirmed cell permeability of 3 by fluorescence microscopy (Fig. 3c). After

incubation with 20 μ M of 3, we visualized fluorescence throughout the cytosol and nucleus of cultured HeLa and EL4 cells. The microscopy results conclusively demonstrate the cell permeability of inhibitor 3. We next determined the subunit specificity of inhibitor 3 by its ability to compete for proteasome subunit labeling with the fully characterized radiolabeled inhibitor Ada¹²⁵IYAhx₃L₃VS (5) (Figs. 1 and 3d), that we have previously shown to react with all active subunits of the proteasome^{14,15}. In a dose-dependent manner inhibitor 3 could compete with Ada¹²⁵IYAhx₃L₃VS for labeling of all active proteasome subunits in extracts from the multiple myeloma cell line MM.1S. We therefore conclude that compound 3 is a cell-permeant proteasome inhibitor that can be used to label all active proteasome subunits.

Probing proteasome activity in cell extracts

To directly visualize the proteins labeled by 3 in EL4 cell extracts, we took advantage of the probe's dansyl-sulfonamido-hexanoyl tag. We incubated cell extracts with increasing concentrations of 3 followed by SDS-PAGE and immunoblotting for the dansyl-sulfonamido-hexanoyl moiety (Fig. 4a). We observed a clear labeling profile, devoid of spurious cross-reactivity attributable to the antibody used. At a concentration of 300 nM, inhibitor 3 labeled all active proteasome subunits to a similar extent, despite weaker reactivity




Figure 2 | Schematic of the 26S proteasome.




Figure 3 | Characterization of compounds **3** and **4**. **(a,b)** FACS analysis results of ubiquitin-Arg-GFP-expressing U373¹² cells upon incubation with proteasome probes and proteasome inhibitors. Incubation with compound **3** resulted in an increase in fluorescence, comparable to that of known proteasome inhibitors, whereas incubation with inactive compound **4** did not **(a)**. **(b)** Graph shows the increase in fluorescence after a 5-h incubation with the indicated concentrations of compound **3** and the parent compound AdaAhx₃L₃VS, from which an approximate IC₅₀ value could be estimated. Graphs are of representative results obtained in two independent experiments. **(c)** Visualization by fluorescence microscopy of the distribution of compound **3** in EL4 and HeLa cells. Bar, 10 μm. **(d)** Competition experiment between inhibitor **3** and inhibitor **5**. Autoradiograph showing the proteasome subunits that have reacted with **5**.

of the probe towards the β_{2i} subunit. In the presence of increasing concentrations of inhibitor, the proteasomal active sites were rapidly saturated, and we observed a nonspecific high-molecular-weight labeling profile at the higher concentrations. This underscores the need for a careful titration of directed activity-based probes. Notably, subunits migrate slightly differently on the gels (Fig. 4a compared to Fig. 3d); the migration of a particular subunit-probe complex on gel is probe-dependent, which explains the subtle differences. To validate the accuracy of the method, we performed proteasome activity measurements with fluorogenic substrates in parallel. We incubated extracts with increasing concentrations of inhibitor **3** and then monitored the cleavage of SucLeuLeuValTyrAMC (chymotryptic-like activity), ZAlaArgArgAMC (trypsin-like activity) and ZLeuLeuGluβNA (caspase-like activity) to determine the remaining activities (Fig. 4b). The activity measurements with fluorogenic substrates corresponded well to the immunoblot profile. At a concentration of 10 μM of **3**, labeling of subunits with this inhibitor reached saturation, and activity against fluorogenic substrates was abolished almost completely, thus confirming the accuracy of the use of inhibitor **3** as a probe to measure proteasome activity *in vitro*.

Probing proteasome activity in living cells

We next probed proteasomal activity in intact cells. We incubated cultured EL4 cells with increasing concentrations of **3**, collected the cells, and immunoblotted for the dansyl group as described above (Fig. 4c). We observed a dose-dependent labeling of active proteasome subunits from EL4 cells, which confirms that probe **3** permeates cell membranes and modifies all active proteasome

subunits. When we compared the labeling pattern obtained in experiments with EL4 extracts to that obtained in experiments with cultured cells, we found that elevated concentrations of probe are needed to efficiently label active proteasome subunits, probably owing to kinetics of membrane permeation. Furthermore, the *in vivo* labeling profile differed in comparison to the labeling pattern in cell extracts. In EL4 cell lysates, the β₁ and β_{1i} subunits appear to be quite active. But *in vivo* the β₁ subunit was labeled only in the presence of high concentrations of inhibitor **3**, whereas the β_{1i} subunit was not labeled at all. In contrast, the β₅ subunits seem to be more active and therefore the target for the majority of the observed *in vivo* active-site labeling. The exact reason for the observed activity difference is not clear. Possible explanations include partial dissociation of associated (19S) regulatory proteasomal cofactors upon preparation of extracts, as well as disruption of transient associations of active proteasomes with cell membranes in detergent extracts. In addition, when we compared *in vivo* labeling experiments to experiments in lysates, virtually no high-molecular-weight adducts were observed despite the presence of equal concentrations

of inhibitor. Thus, inhibitor **3** labels proteasomes with greater selectivity *in vivo*. Next, we validated the *in vivo* activity pattern obtained with probe **3** by performing activity measurements with fluorogenic substrates in extracts prepared from identical samples (Fig. 4d). *In vivo*, the saturation of subunit labeling corresponded to the inhibition of proteasome activity. At 10 μM, compound **3** almost completely inhibited proteasome activity, as was the case *in vitro*, confirming that it is unlikely that an altered bioavailability of compound **3** *in vivo* is responsible for the altered labeling pattern. Next, we used inhibitor **3** to profile the proteasome activity in HeLa cells *in vivo*. Again, the β₅ subunits seemed to be most active and therefore the targets for the majority of the *in vivo* active-site labeling (Supplementary Fig. 1 online). We conclude that proteasome activity profiles obtained by performing measurements in cell extracts are not necessarily representative of the *in vivo* activity patterns, stressing the need for live cell-based assays.

In vivo profiling of bortezomib and MG132

Next, we determined whether the probe could be used to accurately measure the remaining proteasome activity after inhibition by other proteasome inhibitors. We cultured EL4 cells in the presence of bortezomib or MG132, then incubated with inhibitor **3** and immunoblotted as described above. We performed activity measurements with fluorogenic substrates in parallel before and after addition of inhibitor **3**. The addition of bortezomib resulted in a very specific reduction in the ability of compound **3** to modify the proteasomal β₅, β_{5i} and β₁ subunits, with maximum inhibition of these subunits by bortezomib reached at 40 nM (Fig. 5a). The β₂ and β_{2i} subunits remained fully available for modification with

compound **3** at all time points, indicating the lack of a substantial interaction of bortezomib with these subunits *in vivo*. Bortezomib was previously reported to be a specific inhibitor for the chymotryptic-like activity largely provided by the $\beta 5$ subunits. Here, we identified the $\beta 1$ subunits as additional targets for bortezomib.

After incubation of EL4 cells with MG132, compound **3** only labeled the $\beta 2$ subunit suggesting that all other subunits are inhibited by MG132 at high-nM concentrations (Fig. 5b). Notably, MG132 completely inhibited the $\beta 2i$ subunit, which suggests a difference in the active-site chemistry between the $\beta 2$ and $\beta 2i$ subunits. The results we obtained with fluorogenic substrates correlated in both cases to the immunoblot experiments using inhibitor **3** (Fig. 5c,d). Inhibitor **3**, however, can be used to distinguish between constitutive and immunoproteasome subunit activity, whereas fluorogenic substrates cannot. Therefore, we did not observe the difference in labeling between the $\beta 2i$ subunit and the $\beta 2$ subunit using fluorogenic substrates. Combined, these results validate inhibitor **3** as a probe for measuring proteasome activity as well as for characterization of subunit specificity of proteasome inhibitors *in vivo*.

Using inhibitor **3**, we next determined the composition of the active proteasome in the multiple myeloma cell line MM.1S, as well as the proteasomal targets of bortezomib in these cells. We cultured MM.1S cells in the presence or absence of 20 nM bortezomib, a concentration previously demonstrated to be toxic to this cell line¹⁷. In this cell line, we observed that all constitutive

and immunoproteasome catalytic subunits were active in the absence of interferon γ (Fig. 5e). The labeling pattern in MM.1S cells resembled the profile that was observed in EL4 cells. A notable difference was the activity of the $\beta 1i$ subunit in living MM.1S cells: we detected this subunit in extracts of EL4 cells, but not when we performed the assay in living cells. As quickly as in 15 min, 20 nM bortezomib substantially inhibited the activity of




Figure 4 | Profiling of proteasome activity in EL4 cell extracts and cells. (a,c) Immunoblots originating from the same gel showing results of a 1-h labeling experiment in EL4 cell extracts (a) or a 2-h labeling experiment in living EL4 cells (c) with the indicated concentrations of inhibitor **3** or 10 μM inactive control **4**. (b,d) Measurement with fluorogenic substrates of residual proteasome activity in EL4 cell extracts (b) or living EL4 cells (d) after a 1-h (b) or 2-h (d) treatment with the indicated concentrations of inhibitor **3**. Results are representative of two independent experiments.




Figure 5 | *In vivo* profiling of proteasome inhibitors. (a,b) Competition experiment between inhibitor **3** and bortezomib (a) or MG132 (b) in EL4 cells. (c,d) Measurement of residual proteasome activity with fluorogenic substrates after treatment of EL4 cells with bortezomib or a combination of bortezomib and inhibitor **3** (c) or MG132 or a combination of MG132 and inhibitor **3** (d). Results are representative of two independent experiments. CtL: chymotryptic-like, TL, tryptic-like; CL, caspase-like. (e) Competition experiment between bortezomib and inhibitor **3** in MM.1S cells.

all $\beta 5$ and $\beta 1$ subunits with maximum inhibition reached within 2 h (Fig. 5e).

DISCUSSION

Of the proteasome inhibitors described to date, each has a varying ability to inhibit each of the three known catalytic activities^{14,18,19}. With proteasome inhibition becoming widely used in the treatment of multiple myeloma, the composition of proteasomes may be an important determinant of clinical response to treatment with such agents. Most methodologies in common practice for measuring proteasome activity use fluorogenic peptidyl substrates and often involve procedures to reduce potentially cross-reacting protease activities^{5,20,21}. More importantly, procedures that rely on the hydrolysis of fluorogenic substrates do not distinguish between constitutive and immunoproteasome activity and therefore do not reflect the relative activity of different proteasome subunits accurately.

Recently, cell-based assays^{16,22} as well as mouse models^{23,24} have been developed for *in vivo* monitoring of proteasome activity. These methods may be useful in screening for new inhibitors and testing bioavailability, respectively, but their activity readout depends on the balance between synthesis and degradation of fusion proteins, which involves many cellular factors other than the proteasome, including the rate of fusion-protein synthesis. Therefore these methods may not accurately reflect true proteasomal activity. Moreover, the use of expressed fusion proteins makes these methods unfit for monitoring human samples. In addition, they do not provide insight in the constitutive/immunoproteasome activity ratio and cannot be used to characterize the specific activity of proteasome inhibitors *in vivo*. The small molecule-based assay described here overcomes these problems.

We have devised proteasome activity-profiling probe **3** that freely reaches cellular targets in intact cells and that can modify covalently and irreversibly all of the proteasome's exposed catalytic subunits. Use of antibodies directed against the dansyl-sulfonamido-hexanoyl hapten present in **3** allows direct measurement of activity of all proteasome subunits in living cells without the need for prior lysis or purification steps. This ensures not only a fast but also accurate activity readout, as under these conditions proteasomes remain in their native state. Using this probe, we characterized the anti-neoplastic agent bortezomib and revealed that the $\beta 1$ and $\beta 1i$ subunits are targets for this drug, in addition to the previously identified $\beta 5$ subunits²⁰. We also characterized the widely used proteasome inhibitor MG132 and found that it inhibited all subunits, although the $\beta 2$ subunit was inhibited to a lesser extent. Notably, MG132 was found to fully inhibit the immunoproteasome subunit $\beta 2i$.

We have described an easy method for characterizing the specificity of (new) proteasome inhibitors that may be helpful in a clinical setting and could be used to study resistance to this new class of drugs. Differences in labeling profiles observed in experiments with extracts and living cells, which can not be detected using methodology now used to measure proteasome activity, underscores the need for reliable *in vivo* labeling techniques. The technique described here could be adapted for use with other proteases and optimized. Switching to a more chemoselective reactive group could improve both probe reactivity and selectivity, and improving the fluorescence yield may allow for *in vivo*

localization studies of active proteasomes and a truly high-throughput, in-gel readout.

METHODS

Materials. Unless indicated otherwise, chemicals were obtained commercially of the highest-available grade.

Cell culture. The cell lines EL4 (mouse, thymoma) and MM.1S (human, multiple myeloma) were cultured in RPMI 1640 medium. HeLa cells (human, cervix epitheloid carcinoma) and U373 cells (human, astrocytoma) were cultured in DMEM. Both media were supplemented with 10% fetal calf serum (FCS), 100 units/ml penicillin and 100 μ g/ml streptomycin.

Synthesis and characterization of proteasome probes. Detailed information on synthetic procedures and characterization can be found in the **Supplementary Methods** online and **Supplementary Figures 2–4** online. Compounds **3** and **4** were tested and compared to other inhibitors as described previously¹⁶. U373 cells expressing an ubiquitin-Arg-GFP fusion protein were exposed to various concentrations of compounds **3**, **4**, AdaAhx₃L₃VS and various other proteasome inhibitors. After 5 h, we quantitated the accumulation of GFP by FACS. The specificity for proteasomal targets was determined by assaying competition for proteasome inhibition with compound **5**. We incubated equal amounts of protein, obtained by lysis of EL4 cells by glass beads (see below), with the indicated amounts (Fig. 3d) of inhibitor **3** or DMSO for 1 h at 37 °C, and then incubated with compound **5** for 2 h at 37 °C. Proteins were denatured by boiling in reducing sample buffer, separated by 12.5% SDS-PAGE and autoradiographed.

Probing proteasome activity in cell extracts. To obtain lysates for proteasome activity assays, we lysed cells with glass beads as previously described¹⁸. In brief, cells were washed twice with cold phosphate-buffered saline, pelleted and lysed with one volume of glass beads (<106 microns, acid-washed; Sigma) and an equal volume of homogenization buffer (50 mM Tris (pH 7.4), 1 mM dithiothreitol, 5 mM MgCl₂, 2 mM ATP and 250 mM sucrose) by vortexing at high speed for 15–30 min at 4 °C. Beads, membrane fractions, nuclei and cell debris were then removed from the supernatant by centrifugation at 16,000g for 5 min. We quantitated the protein content of extracts using the Bradford assay (BioRad) and then incubated equal amounts of protein with the indicated amounts (Fig. 4a) of probe **3**, 10 μ M control probe **4** or buffer for 1 h at 37 °C. Proteasome activity was assayed as described below.

Probing proteasome activity in living cells. We incubated 10×10^6 EL4 cells in 10 ml RPMI 1640 medium supplemented with 10% FCS and penicillin and streptomycin with the indicated amounts (Fig. 4c) of probe **3** or 10 μ M probe **4**, at 37 °C for 2 h. HeLa cells were incubated with 10 μ M probe **3** for the indicated time periods (**Supplementary Fig. 1**). Cells were collected by centrifugation. We lysed cells using glass beads and assayed protein content using the Bradford assay. Proteasome activity was assayed as described below.

In vivo profiling of bortezomib and MG132. We incubated 10×10^6 EL4 cells in 10 ml RPMI 1640 medium supplemented

with 10% FCS and penicillin and streptomycin with the indicated amounts of bortezomib (Fig. 5a) or MG132 (Fig. 5b) for 1 h at 37 °C. When indicated, this was followed by a 2-h incubation at 37 °C with 10 μM probe 3. We incubated 5×10^6 MM.1S cells in 5 ml RPMI 1640 medium supplemented with 10% FCS and penicillin and streptomycin with 20 nM bortezomib for the indicated time periods at 37 °C, followed by a 1-h incubation with 10 μM probe 3 at 37 °C. We lysed cells using glass beads or by incubating them for 30 min in NP-40 lysis buffer (50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% NP-40) and centrifuged the lysates for 5 min to remove membrane fractions, nuclei and cell debris. Proteasome activity was assayed as described below.

Assaying proteasome activity by immunoblotting. Equal amounts (typically 60 μg) of protein were denatured by boiling in reducing sample buffer, separated by 12.5% SDS-PAGE and electrotransferred onto polyvinylidene difluoride (PVDF) membranes. Immunoblotting was performed using a dansyl-sulfonamido-hexanoyl polyclonal antibody (1:7,500, rabbit; Molecular Probes) and horseradish peroxidase-coupled goat or swine anti-rabbit secondary antibody (Southern Biotech) followed by enhanced chemiluminescence (Western Lightning, Perkin-Elmer).

Assaying proteasome activity with fluorogenic substrates. Assays were performed as previously described²⁵ with minor changes. 10 μg of lysate was added to 100 μl of substrate buffer, containing 20 mM HEPES (pH 8.2), 0.5 mM EDTA, 1% DMSO, 1 mM ATP and 30 μM ZAlaArgArgAMC (tryptic-like), 60 μM ZGlyGly-LeuAMC (chymotryptic-like), 60 μM SucLeuLeuValTyrAMC (chymotryptic-like) or 60 μM ZLeuLeuGlu-βNA (caspase-like). SDS was omitted from the substrate buffer. Fluorescence was measured every minute for 25 min at 37 °C using a Fluostar Optima 96-well plate reader (BMG lab technologies) ($\lambda_{ex}/\lambda_{em} = 355/450$ nm for AMC and 320/405 nm for βNA) and the increase in fluorescence per minute was used to calculate specific activities of each sample. Cells were incubated with 1 μM epoxomicin (Affiniti) at 37 °C for 1 h to determine the nonspecific activity, which was subtracted from each measurement.

Note: Supplementary information is available on the Nature Methods website.

ACKNOWLEDGMENTS

The authors thank M.A. Leeuwenburgh for providing AdaAhx₃L₃VS. This work was financially supported by the National Institutes of Health (H.L.P. and P.J.G.), a grant from AstraZeneca (H.L.P. and P.J.G.), the Netherlands Organization for Scientific Research (H.O.), a National Cancer Institute SPORE grant Career Developmental Award (H.O.), the Dutch Cancer Society (H.O.), a Multiple Myeloma Research Foundation Senior Research Award (B.M.K.), the Dr. Saal van Zwanenberg Stichting (M.V.), the Austrian Academy of Sciences (E.F.), the Koninklijke Hollandsche Maatschappij der Wetenschappen (C.R.B.), and by the Stichting Fonds Doctor Catharine van Tussenbroek (C.R.B.).

COMPETING INTERESTS STATEMENT

The authors declare competing financial interests (see the Nature Methods website for details).

Received 13 October 2004; accepted 24 March 2005

Published online at <http://www.nature.com/naturemethods/>

- Hideshima, T. *et al.* The proteasome inhibitor PS-341 inhibits growth, induces apoptosis, and overcomes drug resistance in human multiple myeloma cells. *Cancer Res.* **61**, 3071–3076 (2001).
- Rock, K.L. & Goldberg, A.L. Degradation of cell proteins and the generation of MHC class I-presented peptides. *Annu. Rev. Immunol.* **17**, 739–797 (1999).
- Glickman, M.H. & Ciechanover, A. The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol. Rev.* **82**, 373–428 (2002).
- Kisselev, A.F. & Goldberg, A.L. Proteasome inhibitors: from research tools to drug candidates. *Chem. Biol.* **8**, 739–758 (2001).
- Adams, J. *et al.* Proteasome inhibitors: a novel class of potent and effective antitumor agents. *Cancer Res.* **59**, 2615–2622 (1999).
- Dahlmann, B., Ruppert, T., Kloetzel, P.M. & Kuehn, L. Subtypes of 20S proteasomes from skeletal muscle. *Biochimie* **83**, 295–299 (2001).
- Driscoll, J., Brown, M.G., Finley, D. & Monaco, J.J. MHC-linked LMP gene products specifically alter peptidase activities of the proteasome. *Nature* **365**, 262–264 (1993).
- Gaczynska, M., Rock, K.L. & Goldberg, A.L. Gamma-interferon and expression of MHC genes regulate peptide hydrolysis by proteasomes. *Nature* **365**, 264–267 (1993).
- Bogyo, M. *et al.* Covalent modification of the active site threonine of proteasomal beta subunits and the *Escherichia coli* homolog HslV by a new class of inhibitors. *Proc. Natl Acad. Sci. USA* **94**, 6629–6634 (1997).
- Greenbaum, D., Medzihradsky, K.F., Burlingame, A. & Bogyo, M. Epoxide electrophiles as activity-dependent cysteine protease profiling and discovery tools. *Chem. Biol.* **7**, 569–581 (2000).
- Liu, Y., Patricelli, M.P. & Cravatt, B.F. Activity-based protein profiling: the serine hydrolases. *Proc. Natl Acad. Sci. USA* **96**, 14694–14699 (1999).
- Adam, G.C., Cravatt, B.F. & Sorensen, E.J. Profiling the specific reactivity of the proteome with non-directed activity-based probes. *Chem. Biol.* **8**, 81–95 (2001).
- Adam, G.C., Sorensen, E.J. & Cravatt, B.F. Proteomic profiling of mechanistically distinct enzyme classes using a common chemotype. *Nat. Biotechnol.* **20**, 805–809 (2002).
- Kessler, B.M. *et al.* Extended peptide-based inhibitors efficiently target the proteasome and reveal overlapping specificities of the catalytic beta-subunits. *Chem. Biol.* **8**, 913–929 (2001).
- Ovaa, H. *et al.* Chemistry in living cells: detection of active proteasomes by a two-step labeling strategy. *Angew. Chem. Int. Edn. Engl.* **42**, 3626–3629 (2003).
- Dantuma, N.P., Lindsten, K., Glas, R., Jellne, M. & Masucci, M.G. Short-lived green fluorescent proteins for quantifying ubiquitin/proteasome-dependent proteolysis in living cells. *Nat. Biotechnol.* **18**, 538–543 (2000).
- Hideshima, T. *et al.* The proteasome inhibitor PS-341 inhibits growth, induces apoptosis, and overcomes drug resistance in human multiple myeloma cells. *Cancer Res.* **61**, 3071–3076 (2001).
- Bogyo, M., Shin, S., McMaster, J.S. & Ploegh, H.L. Substrate binding and sequence preference of the proteasome revealed by active-site-directed affinity probes. *Chem. Biol.* **5**, 307–320 (1998).
- Garcia-Echeverria, C. Recent advances in the identification and development of 20S proteasome inhibitors. *Mini Rev. Med. Chem.* **2**, 247–259 (2002).
- Lightcap, E.S. *et al.* Proteasome inhibition measurements: clinical application. *Clin. Chem.* **46**, 673–683 (2000).
- Stein, R.L., Melandri, F. & Dick, L. Kinetic characterization of the chymotryptic activity of the 20S proteasome. *Biochemistry* **35**, 3899–3908 (1996).
- Fiebigler, E., Story, C., Ploegh, H.L. & Tortorella, D. Visualization of the ER-to-cytosol dislocation reaction of a type I membrane protein. *EMBO J.* **21**, 1041–1053 (2002).
- Lindsten, K., Menendez-Benito, V., Masucci, M.G. & Dantuma, N.P. A transgenic mouse model of the ubiquitin/proteasome system. *Nat. Biotechnol.* **21**, 897–902 (2003).
- Luker, G.D., Pica, C.M., Song, J., Luker, K.E. & Piwnicka-Worms, D. Imaging 26S proteasome activity and inhibition in living mice. *Nat. Med.* **9**, 969–973 (2003).
- Elliott, P.J., Soucy, T.A., Pien, C.S., Adams, J. & Lightcap, E.S. Assays for proteasome inhibition. *Methods Mol. Med.* **85**, 163–172 (2003).