VITAL GUIDE SERIES

3 Infection control: managing the risks.

- How is infection spread in the surgery?
- What can be done to avoid contamination?
- Who is responsible for infection control in the dental practice?

3

VITAL GUIDE TO

Infection control

Welcome to the third article in the Vital Guide Series. At the end of the article are some CPD questions, which are designed to get you thinking about the article and to help you remember some of the key points. Here, **Martin Fulford**¹ explains the risks of infective organisms and how they are best managed by the dental team in the surgery environment.

Introduction

Infection control protocols are a risk management tool used to prevent the transfer of potentially infective organisms from one person to another. It is a very important professional responsibility to ensure the risk of acquiring infection from the dental surgery environment is reduced to the lowest level possible. All members of the dental team should be held accountable to ensure that appropriate protocols are introduced and that compliance to these protocols is adhered to at all times.

What are the risks?

There are a wide range of potentially infective organisms that can contaminate dental equipment and the surgery environment. Blood borne viruses such as Hepatitis B and C and HIV have often been highlighted as high risk. However, there are many other viruses that can be found in the upper respiratory tract, which includes the mouth, such as Herpes simplex, Herpes zoster, Influenza, Mumps, Epstein-Barr and many others. While

many of these viruses may cause minor ailments, some of them have the ability to cause or contribute to severe or life-threatening conditions in susceptible individuals.

There are also many bacteria that may live in the upper respiratory tract and can be a source of risk to individuals, such as tuberculosis (TB – the world's most common infectious disease), meningitis bacteria, syphilis, gonorrhoea and many others that may not be causing disease in the host but have the capability of causing disease in the susceptible individual. Some potentially harmful bacteria such as *Legionella* spp. may be found in the water from the dental unit and therefore will contaminate the dental environment.

More recently identified are very small protein particles known as prions, which may cause a fatal condition called variant Creutzfeldt-Jacob Disease (vCJD), and have been known to be transmitted via contaminated surgical instruments. This may be a particular hazard because they are not destroyed by regular disinfectants or autoclaving.

The infectious agents can be transmitted directly from person to person, via contaminated droplets in the form of aerosols or via contaminated objects such as instruments, work surfaces or contaminated hands.

¹ Dental Advisor to Somerset Health Authority
Email: martin.fulford@btinternet.com

Fig. 1 Eye protection is mandatory in infection control.

Who is at risk?

Any person coming into contact with contamination within the dental surgery is potentially at risk of acquiring an infection or becoming colonised by the organisms found within the contamination. It can be argued that surgery personnel are most at risk based on the frequency and duration of exposure, which is much greater than that, for example, of patients. Other people may also be at risk, such as engineers servicing dental equipment. A duty of care is also owed to dental technicians who may be handling impressions and dentures.

How are the risks transmitted?

The infectious agents can be transmitted directly from person to person, via contaminated droplets in the form of aerosols or via contaminated objects such as instruments, work surfaces or contaminated hands. Infections can be acquired by direct inoculation; for example, by the use of inadequately decontaminated instruments on patients, by sharps injuries to staff with contaminated instruments, by inhalation of infected droplets, by splashing of the eyes with infected droplets or by hand to mouth spread.

Management of risks

It is mandatory that all dental premises have an infection control policy document that sets out protocols that must be followed to manage the risks associated with surgery acquired infection. These protocols should be based on guidelines laid down by competent bodies such as the British Dental Association or the Department of Health.

Specific areas that should be covered by protocols are:

Staff health policies such as vaccination regimes and 'needle-stick' injury protocols; these can be advised by reference to local occupational health physicians.

Personal protective equipment policies: appropriate equipment such as eye protection, gloves, masks/visors and uniforms should be supplied to all personnel exposed to risks and to patients where appropriate. It is also important that this equipment is used appropriately and it is the responsibility of all parties to ensure that they are.

Disinfection and decontamination are a very important component of any infection control protocol and it is vital that this issue is dealt with explicitly. The basis of the protocol must be cleanliness. All infective agents have a minimum infective dose

and therefore all procedures must aim to remove as many organisms and as much bio-soiling as is possible. For example, work surfaces must be cleaned using an appropriate cleaning agent which will be applied using a vigorous scrubbing action with a disposable cloth/tissue. Dental instruments must be scrupulously clean before being subjected to sterilisation in an autoclave – these will only be effective if steam is able to condense on ALL surfaces of the equipment; any residual contamination may prevent this happening, leading to failure of the process. It is important to consider the risks to staff of the decontamination process itself; for example, manual scrubbing of dental instruments may pose a risk to the operator of a sharps injury or splashing of the eyes with contaminated material. The process itself should be effective, reproducible and validated, which may dictate the use of special equipment such as a washer/disinfector. All equipment used in the process should be maintained and tested to ensure that it is performing as it should do; good clinical governance principles dictate that not only should you use appropriate protocols but also that they must be demonstrated to be effective.

Fig. 2 Contaminated aerosols may be a source of infection.

Training

All staff exposed to cross infection risks must be trained in the practice protocols designed to manage these risks. This does not just apply to new staff: all staff should be updated frequently. Records should be kept of staff training. All infection control protocols should be revisited at regular intervals and if necessary should be updated in the light of new guidance or new knowledge in the field. This should be at a minimum of annually or more frequently if necessary; the subject lends itself well for regular practice meetings and clinical governance activity. It may often be the focus of critical incident analysis and much can be learned from such occurrences.

The risk of surgery-acquired infection rises day by day with the increase in prevalence of infectious diseases within the population, such as Hepatitis, HIV and TB. Dental staff owe it to themselves and their patients to adopt best practice on issues relating to infection control.

Useful websites

www.bda.org
www.dh.gov.uk

CPD Answers

Here are the answers to last issue's CPD questions on aesthetic dentistry.

1. Which of the following protocols is a method for avoiding cervical black lines?

- A Subgingival margin placement
- B Reinforcing oral hygiene procedures
- C Providing metal-ceramic restorations
- D Making maxillary teeth longer

Answer A

2. Youthful teeth are characterised by...?

- A Greater mandibular incisor display
- B Longer maxillary lip
- C Greater maxillary incisor display
- D Shorter mandibular lip

Answer C

3. Which of the following contributes to an artificial smile?

- A Bulbous teeth
- B Teeth with identical colour
- C Flat incisal embrasures
- D All of the above

Answer D

4. Which of the following is a culprit of a 'gummy' smile?

- A Maxillary gingival display less than 3 mm
- B Maxillary gingival display greater than 3 mm
- C A low maxillary lip line
- D Inflamed free gingival margins (FGM)

Answer B

Congratulations to the first three correct entries to be drawn from our summer issue.

They were: Mr Jeremiah Daly from Croydon, Mrs Stacy Long from Wolverhampton and Mrs Pamela Benson from Chester. They all win a copy of the book *Holistic oral care*.

the next issue of

vital

for the whole dental team

is out on November 25th

We've attracted the attention of the whole dental team – now you can! To be associated with this new, fresh and innovative magazine, call us today.

Contact Alison Tant at the
VITAL Advertising Department
British Dental Journal,
The Macmillan Building,
4-6 Crinan Street, London N1 9XW
Telephone: 020 7843 4833
Email: vital@nature.com

CPD Answer form

Please **PRINT** your details below:

Title

First Name

Last Name

Address

.....

.....

City

County

Postal Code

E-mail

Your Work

What is the name of your practice/
CDS/hospital?

.....

.....

What is your job title?

.....

Please **TICK (✓)** the answer to
each question

	A	B	C	D
Q.1	☐	☐	☐	☐

	A	B	C	D
Q.2	☐	☐	☐	☐

	A	B	C	D
Q.3	☐	☐	☐	☐

	A	B	C	D
Q.4	☐	☐	☐	☐

☐ If you prefer not to be contacted for purposes other
than this competition, please tick this box. We will
not pass your details on to any third parties.

Please send your completed
answer form to:
CPD Questions, *Vital* magazine,
BDJ, Nature Publishing Group,
4-6 Crinan Street,
London N1 9XW.

CPD Questions

In future, *Vital* will be offering verifiable CPD when it becomes mandatory for DCPs in 2008. In the meantime we have provided some CPD style questions for our new 'Vital Guide to...' series to encourage you to anticipate future CPD requirements. By way of extra encouragement we are pleased to offer *Management of medical emergencies for the dental team* to the senders of the first three correct entries to be drawn on 1 November 2006. The answers to the questions will be published in the next issue of *Vital*. After reading the article have a go at the following questions and send in your completed form to us. In the context of this article, only one of the answers to each of the following questions is correct.

1. Hepatitis is caused by:

- A A bacterium
- B A virus
- C A fungus
- D A prion

2. vCJD is caused by:

- A A bacterium
- B A virus
- C A fungus
- D A prion

3. Who is the most appropriate person to advise on vaccination regimes for dental staff?

- A Dentist
- B General medical practitioner
- C Occupational health physician
- D Maxillofacial surgeon

4. Which of the following is NOT an item of personal protective equipment in the dental surgery?

- A A visor
- B Latex gloves
- C Spectacles
- D An autoclave