The vertebral blocks, as well as the ribs, are the product of the primitive axial series of (invertebral) discs, which, when completely arrayed, each bear five branches, viz., two pair of hæmal arches, two pair of neural arches, and a fascicle of parallel cleets, so to speak, which being cemented together, both in the front and rear, by the superficial ossification of the discs at either end, are fused into the block pieces, as found, e.g. in the young hog; the cementing slab covering the big neural rib head likewise, and not only the pentagonal prismatic block. The first disciform ossification we find in the corals, forming cribrose ethmoidal discs, such as the closely set "sigillate impressions" of the *Astræa*, and afterwards left behind as the coccyx, e.g., of *Byathophyllum*.

SIEMENS' DYNAMO-ELECTRIC LIGHT*

A SERIES of experiments was made last week at Sheerness, with a view of ascertaining the applicability of Siemens' dynamo-electric light to torpedo services in time of war. This scientific combination is produced, as its name signifies, by the application of excessively rapid motion generated from the fly-wheel of a steam-engine to a very powerful set of ordinary galvanic "coils" in connection with soft-iron magnets. The leather strap from a four-horse power engine, encircling a small gun-metal pinion, causes it to revolve with the extreme velocity of 1,600 revolutions per minute, inducing motion in an electric "bobbin" at the side of an apparatus consisting of several sets of strong insulated coils. A stream of electricity consequently passes through them. This stream is conducted to a second series of coils, larger and more powerful than the first, which are also in combination with a pinion revolving 800 times per minute, thus intensifying the stream as it passes through them to a very considerable degree. Both negative and positive currents are now alternately given off from another "bobbin" at the side of the second series of magnetic coils, to the train of insulated wires, which conveys them to the position from which the dynamo-electric light is to be exhibited. Here there is a delicately contrived apparatus for containing the carbon points, between which the light is to be generated, adjusted at the top of a tripod somewhat similar in construction to that of a surveying instrument. At the back of the two carbon points, and "slotted" vertically to admit of their holders passing through it, is a concave reflector of white polished metal, which collects the rays of light into a focus, and transmits them in any required direction by means of an adjusting hand wheel below. A minute aperture in the centre of the reflector, precisely behind the junction of the two carbon points, throws a representation of the flame upon a piece of opal glass in a frame fixed at the back of the reflector; and through the agency of another small hand wheel which causes the carbon points to approach or recede from each other, the flame can be reduced or intensified at pleasure, by simply turning the wheel, care being taken at the same time to keep a watchful eye upon the picture produced, as the withdrawing of the points to too great a distance from each other will extinguish the light. It should have been remarked before that ample means are taken by lubricating the electrical apparatus to counteract the evil effects which might otherwise arise from the excessive friction consequent on the rapidity of motion in the several parts.

The object of instituting the series of experiments which were made on Monday was to ascertain if it was possible to throw such a stream of light upon an enemy's working parties engaged in interrupting communications with a line of torpedoes at night, as would render them sufficiently conspicuous to be fired at and consequently driven off. The place selected was the new fort at Garrison Point, Sheerness. The engine and "coils" were erected in the enclosure of the fort, while the instrument itself was placed in one of the massive embrasures piercing its sides. No sooner was steam got up and the order given to turn ahead, than the burring noise of the machine indicated that electricity was being rapidly generated, sparks and stars of vivid blue light being given off at the various joints. Another instant, and a vivid stream of light shot across the sea to a number of ships lying in the offing at a distance of about two miles, lighting them up with the brilliancy and distinctness of broad moonlight. The effect was magnificent. Clouds of mists, rendered visible by the intensity of the rays shooting through them, rolled across the broad field of bright light from time to time, not, however, interrupting the view in their progress. By shifting the direction of the rays laterally, each object in turn came within the compass

* Reprinted from the *Times*.

of the portion of horizon rendered clear. In fact, it was sufficiently apparent that no objects of any appreciable size, such for instance as an enemy's boats, could come within a mile or more of one of Siemens' dynamo-electric instruments in operation without being rendered conspicuous to any battery in the vicinity, and consequently involving to themselves the most imminent danger. Hence the result of the experiments may be pronounced a success; whether, however, a corresponding effect might not be obtained by a succession of parachute lights thrown from a rocket or mortar is quite an open question.

PHYSICS

Note on the Spectrum of the Aurora

ON the evening of November 9 there appeared one of the most magnificent crimson auroras ever seen at this place. When first observed, at about a quarter before six P.M., it consisted of a brilliant streamer shooting up from the north-western horizon; this was continued in a brilliant red, but rather nebulous mass of light, passing upward and to the north. Its highest points were from 30° to 40° in altitude. A white aurora, consisting of bright streamers, appeared simultaneously, and extended round to the north-east.*

The crimson aurora was examined with the spectroscope at six o'clock. The instrument used was a single glass-prism spectroscope, made by Dubosq, of Paris. On directing the slit toward the brilliant streamer above mentioned, a bright spectrum was observed consisting of five well-marked lines. A millimetre scale attached to the instrument was then illuminated with a gas flame, the auroral lines being readily measured, even when the numbers on the scale were bright enough to be read distinctly. The sodium line was used to adjust the scale, being equally divided by the division 100; the width of the slit was about one millimetre. As thus arranged, the five auroral lines, beginning at the red end, had the following positions:—Scale-Nos. 90, 110.5, 130, 138, 149. The brightness of the lines was, following the above order, 3, 1, 5, 2, 4, the second line from the red end of the spectrum being the brightest. The line marked 90 and the one marked 110.5 were sharp and well defined; the others were all nebulous on the edges. Before the measurements were completely verified by a second comparison, the crimson aurora entirely vanished, having endured less than half an hour. In the white aurora which remained, the spectroscope showed four of the five lines given; the crimson line alone was absent. The measurements are exact to half a division of the scale.

To determine the approximate wave-lengths of these lines, comparison was made both with certain metallic lines and with the lines of the solar spectrum. On the scale of this instrument the metallic lines employed read as follows:—

Ka 63, Lia 79, Srβ 80, H(c) 82, Caa 91, Sra 96, Caβ 113, H(f) 146.5, Srδ 163, Csβ 165, Csa 167, Rba, & β 200, Kβ 218.

The Fraunhofer lines measured as follows:—

a 70.5, B 76, C 82, D 100, E 124.5, δ 130, F 146.5, G 189.

Direct interpolation was used in comparing the wave-lengths of the auroral lines with those given above, the wave-lengths of the Fraunhofer and elemental lines being taken from Gibbs's tables (*Amer. Jour. of Science and Arts*, II. xliii. 1; xlvii. 194). This method was believed capable of giving results as close as the instrumental measurements. In this way the wave-lengths of the five auroral lines were obtained, as given in the following table:

Line.	Scale number.	Wave-length.	Auroral lines.	Other measurements.
B	76	687		
C	82	656		
(1)	90	623	623	627 Zöllner.
D	100	589		
(2)	110.5	562	562	557 Angström.
E	124.5	527		
(3)	130	517	517	520 Winlock.
b	130	517		
(4)	138	502	502	
F	146.5	486		
(5)	149	482	482	485 Alvan Clark, Jr.
G	189	431		

* Professor Newton informs me that he observed an equally brilliant red patch of auroral light in the north-east, five or ten minutes earlier. Since the lower end of the red streamers was much lower than that of the white, it would seem as if the red were seen through the white, the red being most remote.

In this table, column 1 gives the auroral and the Fraunhofer lines; column 2, the number of these as measured upon the scale of the spectroscope used; column 3, the wave-lengths of these lines obtained as above stated; column 4, the wave-lengths of the auroral lines, given by themselves; and column 5, the wave-lengths of what I assume to be the same lines, with their wave-lengths as measured by the observers mentioned.

The point of particular interest in this observation is the fact that the line (4) of wave-length 502 is not laid down in any authority accessible to me as having been observed in the auroral spectrum. Indeed, no previous observer, so far as I know, has seen any auroral line between the Fraunhofer lines *b* and *F*. Professor C. A. Young (*Journal of Science and Art*, III. ii. 332, Nov., 1871) gives two lines—Nos. 56 and 57, or 1866·8 and 1870·3 of Kirchhoff—observed by him in the sun's chromosphere and also by Rayet in the eclipse of 1868, one of which may coincide with this fourth auroral line.

New Haven, Nov. 13

GEORGE F. BARKER

SCIENTIFIC SERIALS

THE *Geological Magazine*, Nos. 86–89, August to November 1871. This valuable magazine continues to furnish us every month with important and interesting articles upon subjects belonging to the various departments of geology. In the first number now before us we find an interesting paper on volcanoes by the editor, Mr. H. Woodward, and a particularly valuable article by Mr. J. W. Judd on the anomalous mode of growth of certain fossil oysters. In the latter, which is illustrated with a plate, the author notices those oysters from various secondary deposits, in which the shell has acquired throughout the peculiar sculpture of some ammonite, *Trigonia*, or other shell, to which its lower valve has adhered during growth.—In the September number the most interesting paper is Mr. Woodward's description of a new Arachnide from the Dudley coal-measures. This animal, to which the author gives the name of *Eophrynus Prestvicii*, is most nearly allied to the existing genus *Phrynus*, and the specimen is remarkable for the beautiful preservation of the casts of both surfaces.—Among the contents of the October number we must call particular attention to Dr. Murie's article on *Sivatherium*, in which the author discusses the characters of that most remarkable animal, which he regards as most nearly allied to the Saiga antelope, the latter being placed by him at the central point of ramification of the hollow-horned ruminants, and leading from the ruminants towards the Pachyderms through the Tapir. This valuable memoir is illustrated with two plates, one representing the skeleton of the animal, the other giving an ideal restoration of the living aspect of the male, female, and young of this gigantic ruminant.—The November number opens with a biographical notice (with a portrait) of Sir Roderick Murchison, followed by a shorter one of Mr. Charles Babbage. The other articles contained in it are on relics of the Carboniferous and other old land-surfaces, by Mr. Woodward; on the prospects of coal south of the Mendips, by Messrs. Bristow and H. B. Woodward; on the futile search for coal near Northampton, by Mr. S. Sharp; and on the Foraminifera of the Cretaceous rocks, by Messrs. T. Rupert Jones and W. K. Parker.

THE *Journal of Botany* for November commences with an interesting contribution to historical botany; in a paper read by the late Robert Brown before the Edinburgh Natural History Society in 1792 on "The Botanical History of Angus" never before printed. It was probably his first contribution to botanical science, having been written when he was about eighteen years old. Prof. Thistelton-Dyer contributes some observations on "Fungi parasitic upon *Vaccinium Vitis-Idaea*," and Mr. A. W. Bennett "Further observations on Protandry and Protogyny," in continuation of his previous researches on this subject. Mr. T. A. Briggs has a note on an undescribed species of *Rubus*, and the remainder of the number is filled up with short notes, abstracts, extracts, and reviews.

The number for December opens with the commencement of a paper by Mr. J. G. Baker "On the Botany of the Lizard Peninsula." Although this district is well known to botanists as the habitat of many very rare and local plants, yet no detailed account has yet been published of the flora of this portion of Cornwall. From the idea that many plants very common in other parts of England would find their limit short of this south-

western extremity of the island, a list is here given of every flowering plant observed during a three days' visit, accompanied by general remarks on the peculiarities of the flora, both in what it includes and in what is absent from it. The only other original paper of importance in the number is a new arrangement by the Rev. J. C. Leeke of the English species of the extremely difficult genus *Salix*.

Journal of the Royal Geological Society of Ireland. Part I, vol. iii. new series (vol. xiii.), has just been published. It contains besides the Report of Council for 1870–71, J. Emerson Reynolds on two remarkable Crystals of Galena; G. H. Kinahan, additional notes on Foliation, and supplementary notes on some of the Drift in Ireland; R. G. Symes, on the Geology and extinct Volcanoes of Clermont, Auvergne—plates i. ii. iii.; W. H. Baily, on the genus *Pleurorhynchus*, and a new species—plate iv.; M. H. Ormsby, Analyses of some Granite Rocks from India, and of their constituent minerals (1668); Edwd. T. Hardman, Analysis of Trachyte Porphyry from Tardree near Antrim, and on the Analysis of a Limestone compared with that of the same rock where it is in close proximity to a Doloritic Dyke; R. C. Tichborne, note on the Geological Formation of some of the Tiroxides.

Journal of the Chemical Society, October.—This number does not contain any papers originally communicated to the Society. The abstracts of foreign papers, however, occupy nearly 100 pages, and comprise many subjects of interest. The importance of the work done this way by the Chemical Society can scarcely be estimated; the journal must now be of great value not only to the chemist, but also to the physicist, physiologist, and to the chemical manufacturer, for many papers in these subjects are abstracted fully. An abstract of M. Berthelot's paper on the heat evolved in the formation of organic derivatives of nitric acid is very interesting. It is shown that in the formation of nitro-glycerine, a very small amount of heat is evolved, as compared with that evolved in the formation of gun-cotton or nitrobenzene. This will account for the ready decomposition of the former, and the formidable effects produced by its decomposition. Amagat has experimented on the compressibility and dilatation of sulphurous and carbonic anhydrides; he finds that they first begin to act as perfect gases at a temperature of 250° C. Several of the abstracts contained in this number have already been noticed in these pages. One of them deserves especial notice, by Friedel and Ladenburg, on silicopropionic acid; this body is the first in which the group CO₂H contained in organic acids has been replaced by the corresponding group SiO₂H. Amato has obtained a curious compound, glucosophosphoric acid, the sodium salt of which has the composition C₆H₁₁O₅Na₂PO₄. Waage has published a paper on the use of bromine in chemical analysis, in which he points out that this reagent can be substituted with advantage for chlorine in many instances; it is very useful in decomposing pyrites, the whole of the sulphur being easily oxidised. We find an abstract of Bischof's paper on Fire Clays, which appears to deal very practically with this most important subject.

THE part just issued of the *Transactions of the Linnean Society*, completing vol. xxvii., contains three papers, of two of these, "Revision of the Genus *Cassia*," by Mr. G. Bentham, the president, and "Contributions to the Natural History of the *Passiflora*," by Dr. M. T. Masters, abstracts have already appeared in our columns. The remaining paper, "Notes on the Reptiles, Amphibia, Fishes, Mollusca, and Cretacea, obtained during the Voyage of H.M.S. *Nassau* in the years 1866–69," by Dr. Cunningham, contains descriptions of several new species collected and named by the naturalist to the expedition, and notes on the habits, localities, and characters of many other species. All these papers are illustrated by plates.

The first part of Volume xxviii. is also published, consisting of only a single paper, Dr. Triana's monograph of the *Melastomaceae*. After some general remarks on the order, and on each of the genera comprised within it in French, follows an enumeration of the species, with the synonymy, references to type specimens in the principal herbaria, and fresh descriptions of new or badly-described species. It is illustrated by seven plates.

THE *Bulletin of the Royal Academy of Sciences of Belgium* for September and October, 1871 (Tom. xxxii., Nos. 9 and 10), contains but little scientific matter.—M. J. C. Houzeau communicates a description of a method of measuring directly, the