upon a surface of calm water, they are, in part, reflected according to the usual law, and a rainbow is then seen by reflection. But the absolute position of the bow changes with every change in the position of the observer's eye; hence the bow seen mirrored in the pool is *not* the reflection of that seen at the same time directly in the heavens. Suppose the shower to be fixed in space, then the drops which produce the bow seen directly, would not be those which produce the bow as seen by reflection.

In the paragraph to which your correspondent "Z.X.Y." has called attention, I meant to combat the notion, entertained by many, that the rainbow is reflected after the fashion of an ordinary floating cloud which emits light in all directions, and which, by the light thus emitted, paints its image in the water. A few additional words might have made my meaning clearer; but as I was dealing at the time more with historic statement than with scientific exposition, I desired to be brief. I can hardly think, however, that your correspondent will be angry with me for giving him what must have been agreeable as well as successful occupation at the Falls of the Rhine.

Royal Institution, Sept. 15

JOHN TYNDALL

Original Research at the Universities

My attention has been arrested by the following sentence in the extract given by you from Prof. Frankland's evidence before the Science Commission:—"I believe that one cause (of the slow progress of original research in England) lies in the entire non-recognition of original research by any of our Universities. Even the University of London, which has been foremost in advancing instruction in experimental science, gives its highest degree in Science without requiring any proof that the candidate possesses the faculty of original research, or is competent to extend the boundaries of the science in which he graduates."

It may interest Dr. Frankland and those who take the same view as he does, to know that this subject has engaged the attention of the graduates of the University of London. At a meeting of the Annual Committee of Convocation in December last, it was moved by Prof. Guthrie—

"That every candidate for the degree of Doctor of Science shall be required to submit to his respective Examiners a written dissertation embodying some original research in one or more of the subjects of his intended examination; and that such dissertation be approved before the candidate be allowed to proceed to examination."

This motion I had the honour of seconding; but the degree of acceptance which the principle involved in it met with from the Committee is seen by the sequel, as stated in the printed minutes, that it was "rejected by a large majority." The exact numbers, if my memory serves me rightly, were Ayes, 3; Noes, 16; among the Noes were two Doctors and one Bachelor of Science, and at least five Doctors of Medicine. The "Annual Committee," it may be stated, is a representative body elected annually by the graduates in Convocation, but has no legislative or administrative power, this resting entirely with the Senate.

ALFRED W. BENNETT

Endowment of Research

WITH regard to the Endowment of Scientific Research, could not this be well placed in the hands (as it now is, to a very limited extent) of a Committee of the British Association? the committee being authorised to supply funds for experimental purposes, and the members, say three or four in number, to have a permanent salary for the time spent in the examination of claims from applicants.

It might possibly be desirable that one or more of the committee should retire every two or three years and not be eligible for re-election until after the lapse of three years; and also, to prevent waste of time, that all applications for help should be presented only through one or more gentlemen of known scientific attainments, and not of necessity at the instigation of the person to whom the assistance was to be rendered. I believe that this would be a good practical arrangement as regards the poorer class, who are compelled to throw up valuable original researches to supply themselves and those depending on them with homes and food.

The abuse of a trust of this kind would hardly be possible, as

the help would of necessity be given in those cases where a certain amount of work had already been done under difficulties, and where the natural instinct for original research was of necessity strongly developed. The presentation of an annual sum for, say five years, renewable at the end of the time if necessary, would be a godsend to many a man who has allowed himself to starve for the benefit of posterity.

THOS. FLETCHER

FERTILISATION OF FLOWERS BY INSECTS* III.

On the co-existence of two forms of flowers in the same species or genus,—a more conspicuous one adapted to cross-fertilisation by insects, and a less conspicuous one adapted to self-fertilisation.

SINCE Darwin, in his admirable work on Orchids,† had proved that the flowers of this family are endowed with an immense variety of contrivances for cross-fertilisation by insects, it was almost generally admitted by botanists that cross-fertilisation is the rule throughout the whole vegetable kingdom. Darwin's well-known aphorism, that "Nature abhors perpetual self-fertilisation" was exaggerated by his successors in this field of research, Hildebrand in Germany and Delpino in Italy, who, in their various elaborate memoirs on the fertilisation of flowers, repeatedly expressed their strong belief that nature abhors self-fertilisation at all. In direct opposition to this opinion, Axell‡ propounded the doctrine that the development of the fertilising arrangements in phanerogams has been always an advance, and still continues to advance, in one and the same direction, towards a perfection which affords more and more facilities for self-fertilisation.

My own observations on the contrivances of our flowers and on the insects really visiting and fertilising them, have convinced me, that neither Hildebrand's and Delpino's, nor Axell's opinion is a thoroughly adequate one, but that under certain conditions the facility for self-fertilisation is most advantageous to a plant, while, under other conditions, the inevitableness of cross-fertilisation by the visits of insects is the more advantageous.

To all plants the flowers of which possess such a degree of attractiveness for insects that cross-fertilisation by these transporters of pollen is never wanting, the possibility of self-fertilisation is quite useless, and from this cause, not being subjected to the effects of natural selection, may be lost, like any useless peculiarity, and in many instances, indeed, has been lost. On the contrary, to those plants the flowers of which possess so slight a degree of attractiveness for insects, that the transportation of the pollen to the stigma by insects is effected in but very few cases, the possibility of self-fertilisation is most advantageous, and indeed we find in most cases such plants well adapted for self-fertilisation.

Among many facts which I could appeal to as proofs of my statements, there are, I believe, none more instructive than those alluded to in the superscription of this article.

In some species of our wild plants I have found on different plants two different forms of flowers, evidently showing the connection above stated between attractiveness for insects and adaptation for inter-crossing or for self-fertilisation. As nobody before, for aught I know, has observed this phenomenon, I will give some details of the most important instances hitherto observed.

Lysimachia vulgaris

Of this species specimens with more conspicuous flowers are found in sunny localities. The petals of this form are dark yellow with red at the base, on an average about 12 mm. long, and 6 mm. wide, opening widely and

* Continued from p. 206.

† "On the Various Contrivances by which British and Foreign Orchids are Fertilised by Insects." (London, 1863.)

‡ In his work: "Om anordningarna för växtens växtens befruktning." (Stockholm, 1869.)

bending outwards and backwards; the filaments are red-coloured towards their end; the style overtops the longest stamens by some millimetres. A species of bee, *Macropis labiata* Pz., frequently visits these flowers for pollen. It comes first into contact with the stigma, and supplies it with pollen from previously visited flowers, thus regularly effecting cross-fertilisation. But if we prevent the visits of insects by covering over the stems by a

FIG. 9.—*Euphrasia officinalis*. Lateral view of a flower of the largest form just opened.

FIG. 10.—Position of the stigma (*st*), and of the anthers (*a¹*, *a²*) of the same flower in a more advanced state.

FIG. 11.—Two anthers, seen from the inner side, showing the slits fringed with hairs.

net, self-fertilisation scarcely takes place, in consequence of the style overtopping all the stamens.

Specimens of the same species with less conspicuous flowers are found in shady ditches. The petals of these plants are lighter yellow, uniform in colour, without any red at the base, on an average 10 mm. long, and 5 mm. wide; they only open slightly, remaining nearly upright,

FIG. 12.—Lateral view of a flower of the smallest form, just opening.

FIG. 13.—Position of the stigma (*st*) and the anthers (*a¹*, *a²*) in this flower.

FIG. 14.—Front view of a flower of the same form, in a more advanced state.

(All the figures are magnified in the proportion of 7:1. The hairs of the calyx and the coloured spots and lines of the corolla are omitted.)

but diverging obliquely; the filaments are greenish yellow, without any red towards their end; the style hardly equals the two lowest and longest stamens. The stigma comes without any external agency into contact with the pollen of the same flower thus regularly experiencing self-fertilisation. This manner of producing seeds is an indispensable condition for preservation of

this variety of *Lysimachia vulgaris*. For in consequence of its shady habitat, and of its lower degree of attractiveness for insects, its flowers are but very rarely visited, and it would be exposed to extinction without the possibility of propagation by self-fertilisation. I but once observed a little fly of the family of Syrphidæ, *Syrphia pipiens* L., eating the pollen of this shady form of flowers. Although this fly might possibly transport pollen from one flower to the stigma of another, cross-fertilisation was nevertheless by no means more probable than self-fertilisation.

The two forms here described of the flowers of *Lysimachia vulgaris* graduate into each other by connecting forms, which are met with in intermediate localities, for instance on the sunny edges of ditches.

Another example of the same sort of dimorphism, even more striking than that just mentioned, is presented by *Euphrasia officinalis*. Of this species flowers are found in different localities of a very different size. But the more the attractiveness for insects is increased by the size of the corolla, the more is cross-fertilisation secured in case insects visit the flowers, self-fertilisation at the same time being prevented; while on the contrary, the smallest flowers regularly fertilise themselves, even without the visits of insects. I will attempt to explain these peculiarities by drawings of the largest and of the smallest form of flowers I have hitherto been able to find.

In the flower just opened of the largest form (as shown in Fig. 9), the stigma, already in a mature condition, greatly overtops the anthers. Therefore an insect,* inserting its proboscis into the tubular corolla in order to gain the nectar contained at the bottom of its tube, first grazes the stigma charging it with pollen-grains from flowers previously visited, and then pushes against the two hairs (*sp*) which project from the two lower anthers (*a²*) into the middle of the entrance to the corolla. This shaking of the hairs is transmitted to all the four anthers, which lie close together and are soldered together by their upper margins, and a small quantity of the smooth powdery pollen-grains falls out of all the pollen sacs. The slits in the pollen sacs being fringed with hairs directed downwards (as shown in Fig. 11) a lateral dispersion of the pollen-grains is prevented; all the pollen-grains shaken out fall directly downwards upon the proboscis, enabling it to fertilise the next flower visited by the insect.

In the state just described the corolla has not yet attained its full size. Growing farther, it at length equals the stigma by which it was at first so much overtopped, and now the mutual position of the stigma and the anthers is that shown in Fig. 10. When occupying this position, the stigma is always already shrivelled and brownish coloured, and is no longer capable of being fertilised. Self-fertilisation is therefore quite impossible.

The probability of cross-fertilisation and of self-fertilisation is directly opposite in the flowers of the smallest form, presented by Fig. 12–14. Whilst in the flowers of the largest form, as just described, the anthers remain soldered together, and do not scatter their pollen unless the hairs are shaken, in the flowers of the smallest form the anthers separate from each other, and scatter nearly all their pollen long before the corolla has fully opened. The end of the style, moreover, bends inwards so much as to bring the stigma (as Fig. 13 shows) close beneath the upper anthers. Therefore, on examining a flower hardly half-opened (Fig. 12), we always find the stigma already largely charged with pollen-grains of the same flower. When fully opened, the flowers of the smallest form show the stigma in a shrivelled and brownish coloured condition, lying between the separated and emptied pollen-sacs (as shown in Fig. 14). Hence cross-fertilisation could scarcely be effected, even if insects (which I never

* I observed four species of bees and three species of Diptera visiting the flowers of *Euphrasia officinalis* for honey.

observed) should visit these very inconspicuous flowers. The fringing hairs in the flowers of the largest form, so nicely securing the perpendicular falling of the pollen-grains upon the proboscis, are quite useless in a flower regularly restricted to self-fertilisation; indeed, in the anthers of the smallest form we find no fringing hairs at all, or only a few isolated ones.

The two extreme forms here described graduate into each other by various intermediate forms. When publishing my book on "Fertilisation of Flowers by Insects," I had never observed either the largest or the smallest form here described. From this cause the figures in page 291 of my work, drawn from other varieties, differ in some points from the description here given.

In *Lysimachia vulgaris* the two forms here described are so closely allied, that no botanist, for aught I know, has considered them worthy of being distinguished as varieties by separate names; in *Euphrasia officinalis* the difference between the two forms is somewhat greater, and some botanists, although overlooking the different manner of fertilisation, have distinguished them as varieties, (for instance, Ascherson in his "Flora der Provinz Brandenburg").

In a third example of the same dimorphism of flowers, presented by *Rhinanthus crista galli*, the divergence of the two forms has proceeded so far that most botanists distinguish them by separate names, some as varieties (*Rh. crista galli* α and β of Linnæus), others as distinct species (*Rh. major* Ehrh. and *Rh. minor* Ehrh.). These two forms differ with respect to their fertilisation, nearly in the same manner as the largest and the smallest form of *Euphrasia officinalis*. *Rh. minor* having a smaller corolla, and therefore being but rarely visited by insects, regularly fertilises itself when insects do not visit it, by bending the stigma beneath the pollen-sac, which at last opens spontaneously, and covers the stigma with its pollen-grains. In *Rh. major* the stigma so far overtops the pollen-sac that self-fertilisation is excluded. It is, however, a remarkable difference between *Rh. minor* and the smallest form of *Euphrasia officinalis*, that the former is regularly cross-fertilised, when visited by insects, if this happens not too late, and that it only has recourse to self-fertilisation if altogether unvisited by insects.*

Lippstadt, Sept. 9

HERMANN MÜLLER

THE 'POLARIS' ARCTIC EXPEDITION

THE missing link in the story of the *Polaris* Expedition has been picked up, and the narrative, as a whole, is one of the strangest in the whole history of Arctic adventure. Our readers may remember the story we gave of the 19 persons who were left on the ice-floe when the *Polaris* broke from her moorings in about N. lat. 79°, on the night of October 19, 1872, and who were all miraculously rescued six months later off the coast of Labrador. Eleven more of the crew arrived at Dundee last Friday afternoon in the whaling vessel, *Arctic*, Capt. Adams. Among these eleven are, Capt. S. O. Buddington, sailing and ice master, Dr. Emil Bessels, H. C. Chester, first mate, W. Martin, second mate, Emil Schumann, chief engineer, A. Odell, second engineer, besides a fireman, the carpenter, and three seamen.

After the ship drifted away from the floe she ultimately reached Lifeboat Cove, where it was resolved to beach her, which was done after much trouble. From the timbers of the ship a house was constructed on shore, and by the help of a few friendly Esquimaux, and the provisions and coals saved from the *Polaris*, the fourteen men spent the winter much more comfortably than might have been expected under the circumstances. Towards the end of the winter, however, it was resolved to make an

attempt to push southwards, and for this purpose under the superintendence of the energetic first mate, Mr. Chester, of whom all the crew speak in high terms, two boats were, amid many hardships, constructed out of some of the cabin-timbers of the *Polaris*. About the middle of last June, the boats having been completed and packed with what provision could be had, as well as ammunition, the party bade adieu to Lifeboat Cove and proceeded to make their way southwards. After many anxieties Cape York was reached on June 21. Here the boats were quite beset among the ice, but the greatest possible excitement and fear were experienced when, on the 23rd, a vessel was espied. She turned out to be the *Ravenscraig* whaler, of Dundee, Capt. Allan. All hands determined to reach the ship with the least possible delay, but in doing so they were greatly assisted by Capt. Allan, who had sent his crew to help them in carrying what things they had in their possession. They brought one boat with them and left the other. On reaching the ship they were very kindly treated, but subsequently, so that the fishing operations might be interrupted as little as possible, Capt. Allan shipped a few on the *Arctic*. The latter vessel having completed her fishing earlier than expected, and knowing that the crew of the *Polaris* would be anxious to return home as speedily as possible, Captain Adams, her commander, went in search of the *Ravenscraig*. Finding her, he took on board those of the survivors it contained, but Capt. Allan had previously put on board the *Intrepid*—R. W. D. Bryan, astronomer and chaplain; J. B. March, seaman; and John W. Booth, fireman. The *Intrepid* is expected in the course of a few weeks. The men state that the privations which they suffered were by no means of a serious character. The life was rough, laborious, and monotonous, and although danger occasionally presented itself in a way well calculated to inspire the greatest fear, yet no accident of any importance occurred to the adventurers.

Capt. Markham, R.N., accompanied Capt. Adams, of the *Arctic*, on his whaling voyage with the view of making investigations in the northern regions. The captain left Dundee on Friday, and was present in the Geographical Section at Bradford on Saturday, where he was received with great enthusiasm, and where he announced himself as heart and soul a convert to the Smith Sound route to the Pole.

The men connected with the *Polaris* Arctic expedition left Dundee on Monday, and Liverpool on Tuesday, for New York. All were in excellent health and spirits, and some of them say that they would have no objection to go on another such enterprise. Capt. Buddington states that Capt. Hall was buried in lat. 81°38' N., and long. 61°44' W. The vice-consul examined the crew of the *Polaris* on Monday, and transmitted their depositions to America, so that their statements may be extant should any accident befall themselves.

Dr. Bessel, who was the chief of the scientific party connected with the expedition, states that zoological, meteorological, botanical, and geological specimens were collected, but many of them were lost when the crew separated in October last. Careful and minute observations were also made, and after the explorers were picked up by the *Ravenscraig* they were continued. These surveys, of course, were not so exact as was to be desired, there being little convenience and very few instruments. The specimens taken on board the whalers are all preserved, and it is believed that, from a scientific point of view, they will be of very great value. The opinion of Dr. Bessel is that, had no accident occurred to the *Polaris*, the expedition would have been prosecuted. Regarding statements which had been made respecting the causes which led to the death of Captain Hall, he asserts that the captain was carried off by an attack of apoplexy. The doctor declines to enter into the question as to the management of the expedition after

* A further explanation of these two forms is given in my book "Die Befruchtung der Blumen durch Insekten," pp. 294-296.