

foreign material filling the fissure reflecting light less perfectly than the general surface of the berg. I conceive that the upper surface of one of these great tabular southern icebergs, including by far the greater part of its bulk, and culminating in the portion exposed above the surface of the sea, was formed by the piling up of successive layers of snow during the period, amounting perhaps to several centuries, during which the ice-cap was slowly forcing itself over the low land and out to sea over a long extent of gentle slope, until it reached a depth considerably above 200 fathoms, when the lower specific weight of the ice caused an upward strain which at length overcame the cohesion of the mass, and portions were rent off and floated away. If this be the true history of the formation of these icebergs, the absence of all land débris in the portion exposed above the surface of the sea is readily understood. If any such exist, it must be confined to the lower part of the berg, to that part which has at one time or other moved on the floor of the ice-cap.

The icebergs, when they are first dispersed, float in from 200 to 250 fathoms. When, therefore, they have been drifted to latitudes of 65° or 64° S., the bottom of the berg just reaches the layer at which the temperature of the water is distinctly rising, and it is rapidly melted, and the mud and pebbles with which it is more or less charged are precipitated. That this precipitation takes place all over the area where the icebergs are breaking up constantly, and to a considerable extent, is evident from the fact of the soundings being entirely composed of such deposits; for the diatoms, Globigerina, and radiolarians are present on the surface in large numbers; and unless the deposit from the ice were abundant it would soon be covered and masked by a layer of the exuviae of surface organisms.

SCIENTIFIC SERIALS

Poggendorff's Annalen der Physik und Chemie, No. 6.—The first paper, by G. Lundquist, On the reflection of light from the outside of isotropic bodies, is reprinted from the "Transactions" of the Royal Society of Upsala.—Dr. H. Brongersma contributes a memoir On the Medium in electrical influence, in which, following up the experiments of Riess, he examines Faraday's theory of electrical induction by polarisation of the medium, and describes in detail the apparatus with which he experimented, tabulates his results, and agrees with Riess's induction as to a direct influx of electricity.—The next article comprises a series of communications from the Mineralogical Institute of the University of Strasburg, in which Paul Groth treats of the crystalline form and thermo-electric properties of smaltine or arsenical cobalt. Its chemical formula is very variable, $R = (\text{Co}, \text{Ni}, \text{Fe}) \text{As}_2$. Eight samples from different localities, which contribute a better knowledge of the hemihedral forms of the species, are discussed. He concludes that some of the forms hitherto regarded as holohedral are hemihedral forms with parallel sides. As with iron pyrites and cobalt-glance, with which it is isomorphous, one part of the crystal is negative towards copper, while the other half is positive.—Dr. Hintze treats of the chemical composition of leadhillite. Prof. Laspeyres two years ago described a mineral from Iglesias, in Sardinia, under the name of maxite, of which the formula was $5\text{PbSO}_4 + 9\text{PbCO}_3 + 4\text{PbO} + 5\text{H}_2\text{O}$. M. Bertrand, of Paris, soon after published an account of leadhillite from the same neighbourhood. The author proves that the two minerals are identical, and that the formula of leadhillite is not $\text{PbSO}_4 + 3\text{PbCO}_3$, as hitherto believed, but $2\text{PbSO}_4 + 4\text{PbCO}_3 + \text{PbO} + 2\text{H}_2\text{O}$. The next paper, by the same author, is crystallographic researches on the combination of aldehyde with the aromatic hydrocarbons. The chemical composition and crystalline forms are given of ditolyltrichloroethan, diphenyltribromoethan, diphenyltrichloroethylen, diphenyldibromoethan, dimonobromophenyltrichloroethan, &c. All these bodies belong to the monoclinic system.—Dr. A. Arzruni gives a short note on twin-growth in willemite. The next paper by the same author is "Optical researches on the turpentinehydrates," which he follows by crystallographic and optical researches on compounds of urea. His last paper is on two isomorphous substances derived from benzol.—F. Zollner prints his important paper on the aggregation and position of the sun-spots, and concludes that they are cooled scoraceous products.—Dr. Karl Braun contributes studies on the earth's magnetism.—Among the reprinted papers are Dr. Andrews' on ozone and Prof. Wright's on the polarisation of the zodiacal light.

Memorie della Societa Spettroscopisti Italiani, September 1874.—This number contains a paper by Mr. J. N. Lockyer, describing certain phenomena seen when examining the spectrum of the electric light through a mass of sodium vapour in a tube. When this is done, the sodium lines are seen to shade gradually off, sometimes on one side, sometimes on both, the boundary of the shading being curved and sometimes limited by a bright line.—There is also another paper by the same author, On experiments on the absorption of a great thickness of sodium and iodine vapour in a tube 5 ft. long. After mentioning that it had been hitherto assumed that a great thickness of gas causes its radiation, and therefore its absorption, to become more continuous, he states that, on generalising his work, it appears that when the density of a vapour is increased, a continuous spectrum is approached in the case of the metallic elements of low specific gravity by the widening of their lines, and in that of the elements of high sp. gr. by the increase of the number of lines. To test this, the absorption of sodium vapour in a 5 ft. tube was observed, and the D line was found to be no thicker than the same line produced by a test-tube full of the vapour, and the line was thicker than the D-line in the solar spectrum, in which spectrum all the short lines are reversed.—Father Secchi communicates a letter of A. T. Arcimis, detailing observations on the spectra of meteorites. The spectra of all seem to be continuous, but wanting in the violet, that colour of the spectrum predominating according to its colour to the naked eye. The sodium line was visible in the trail of some, as also were the lines of magnesium.—G. De Sisa gives a table of the solar spots observed at Palermo from June to September.—A table of the chromosphere, as seen during February and March last at Palermo, is added to this number.—E. Fergola contributes a lengthy paper on the position of the axis of rotation of the earth with respect to its axis of figure.

SOCIETIES AND ACADEMIES

LONDON

Linnean Society, Dec. 3.—Dr. G. J. Allman, F.R.S., president, in the chair.—Mr. Jas. Brogden, Sir Edmund Buckley, Bart., M.P., Messrs. Jas. Cowherd, P. Duffy, C. C. Dupré, A. M. Ross, and J. W. Silver were severally elected Fellows of the Society.—Prof. Huxley read a paper On the classification of the animal kingdom, which will be found in another column. An interesting discussion followed, in which the President, Mr. Busk, Mr. H. G. Seeley, Mr. Stewart, Dr. Murie, and others took part.

Chemical Society, Dec. 3.—Mr. W. H. Perkins, F.R.S., in the chair.—A paper was read by Mr. S. Lupton On the formulae of the alums; the next was a notice On the colour of cupric chloride, by Mr. W. N. Hartley, who finds that the crystals of the salt when quite dry have a blue colour, and not a green, as they usually appear when slightly moist.—Papers were also read On the oxidation of the essential oils, Part II. by Mr. C. T. Kingzett; On the purification and boiling-point of methyl hexyl carbinol, by Mr. E. Neison; and a note on the boiling-point of methyl hexyl carbinol, by Dr. C. Schorlemmer, F.R.S.

Zoological Society, Dec. 1.—Dr. A. Günther, F.R.S., in the chair.—A letter was read from the Rev. S. J. Whitmee, of Samoa, stating that he had sent home for the Society some birds and a pair of the Samoan Bat, which had lately been described by Mr. Alston as *Pteropus whitmeei*. Particulars were given as to the habits of the latter.—A communication was read from Mr. Henry W. Piers, of Capetown, containing remarks on some specimens of *Gymnetrus* in the museum at Capetown.—The Secretary announced that Col. R. S. Tickell, late of H.M. Indian Army, had presented to the Society's library a very finely illustrated MS. work, in seven small folio volumes, on the Ornithology of India.—A communication was read from Mr. J. Brazier, of Sydney, N.S.W., giving descriptions of eleven new species of terrestrial and marine shells from North-east Australia.—A paper, by Messrs. P. L. Sclater and O. Salvin, was read on birds collected by Mr. Whitely in Western Peru, being the eighth communication made by the authors on this subject.—A communication was read from Mr. H. Whitely, containing some further notes on Humming Birds collected by him in High Peru.—Mr. A. G. Butler read a paper in which he gave descriptions of three new species of homopterous insects from various parts of the world.—Mr. A. H. Garrod gave some further particulars on the mechanism of the "shew off" in the Bustards, and described the peculiar structure of the *frenum linguae* recently noticed in a young male of the Great Bustard.