

the matter is not free from difficulty, for the difference of intensity with which a lateral sound is perceived by the two ears is not great. The experiment may easily be tried roughly by stopping one ear with the hand, and turning round backwards and forwards while listening to a sound held steadily. Calculation shows, moreover, that the human head, considered as an obstacle to the waves of sound, is scarcely big enough in relation to the wave-length to give a sensible shadow. To throw light on this subject I have calculated the intensity of sound due to a distant source at the various points on the surface of a fixed spherical obstacle. The result depends on the ratio (a) between the circumference of the sphere and the length of the wave. If we call the point on the spherical surface nearest to the source the anterior pole, and the opposite point (where the shadow might be expected to be most intense) the posterior pole, the results on three suppositions as to the relative magnitudes of the sphere and wave-length are given in the following table:—

	Intensity.
$a = 2$	Anterior pole '690
	Posterior pole '318
	Equator '356
$a = 1$	Anterior pole '503
	Posterior pole '285
	Equator '237
$a = \frac{1}{2}$	Anterior pole '294
	Posterior pole '260
	Equator '232

When, for example, the circumference of the sphere is but half the wave-length, the intensity at the posterior pole is only about a tenth part less than at the anterior pole, while the intensity is least of all in a lateral direction. When a is less than $\frac{1}{2}$, the difference of the intensities at the two poles is still less important, amounting to about one per cent. when $a = \frac{1}{3}$.

The value of a depends on the wave-length, which may vary within pretty wide limits, and it might be expected that the facility of distinguishing a lateral sound would diminish when the sound is grave. Experiments were accordingly tried with forks of a frequency of 128, but no greater difficulty was experienced than with forks of a frequency of 256, except such as might be attributed to the inferior loudness of the former. According to calculation the difference of intensity would here be too small to account for the power of discrimination.

PROF. HUXLEY'S LECTURES ON THE EVIDENCE AS TO THE ORIGIN OF EXISTING VERTEBRATE ANIMALS¹

VI.

IN the highest group of Vertebrates, the Mammalia, the perfection of animal structure is attained. It will hardly be necessary, indeed it will be impossible, in the time at our disposal, to give the general characters of the group, but our purpose will be answered as well by devoting a short time to considering the peculiarities of a single well-known animal, the evidence as to the origin of which approaches precision.

The horse is one of the most specialised and peculiar of animals, its whole structure being so modified as to make it the most perfect living locomotive engine which it is possible to imagine. The chief points in which its structure is modified to bring about this specialisation, and in which, therefore, it differs most markedly from other mammals, we must now consider.

In the skull the orbit is completely closed behind by bone, a character found only in the most modified mammals. The teeth have a very peculiar character. There

are, first of all, in the front part of each jaw, six long curved incisors or cutting teeth, which present a singular dark mark on their biting surfaces, caused by the filling in of a deep groove on the crown of each tooth, by the substances on which the animal feeds. After the incisors, comes on both sides of each jaw a considerable toothless interval, or *diastema*, and then six large grinding teeth, or molars and premolars. In the young horse a small extra premolar is found to exist at the hinder end of the diastema, so that there are, in reality, seven grinders on each side above and below; furthermore, the male horse has a tusk-like tooth, or canine, in the front part of the diastema immediately following the last incisor. Thus, the horse has, on each side of each jaw, three incisors, one canine, and seven grinders, making a total of forty-four teeth.

The grinding surfaces of the molars and premolars are very curious. In the upper jaw, each tooth is marked by four crescentic elevations, concave externally, the inner pair having each a curious folded mass connected with it. These projecting marks are formed of dentine and enamel, and, consequently, wear away more slowly than the intervening portions of the tooth, which are composed of cement. The lower grinders are marked with two crescents and two accessory masses, but the crescents are convex externally, and, consequently, when the opposite teeth bite together, the elevations do not correspond at any point. In this way a very perfect grinding surface is obtained. The teeth are of great length, and go on growing for a long time, only forming roots in old animals. All these points contribute to the perfection of the horse as a machine, by rendering the mastication of the food, and its consequent preparation for digestion in the stomach, as rapid and complete a process as possible.

It is, however, in the limbs that the most striking deviation from the typical mammalian structure is seen, the most singular modifications having taken place to produce a set of long, jointed levers, combining great strength with the utmost possible spring and lightness.

The humerus is a comparatively short bone inclined backwards: the radius is stout and strong, but the ulna seems to be reduced to its upper end—the olecranon or elbow; as a matter of fact, however, its distal end is left, fused to the radius, but the middle part has entirely disappeared: the carpus or wrist—the so-called “knee” of the horse—is followed by a long “cannon-bone,” attached to the sides of which are two small “splint-bones”; the three together evidently represent the metacarpus, and it can be readily shown that the great cannon-bone is the metacarpal of the third finger, the splint-bones those of the second and fourth. The splint-bones taper away at their lower ends and have no phalanges attached to them, but the cannon-bone is followed by the usual three phalanges, the last of which, the “coffin-bone,” is ensheathed by the great nail or hoof.

The femur, like the humerus, is a short bone, but is directed forwards; the tibia turns backwards, and has the upper end of the rudimentary fibula attached to its outer angle. The latter bone, like the ulna, has disappeared altogether as to its middle portion, and its distal end is firmly united to the tibia. The foot has the same structure as the corresponding part in the fore-limb—a great cannon-bone, the third metatarsal; two splints, the second and fourth; and the three phalanges of the third digit, the last of which bears a hoof.

Thus, in both fore and hind limb one toe is selected, becomes greatly modified and enlarged at the expense of the others, and forms a great lever, which, in combination with the levers constituted by the upper and middle divisions of the limb, forms a sort of double C-spring arrangement, and thus gives to the horse its wonderful galloping power.

In the river-beds of the Quaternary age—a time when England formed part of the Continent of Europe—

¹ A course of six lectures to working men, delivered in the theatre of the Royal School of Mines, Lecture VI., April 3. Continued from vol. xiii. p. 516.

abundant remains of horses are found, which horses resembled altogether our own species, or perhaps are still more nearly allied to the wild ass. The same is the case in America, where the species was very abundant in the Quaternary epoch—a curious fact, as, when first discovered by Europeans, there was not a horse from one end of the vast continent to the other.

In the Pliocene and older Miocene, both of Europe and America, are found a number of horse-like animals, resembling the existing horse in the pattern and number of the teeth, but differing in other particulars, especially the structure of the limbs. They belong to the genera *Protohippus*, *Hipparion*, &c., and are the immediate predecessors of the Quaternary horses.

In these animals the bones of the fore-arm are essentially like those of the horse, but the ulna is stouter and larger, can be traced from one end to the other, and, although firmly united to the radius, was not ankylosed with it. The same is true, though to a less marked extent, of the fibula.

But the most curious change is to be found in the toes. The third toe though still by far the largest, is proportionally smaller than in the horse, and each of the splint bones bears its own proper number of phalanges; a pair of "dew-claws," like those of the reindeer, being thus formed, one on either side of the great central toe. These accessory toes, however, by no means reached the ground, and could have been of no possible use, except in progression through marshes.

The teeth are quite like those of the existing horse, as to pattern, number, presence of cement, &c.; the orbit also is complete, but there is a curious depression on the face-bones, just beneath the orbit, a rudiment of which is, however, found in some of the older horses.

On passing to the older Miocene, we find an animal, known as *Anchitherium*, which bears, in many respects, a close resemblance to *Hipparion*, but is shorter-legged, stouter-bodied, and altogether more awkward in appearance. Its skull exhibits the depression mentioned as existing in *Hipparion*, but the orbit is incomplete behind, thus deviating from the specialised structure found in the horse, and approaching nearer to an ordinary typical mammal. The same is the case with the teeth, which are short and formed roots at an early period; their pattern also is simplified, although all the essential features are still retained. The valleys between the various ridges are not filled up with cement, and the little anterior premolar of the horse has become as large as the other grinders, so that the whole forty-four teeth of the typical mammalian dentition are well developed. The diastema is still present between the canines and the anterior grinding teeth—a curious fact in relation to the theory that the corresponding space in the horse was specially constructed for the insertion of the bit; for, if the Miocene men were in the habit of riding the *Anchitherium*, they were probably able to hold on so well with their hind legs as to be in no need of a bit.

The fibula is a complete bone, though still ankylosed below to the tibia; the ulna also is far stouter and more distinct than in *Hipparion*. In both fore and hind foot the middle toe is smaller, in relation to the size of the animal, than in either the horse or the *Hipparion*, and the second and fourth toes, though still smaller than the third, are so large that they must have reached the ground in walking. Thus, it is only necessary for the second and fourth toes, and the ulna and fibula to get smaller and smaller for the limb of *Anchitherium* to be converted into that of *Hipparion*, and this again into that of the horse.

Up to the year 1870 this was all the evidence we had about the matter, except for the fact that a species of *Palæotherium* from the older Eocene was, in many respects, so horse-like, having, however, well-developed ulna and fibula, and the second and fourth toes larger even than in *Anchitherium*, that it had every appearance of

being the original stock of the horse. But within the last six years some remarkable discoveries in central and western North America, have brought to light forms which are, probably, nearer the direct line of descent than any we have hitherto known.

In the Eocene rocks of these localities, a horse-like animal has been found, with three toes, like those of *Anchitherium*, but having, in addition, a little style of bone on the outer side of the fore foot, evidently representing the fifth digit. This is the little *Orohippus*, the lowest member of the Equine series.

This evidence is conclusive as far as the fact of evolution is concerned, for it is preposterous to assume that each member of this perfect series of forms has been specially created; and if it can be proved—as the facts adduced above certainly do prove—that a complicated animal like the horse may have arisen by gradual modification of a lower and less specialised form, there is surely no reason to think that other animals have arisen in a different way.

This case, moreover, is not isolated. Every new investigation into the Tertiary mammalian fauna brings fresh evidence, tending to show how the rhinoceros, the pigs, the ruminants, have come about. Similar light is being thrown on the origin of the carnivora, and also, in a less degree, on that of all the other groups of mammals.

It may well be asked why such clear evidence should be obtainable as to the origin of mammals, while in the case of many other groups—fish, for instance—all the evidence seems to point the other way. This question cannot be satisfactorily answered at present, but the fact is probably connected with the great uniformity of conditions to which the lower animals are exposed, for it is invariably the case that the higher the position of any given animal in the scale of being, the more complex are the conditions acting on it.

It is not, however, to be expected that there should be, as yet, an answer to every difficulty, for we are only just beginning the study of biological facts from the evolutionary point of view. Still, when we look back twenty years to the publication of the "Origin of Species," we are filled with astonishment at the progress of our knowledge, and especially at the immense strides it has made in the region of palæontological research. The accurate information obtained in this department of science has put the fact of evolution beyond a doubt; formerly, the great reproach to the theory was, that no support was lent to it by the geological history of living things; now, whatever happens, the fact remains that the hypothesis is founded on the firm basis of palæontological evidence.

THE LOAN COLLECTION CONFERENCES

CONSIDERABLE progress has been made in the arrangements for holding conferences in connection with the approaching Loan Collection of Scientific Apparatus at South Kensington.

In the Section of Mechanics, which includes Pure and Applied Mathematics and Measurement, the conferences will be held on May 17, 22, and 25, and the following gentlemen have promised to give addresses or to take part:—

Dr. Siemens, F.R.S., general address with special reference to Measurement.

Mr. F. J. Bramwell, F.R.S., on Prime Movers.

Mr. Barnaby, Director of Naval Construction to the Admiralty.—Naval Architecture.

Mr. W. Froude, M.A., F.R.S.—Fluid Resistance.

M. Tresca, Sous-Directeur du Conservatoire des Arts et Métiers, Paris.—Flow of Solids.

M. le General Morin, Directeur du Conservatoire des Arts et Métiers, Paris.—Ventilation of Public Buildings.

Sir Joseph Whitworth, Bart., F.R.S.—Linear Measurement.