	Droughts.	Floods.
Minimum Group (11th, 1st, and 2nd Series) ...	13	2
Intermediate Group (3rd, 4th, 6th, and 10th Series) ...	10	4
Maximum Group (5th, 6th, 7th, and 8th Series) ...	7	6

Of the seventeen cycles of eleven years, between 1711 and 1877, only three have no record of one or more dry years. The great droughts were those of 1722, 1778, 1792, 1825, 1845, and 1877, of which four occurred in the minimum group, one commenced in that group, but culminated in the intermediate group, and one was confined to the latter group.

Rio de Janeiro, June 12

ORVILLE A. DERBY

The Cell of the Bee

THE following simple construction shows in one figure all the elements of a cell of a honey-comb. On two rectangular axes take OA and OB equal to the side and diagonal of a square. Join AB and bisect it in C. Draw CD normal to AB. Join AD, and complete the rhombus whose sides are AD, DB. Then if OA be a side of the hexagon, ADBE is one of the three equal planes forming the trihedral angle which closes the cell. The three short diagonals DE meet in the vertex of the cell, and are

normal to each other. The three long diagonals AB form an equilateral triangle. OD is the height of the vertex above the hexagonal face of the prism. AED is the angle which the axis of the prism makes with each of the diagonals DE. ADO is the angle which the axis of the prism makes with each of the edges of the trihedral angle. The diagonals DE and AB are in the ratio of the side and diagonal of a square. Such a cell contains a maximum volume with a minimum surface.

Bardsea

EDWARD GEOGHEGAN

OUR ASTRONOMICAL COLUMN

THE REPORTED OBSERVATION OF "VULCAN."—In the telegram received from Mr. Lockyer relating to the solar eclipse which appeared in NATURE last week (p. 353), and which, like many other similar messages, had suffered in course of transmission, mention was made of Prof. Watson's observation of an object of $4\frac{1}{2}$ magnitude in R.A. 8h. 26m., and declination 18° 0' north, which was neither a known planet nor a star. θ Cancri, a star of the fifth magnitude, is less than a degree from this position, but the observer would doubtless be aware of its presence. A telegram to the same purport was received by M. Mouchez, the director of the Observatory at Paris. By the formula deduced by Leverrier from the observations of suspicious objects in transit across the sun's disc, if the indeterminate k be put = 0, the elongation in longitude of his hypothetical body from the sun's centre at the time of totality at Prof. Watson's station would be $5^\circ 9'$ eastward, and if $k = -1$, $9^\circ 5'$ westward, neither of which, it will be seen, accords with the position given in the telegram. The fourth-magnitude star, δ Cancri, must have been within the limits of the coronal surroundings of the sun, and was only just beyond them during the total eclipse of July 28, 1851, when no observer, to our knowledge, remarked the star. In the instructions for observing the eclipse issued from the United States Naval Observatory, Washington, and pre-

pared by Prof. Harkness at the instance of Admiral Rodgers, the superintendent, it is remarked: "As the truth of Leverrier's discovery of an apparently unexplained motion of the perihelion of Mercury is now established beyond all doubt, it is important to renew the search for an intra-Mercurial planet or planets." And to facilitate the work of such astronomers as might institute a search with considerable telescopic power, a chart was appended to the instructions showing every star so large as the seventh magnitude in that portion of the heavens occupied by the sun at the time. This chart extends from 7h. 32m. to 9h. 40m. in right ascension, and from 11° to 26° in declination. θ Cancri, the only star which appears near the position indicated for Prof. Watson's object, is marked on the chart as a sixth magnitude, which is the estimate of the *Uranometria* and *Durchmusterung*, but the star has been occasionally rated a fifth magnitude as in the first Radcliffe Catalogue, wherein particular attention was paid to the brightness of the stars. This is only a half-magnitude below Prof. Watson's estimate, but it remains to be seen from further intelligence whether there was any possibility of the star having been the object really noted; if it were separately remarked, or if the observed position does not admit of such change as would be necessary for identification, then it may truly be said that the American astronomer will have rendered the occasion of this eclipse a memorable one in the history of the science. Leverrier's confidence in the existence of an unexplained motion in the perihelion of Mercury and the necessity of accounting for it, by admitting the presence of matter in some form within the orbit of the planet, continued undiminished up to the time of his decease. One of his last communications to the writer of these lines was upon this subject.

THE LUNAR ECLIPSE ON AUGUST 12.—The eclipse of the moon on August 12 is the only one that will be wholly visible in this country until the year 1884; first contact with the earth's dark shadow at 10h. 42m., the middle of the eclipse at 12h. 8m., magnitude 0.59, and last contact at 13h. 34m. On October 4, 1884, there will be a total eclipse of the moon, the middle near 10 P.M., and the passage through the shadow nearly central.

THE AUGUST METEORS.—The earth will arrive at the descending node of the orbit of the comet 1862 III., in the track of which the meteors of the August period are found to move, soon after noon on Saturday next; the comet itself has now receded from the sun to a distance nearly equal to the mean distance of Neptune, to return to these parts of the system probably between 1980 and 1985. Moonlight interferes this year with observation of the smaller meteors August 9-11, during a part of the night.

TEMPEL'S COMET.—The following places of this comet are deduced from M. Schulhof's elements, with the time of perihelion passage, corrected by the early observations at Strasburg by Prof. Winnecke:—

12h. G.M.T.	Right Ascension.	North Polar Distance.	Distance from Earth.	Intensity of Light.
	h. m. s.			
August 23	16 20 30	109 3	9'018	0.86
" 25	25 55	109 52		
" 27	31 28	110 40	9'079	0.84
" 29	37 12	111 27		
" 31	43 6	112 13	9'144	0.82
September 2	49 10	112 58		
" 4	16 55	113 42	9'213	0.80
" 6	17 14	114 24		
" 8	8 22	115 5	9'286	0.77
" 10	15 5	115 44		
" 12	21 58	116 22	9'365	0.74
" 14	29 0	116 58		
" 16	17 36	117 33	9'448	0.71

On September 10 the comet passes very near to the orbit of Mars, but the planet is far distant. The dimen-