

ing this to the ethnogeny of Wales. Does it not seem probable that the early short race of long-skulled, mild-featured, stone-using people may have been the ancestors of the swarthy Silurians of Tacitus; while the later tall race of round-skulled, rugged-featured, bronze-using men may have represented the broad-headed, Keltic-speaking folk of history? At any rate, the evidence of craniology does not run counter to this hypothesis. For Dr. Beddoe's observations on head-forms in the west of England have shown that "heads which are ordinarily called brachycephalic belonged for the most part to individuals with light hair," while the short dark-haired people whom he examined were markedly dolichocephalic.<sup>1</sup> At the same time it must be admitted that his observations lend "no support to the view that the Keltic skull has been, or would be, narrowed by an admixture of the Iberian type." It should not, however, be forgotten that the same observer, in referring to a collection of crania from the Basque country preserved in Paris, says: "The form of M. Broca's Basque crania was very much that of some modern Silurian heads."<sup>2</sup>

According to the view advocated by Thurnam we have a right to anticipate that the oldest skulls found in this country would be of dolichocephalous type; and such I believe to be actually the case. Dr. Barnard Davis, it is true, has stated in the "Crania Britannica that the ancient British skull must be referred to the brachycephalic type; and such an induction was perfectly legitimate so long as the craniologist dealt only with skulls from the round barrows or from similar interments. But the long-barrow skulls examined by Prof. Rolleston,<sup>3</sup> and the Cissbury skulls recently studied by the same anatomist,<sup>4</sup> are decidedly dolichocephalic, as also are all the early prehistoric skulls which have been found of late years in France. While referring to craniology in this country, I may perhaps be allowed to remark that the eminent Italian anthropologist, Dr. Paolo Mantegazza, in a suggestive paper which has just appeared in his valuable journal, the *Archivio per l'Antropologia*, has referred to the Englishman's contempt for craniological work—work but little worthy of the practical spirit of the Anglo-Saxon race.<sup>5</sup> No doubt it is desirable to increase the number of our observations, but still the good-humoured remark about despising craniology can hardly be applied to a country which numbers among its living men of science such eminent craniologists as Prof. Busk, Prof. Cleland, Dr. Barnard Davis, and Professors Flower, Huxley, and Rolleston.

It may naturally be asked whether the researches of archaeologists in Wales lend any support to Thurnam's hypothesis. Nothing, I conceive, would be easier than to show that very material support has come from this quarter; but I have abstained, of set purpose, from introducing into this address any remarks on the prehistoric archaeology of Wales. For I have not forgotten that we are to have the privilege of hearing an evening lecture on "Primæval Man" by so distinguished an archaeologist and naturalist as Prof. Boyd Dawkins. No one has done more in this country to forward Thurnam's views, whether by actual exploration or by writing, than Prof. Boyd Dawkins; and if I have not referred to his work, especially to his discoveries in Denbighshire, it has been simply because I was anxious to avoid trespassing on any subject which he is likely to bring forward.<sup>6</sup>

Setting aside, then, any archaeological evidence derived from the bone-caves, barrows, or other sepulchres in Wales, we may finally look at the outcome of our inquiry into Welsh ethnogeny. If we admit, as it seems to me we are bound to admit, the existence of two distinct ethnical elements in the Welsh population, one of which is short, dark, and dolichocephalic—call it Silurian, Atlantean, Iberian, Basque, or what you will; and the other of which is tall, fair, and brachycephalic, such as some term Cymric and others Ligurian; then it follows that by

the crossing of these two races we may obtain not only individuals of intermediate character, but occasionally more complex combinations; for example, an individual may have the short stature and long head of the one race, associated with the lighter hair of the other; or again, the tall stature of one may be found in association with the melanism and dolichocephalism of the other race. It is therefore no objection to the views herein expressed if we can point to a living Welshman who happens to be at once tall and dark, or to another who is short and fair.

At the same time I am by no means disposed to admit that when we have recognised the union of the xanthous and melanic elements in Wales, with a predominance of the latter in the south, we have approached to anything like the exhausting limit of the subject. Still earlier races may have dwelt in the land, and have contributed something to the composition of the Welsh. In fact the anthropologist may say of a Welshman, as a character in "Cymbeline" says of Posthumus when doubtful about his pedigree—

"I cannot delve him to the root."

It is possible that the roots of the Welsh may reach far down into some hidden primitive stock, older mayhap than the Neolithic ancestors of the Silurians; but of such pristine people we have no direct evidence. So far however as positive investigation has gone, we may safely conclude that the Welsh are the representatives, in large proportion, of a very ancient race or races; and that they are a composite people who may perhaps be best defined as *Siluro-Cymric*.

Many other questions relating to Welsh ethnology press for consideration—such as the hypothesis that the Kymro was preceded, in parts at least of Wales, by the Gael; but such questions must be dismissed from present discussion, for I fear that my remarks have already overrun the limits of a departmental address. Let us hope however that much light may be thrown upon a variety of questions bearing upon local anthropology in the course of the discussions which will arise in this department during the present session of the Association.

#### PALÆONTOLOGICAL AND EMBRYOLOGICAL DEVELOPMENT<sup>1</sup>

SINCE the publication of the "Poissons Fossiles" by Agassiz and of the "Embryologie des Salmonidées" by Vogt, the similarity, traced by the former between certain stages in the growth of young fishes and the fossil representatives of extinct members of the group, has also been observed in nearly every class of the animal kingdom, and the fact has become a most convenient axiom in the study of palæontological and embryological development. This parallelism, which has been on the one side a strong argument in favour of design in the plan of creation, is now, with slight emendations, doing duty on the other as a newly-discovered article of faith in the new biology.

But while in a general way we accept the truth of the proposition that there is a remarkable parallelism between the embryonic development of a group and its palæontological history, yet no one has attempted to demonstrate this, or rather to show how far the parallelism extends. We have up to the present time been satisfied with tracing the general coincidence, or with striking individual cases.

The resemblance between the pupa stage of some Insects and of adult Crustacea, the earlier existence of the latter, and the subsequent appearance of the former in palæontological history, furnished one of the first and most natural illustrations of this parallelism; while theoretically the necessary development of the higher tracheate insects from their early branchiate aquatic ancestors seemed to form an additional link in the chain, and point to the Worms, the representatives of the larval condition of Insects, as a still earlier embryonic stage of the Articulates.

Indeed there is not a single group of the animal kingdom in which embryology has not played a most important part in demonstrating affinities little suspected before. The development of our frogs, our salamanders, has given us the key to much that was unexplained in the history of Reptiles and Batrachians. The little that has been done in the embryology of Birds has revolutionised our ideas of a class which at the beginning of the century seemed to be the most naturally circumscribed of all. Embryology and palæontology combined have led to the recognition of a natural classification uniting Birds and Reptiles on

<sup>1</sup> Address by Prof. Alexander Agassiz, vice-president, Section B, at the Boston Meeting of the American Association, August, 1880.

<sup>1</sup> *Ment. Anthropol. Soc. Lond.*, vol. ii. 1866, p. 350.

<sup>2</sup> *Ibid.*, p. 356.

<sup>3</sup> "On the People of the Long Barrow Period," *Journ. Anthropol. Inst.*, vol. v. 1876, p. 120.

<sup>4</sup> *Ibid.*, vol. vi. 1877, p. 20; vol. viii. 1879, p. 377.

<sup>5</sup> The whole passage so amusingly refers to the national idiosyncracies of craniologists, that it is well worth reproduction. "In Francia, Broca, il pontefice massimo dell' ipercraniologia moderna, col suo ardore eternamente giovanile, non studia più i crani, ma i cervelli; in Germania si prendono ancora misure sui teschi, ma con *rationabile obsequio*, quasi si dovesse adempiere ad un dovere noioso; in Inghilterra si continua a sprezzare la craniologia, come cosa poco degna dello spirito pratico della razza anglosassone; e in Italia, paese più scettico di tutti, perchè più antico e più stanco di tutti, si continua a misurare, pur sorridendo dell' improba e pur inutile fatica."

—"La Riforma Craniologica," *Archivio*, vol. x., 1880, p. 117.

<sup>6</sup> For Prof. Boyd Dawkins' contributions to the subject see his interesting works on "Cave-hunting," 1874, and on "Early Man in Britain," 1880.

the one side, and Batrachians and Fishes on the other. It is to embryology that we owe the explanation of the affinities of the Old Fishes in which Agassiz first recognised the similarity to the embryo of Fishes now living, and by its aid we may hope to understand the relationship of the oldest representatives of the class. It has given us the only explanation of the early appearance of the Cartilaginous Fishes, and of the probable formation of the earliest vertebrate limb from the lateral embryonic fold, still to be traced in the young of the Osseous Fishes of to-day.

Embryology has helped us to understand the changes aquatic animals must gradually undergo in order to become capable of living upon dry land. It has given us pictures of swimming-bladders existing as rudimentary lungs in Fishes with a branchial system; in Batrachians it has shown us the persistence of a branchial system side by side with a veritable lung. We find among the earliest terrestrial vertebrates types having manifest affinities with the Fishes on one side and Batrachians on the other, and we call these types Reptiles; but we should nevertheless do so with a reservation, looking to embryology for the true meaning of these half-fledged Reptiles, which lived at the period of transition between an aquatic and a terrestrial life, and must therefore always retain an unusual importance in the study of the development of animal life.

When we come to the embryology of the marine Invertebrates the history of the development of the barnacles is too familiar to be dwelt upon, and I need only allude to the well-known transformations of the Echinoderms, of the Acalephs, Polyps, in fact of every single class of Invertebrates, and perhaps in none more than in the Brachiopods, to show how far-reaching has been the influence of embryology in guiding us to a correct reading of the relations between the fossils of successive formations. There is scarcely an embryological monograph now published dealing with any of the later stages of growth which does not speak of their resemblance to some type of the group long ago extinct. It has therefore been most natural to combine with the attempts constantly made to establish the genetic sequence between the genera of successive formations an effort to establish also a correspondence between their palæontological sequence and that of the embryonic stages of development of the same, thus extending the mere similarity first observed between certain stages to a far broader generalisation.

It would carry me too far to sketch out, except in a most general way, even for a single class, the agreement known to exist in certain groups between their embryonic development and their palæontological history. It is hinted at in the succession of animal life of any period we may take up, and perhaps cannot be better expressed than by comparing the fauna of any period as a whole with that of following epochs;—a zoological system of the Jura, for instance, compared with one made up for the Cretaceous; next, one for the Tertiary, compared with the fauna of the present day. In no case could we find any class of the animal kingdom bearing the same definitions or characterised in the same manner. But apply to this comparison the data obtained from the embryological development of our present fauna, and what a flood of light is thrown upon the meaning of the succession of these apparently disconnected animal kingdoms, belonging to different geological periods, especially in connection with the study of the few ancient types which have survived to the present day from the earliest times in the history of our earth!

Although there is hardly a class of the animal kingdom in which some most interesting parallelism could not be drawn, and while the material for an examination of this parallelism is partially available for the Fishes, Molluscs, Crustacea, Corals, and Crinoids, yet for the illustration and critical examination of this parallelism I have been led to choose to-day a very limited group, that of Sea-urchins, both on account of the nature of the material and of my own familiarity with their development and with the living and extinct species of Echini. The number of living species is not very great,—less than three hundred,—and the number of fossil species thus far known is not, according to Zittel, more than about two thousand. It is therefore possible for a specialist to know of his own knowledge the greater part of the species of the group. It has been my good fortune to examine all but a few of the species now known to exist, and the collections to which I have had access contain representatives of the majority of the fossil species. Sea-urchins are found in the oldest fossiliferous rocks; they have continued to exist without interruption in all the strata up to the present time. While it is true that our knowledge of the Sea-urchins occurring

before the Jurassic period is not very satisfactory, it is yet complete enough for the purposes of the present essay, as it will enable me, starting from the Jurassic period, to call your attention to the palæontological history of the group, and to compare the succession of its members with the embryological development of the types now living in our seas. Ample material for making this comparison is fortunately at hand; it is material of a peculiar kind, not easily obtained, and which thus far has not greatly attracted the attention of zoologists.

Interesting and important as are the earliest stages of embryonic development in the different classes of the animal kingdom, as bearing upon the history of the first appearance of any organ and its subsequent modifications, they throw but little light on the subject before us. What we need for our comparisons are the various stages of growth through which the young Sea-urchins of different families pass from the time they have practically become Sea-urchins until they have attained the stage which we now dignify with the name of species. Few embryologists have carried their investigations into the more extended field of the changes the embryo undergoes when it begins to be recognised as belonging to a special class, and when the knowledge of the specialist is absolutely needed to trace the bearing of the changes undergone, and to understand their full meaning. Fortunately the growth of the young Echini has been traced in a sufficient number of families to enable me to draw the parallelism between these various stages of growth and the palæontological stages in a very different manner from what is possible in other groups of the animal kingdom, where we are overwhelmed with the number of species, as in the Insects or Mollusks, or where the palæontological or the embryological terms of comparison are wanting or very imperfect.

Beginning with the palæontological history of the regular sea-urchins at the time of the Trias, when they constituted an unimportant group as compared with the Crinoids, we find the Echini of that time limited to representatives of two families. One of these, the genus *Cidaris*, has continued to exist, with slight modifications, up to the present time, and not less than one-tenth of all the known species of fossil Echini belong to this important genus, which in our tropical seas is still a prominent one. It is interesting here to note that in the *Cidaridæ* the modifications of the test are not striking, and the fossil genera appearing in the successive formations are distinguished by characters which often leave us in doubt as to the genus to which many species should be referred. In the genus *Rhabdocidaris*, which appears in the lower Jura, and which is mainly characterised by the extraordinary development of the radioles, we find the extreme of the variations of the spines in this family. From that time to the present day the most striking differences have existed in the shape of the spines, not only of closely allied genera, but even in specimens of the same species; differences which in some of the species of to-day are as great as in older geological periods. The oldest *Cidaridæ* are remarkable for their narrow poriferous zones. It is only in the Jura that they widen somewhat; subsequently the pores become conjugated, and only later, during the Cretaceous period, do we find the first traces of any ornamentation of the test (*Temnocidaris*) so marked at the present day in the genus *Goniocidaris*. As far, then, as the *Cidaridæ* are concerned, the modifications which take place from their earliest appearance are restricted to slight changes in the poriferous zone and in the ornamentation of the test, accompanied with great variability in the shape of the primary radioles. We must except from this statement the genera *Diplocidaris* and *Tetracidaris*, to which I shall refer again. The representatives of the other Triassic family become extinct in the lower tertiary. The oldest genus, *Hemicidaris*, undoubtedly represents the earliest deviations from the true *Cidaris* type; modifications which affect not only the poriferous zone, but the test, the actinal and the abactinal systems, while from the extent of these minor changes we can trace out the gradual development of some of the characteristics in families of the regular Echini now living. The genus *Hemicidaris* may be considered as a *Cidaris* in which the poriferous zone is narrow and undulating, in which the granules of the ambulacral system have become minute tubercles in the upper portion of the zone and small primary tubercles in its actinal region, in which many of the inter-ambulacral granules become small secondaries, in which the plates of the actinal system have become reduced in number, and the apical system has become a narrow ring, and finally, in which the primary radioles no longer assume the fantastic shapes so common among the *Cidaridæ*.

We can trace in this genus the origin of the modifications of the poriferous zone, leading us, on the one side, through genera with merely undulating lines of pores to more or less distinct confluent arcs of pores, formed round the primary ambulacral tubercles, and, on the other, to the formation of open arcs of three or more pairs of pores. The first type culminates at the present day with the Arbaciadæ, the other with the Diadematiadæ, Triplechinidæ, and Echinometradæ. This specialisation very early takes place, for already in the lower Jura Stomechinus has assumed the principal characteristics of the Triplechinidæ of to-day.

Although in Hemicidaridæ the number of the coronal plates has increased as compared with the Cidaridæ, and while we find that in many genera, even of those of the present day, the number of the coronal plates is still comparatively small, yet, as a general rule, the more recent formations contain genera in which the increase in number of the interambulacral plates is accompanied by a corresponding decrease in the number of plates of the interambulacral area so characteristic thus far of the Cidaridæ and Hemicidaridæ, a change also affecting the size of the primary ambulacral tubercles. This increase in the number of the coronal plates is likewise accompanied by the development of irregular secondary and miliary tubercles, and the disappearance in this group of the granular tuberculation, so important a character in the Cidaridæ. With the increase in the number of the interambulacral coronal plates, the Pseudodiadematiadæ still retain prominent primary tubercles, recalling the earlier Hemicidaridæ and Cidaridæ, and, as in the Cidaridæ proper, the test is frequently ornamented by deep pits or by ridges formed by the junction of adjoining tubercles. The genital ring becomes narrower, and the tendency to the specialisation of one of its plates, the madreporite, more and more marked.

With the appearance of Stomechinus, the Echinidæ proper already assume in the Jura the open arcs of pores, the large number of coronal interambulacral plates, the specialisation of the secondary tubercles, and the large number of primary tubercles in each plate. With the appearance of Sphærechinus in the early Tertiary come in all the elements for the greater multiplication of the pairs of pores in the arcs of the poriferous zones, while the gigantic primary spines of some of the genera (Heterocentrotus), and the small number of primary tubercles are structural features which had completely disappeared in the group preceding the Echinometradæ, to which they appear most closely allied.

Going back again to the Hemicidaridæ, it requires but slight changes to pass from them to Acrosalenia and to the Saleniæ proper; the latter have continued to the present day, and have, like the Cidaridæ, retained almost unchanged the characters of the genera which preceded them, combined however with a few Cidaridian and Echinid features which date back to the Triassic period. We can thus trace the modifications which have taken place in the poriferous zone, the apical and actinal systems, the coronal plates, the ambulacral and interambulacral tubercles, as well as in the radioles, and in the most direct manner possible indicate the origin of the peculiar combination of structural features which we find at any geological horizon. On taking in succession the modifications undergone by the different parts of the test, we can trace each one singly, without the endless complication of combinations which any attempt to trace the whole of any special generic combination would imply.

Leaving out of the question for the moment the Palæchinidæ, we find no difficulty in tracing the history of the characters of the genera of the regular Echini which have existed from the time of the Trias and are now living, provided we take up each character independently. Nothing can be more direct than the gradual modification of the simple, barely undulating poriferous zone, made up of numerous ambulacral plates covered by granules, such as we find it among the Cidaridæ of the Trias, first into the slightly undulating poriferous zone of the Hemicidaridæ, next into the indistinct arcs of pores of the Pseudodiadematiadæ, then into the arcs with a limited number of pores of the Triplechinidæ, and finally to the polyporous arcs of the Echinometradæ. What can be more direct than the gradual modification to be traced in the development of the primary ambulacral tubercles, such as are characteristic of the Echinidæ of the present day, from their first appearance at the oral extremity of the ambulacral system of the Hemicidaridæ, and the increase in the number of primary interambulacral tubercles, accompanied by the growth of secondaries and miliaries, which we can trace in Hemicidaridæ, Acrosalenia, and Stomechinus—the

increase in number of the primary and secondary tubercles being accompanied by a reduction in the size of the radioles and a greater uniformity in their size and shape?

But while these modifications take place the original structural feature may be retained in an allied group. Thus the Cidaridæ retain unchanged from the earliest time to the present day the few primary tubercles, the secondary granules, the simple poriferous zone, the imbricating actinal system, and the few coronal plates, with the large apical system and many-shaped radioles; while in the Saleniæ the primary interambulacral tubercles, the secondary granules, the radioles, the genital ring, are recognised features of the Cidaridæ, associated, however, with an Echinid actinal and anal system, Hemicidarid primary ambulacral tubercles, and an Echinid poriferous zone. In the same way in the Diadematiadæ, the large primary interambulacral tubercles are Cidaridian features, while the structure of the ambulacral tubercles is Hemicidaridian. The existence of two kinds of spines is another Cidaridian feature, while the apical and actinal systems have become modified in the same direction as that of the Echinidæ. The more recent the genus the greater is the difficulty of tracing in a direct manner the origin of any one structural feature, owing to the difficulty of disassociating structural elements characteristic of genera which may be derived from totally different sources. This is particularly the case with genera having a great geological age. Many of them, especially among the Spatangoids, show affinities with genera following them in time, to be explained at present only on the supposition that when a structural feature has once made its appearance it may reappear subsequently apparently as a new creation, while in reality it is only its peculiar combination with structural features with which it had not before been associated (a new genus), which conceals in that instance the fact of its previous existence. A careful analysis, not only of the genera of the order, but sometimes of other orders which have preceded this combination in time, may often reveal the elements from which have been produced apparently unintelligible modifications.

There is, however, not one of the simple structural features in the few types of the Triassic and Liassic Echini from which we can so easily trace the origin of the structural features of all the subsequent Echinid genera, which is not also itself continued to the present day in some generic type of the present epoch, fully as well characterised as it was in the beginning. In fact, the very existence to-day of these early structural features seems to be as positive a proof of the unbroken systematic affinity between the Echini of our seas and those of the Trias as the uninterrupted existence of the genus *Pygaster* or *Cidaris* from the Trias down to the present epoch, or of the connection of many of the genera of the Chalk with those of our epoch (*Salenia*, *Cyphosoma*, *Psammechinus*, &c.).

Passing to the Clypeastridæ, we find there, as among the *Desmosticha*, that the earliest type, *Pygaster*, has existed from the Trias to the present time; and that, while we can readily reconstruct, on embryological grounds, the modifications the earliest *Desmosticha*-like Echini should undergo in order to assume the structural features of *Pygaster*, yet the early periods in which the precursors of the Echinoconidæ and Clypeastridæ are found have thus far not produced the genera in which these modifications actually take place. But, starting from *Pygaster*, we naturally pass to *Holactypus*, to *Discoidea*, to *Conoclypus*, on the one side, while on the other, from *Holactypus* to *Echinocyamus*, *Sismondia*, *Fibularia*, and *Mortonia*, we have the natural sequence of the characters of the existing Echinanthidæ, *Laganidæ*, and *Scutellidæ*, the greater number of which are characteristic of the present epoch. If we were to take in turn the changes undergone in the arrangement of the plates of the test, as we pass from *Pygaster* to *Holactypus*, to *Echinocyamus*, and the Echinanthidæ, we should have in the genera which follow each other in the palæontological record an unbroken series showing exactly what these modifications have been. In the same way the modifications of the abactinal and anal systems, and those of the poriferous zone, can equally well be followed to *Echinocyamus*, and thence to the Clypeastridæ; while a similar sequence in the modifications of these structural features can be followed from *Mortonia* to the *Scutellidæ* of the present period.

Passing finally to the *Petalosticha*, we find no difficulty in tracing theoretically the modifications which our early Echinoconidæ of the Liass should primarily undergo previous to the appearance of *Galeropygus*. The similarity of the early *Cassiduloid* and *Echinoneoid* types points to the same systematic

affinity, and perhaps even to a direct and not very distant relationship with the Palæchinidæ. For if we analyse the Echinothuria of the present day, we find in genera like *Phormosoma* many structural features, such as the shape of the test, the character of the spines, the structure of the apical system, that of the poriferous zone, indicative of possible modifications in the direction of *Pygaster* or of *Galeropygus*, which have as yet not been taken into account.

Adopting for the Petalosticha the same method of tracing the modifications of single structural features in their palæontological succession, we trace the comparatively little modified palæontological history of the Echinoneidæ of the present day from the *Pyrina* of the lower Jura. This, in its turn, has been preceded by *Hyboclypus* and *Galeropygus*, while the Echinolampadæ of the present day date back, with but trifling modifications, to the Echinobrissus of the Lias, itself preceded by *Clypeus*: and they have been subject only to slight generic changes since that time, Echinobrissus being still extant, while such closely-allied genera as *Catopygus* and *Cassidulus* of the earlier Cretaceous are still represented at the present day; the modifications taking place in the actinal system, in the ambulacral zones of the Echinoneidæ and of the Echinolampadæ showing the closest possible systematic affinity in these families. Starting again from *Hyboclypus*, with its elongate apical system, we naturally pass to *Collyrites* and the strange *Dysasteridæ* forms which in their turn are closely allied to the *Holasteridæ*. From *Holaster* on the one side and from *Toxaster* on the other, we find an unbroken sequence of structural characters uniting the successive genera of *Holasteridæ*, such as *Cardiaster*, *Offaster*, *Stenonia*, *Ananchytes*, and *Asterostoma*, with *Palæopneustes*, *Homolampas*, and the *Pourtalesidæ* of the present day, while from the genera of the *Toxasteridæ* we naturally pass to the cretaceous *Hemiaster*; in this genus and the subsequent *Micraster* we find all the elements necessary for the modifications which appear in the *Spatanginæ* from the time of the Chalk to the present day. These modifications result in genera in which we trace the development of the fascioles, of the actinal, anal, and abactinal plastrons, of the beak, the formation of the petaloid ambulacra, first flush with the test, and little by little changed into marsupial pouches, the growth of the anterior groove and the manifold modifications of the ambulacral system in *Spatangus*, *Agassizia*, and *Echinocardium*, often recalling in some of its features structural characters of families which have preceded this in time.

Apparently in striking contrast with the Echini of the secondary period and those which have succeeded them stand the Palæozoic Echini; but when we have examined the embryology of Echini, we shall be better prepared to understand their structure and the affinities of the Palæchinidæ with the Echini of the present day, and their immediate predecessors.

Taking up now the embryological development of the several families which will form the basis of our comparisons, beginning with the *Cidaridæ*, we find that in the earliest stages they very soon assume the characters of the adult, the changes being limited to the development of the abactinal system, the increase in number of the coronal plates, and the modifications of the proportionally gigantic primary radioles.

In the *Diadematiidæ* the changes undergone by the young are limited to the gradual transformation of the embryonic spines to those which characterise the family, to the changes of the vertical row of pores in the ambulacral area into arcs of three or four pairs of pores, and to the specialisation of the actinal and abactinal systems.

In the *Arbaciadæ* the young stages are remarkable for the prominent sculpture of the test, for the flattened spines, for their simple poriferous zone, for their actinal system, and for their genital ring. The anal plates appear before the genital ring.

In the *Echinometradæ* the young thus far observed are characterised by the small number of their primary tubercles, the large size of the spines, the simple vertical row of pores, the closing of the anal ring by a single plate, and the turban-shaped outline of the test. Little by little the test loses with increasing age this *Cidaridæ*-like character; it reminds us, from the increase in the number of its plates, more of *Hemicidaridæ*; then, with their still greater increase, of the *Pseudodiadematidæ*; and, finally, of the *Echinometradæ* proper. The spines, following *pari passu* the changes of the test, lose little by little their fantastic embryonic, or rather *Cidaridæ*-like appearance, and become more solid and shorter, till they finally assume the delicately fluted structure characteristic of the *Echinometradæ*. The vertical poriferous zone is first changed into a series of

connected vertical arcs, which become disjointed, and form, with increasing age, the independent arcs of pores, of three or more pairs of pores, of the *Echinometradæ*.

In the *Echinidæ* proper we find in the young stages the same unbroken vertical line of pores, which gradually becomes changed to the characteristic generic types. We find, as in the *Echinometradæ*, an anal system closed with a single plate, and an abactinal system separating in somewhat more advanced stages from the coronal plates of the test. This is as yet made up of a comparatively small number of plates, carrying but few large primary tubercles, with fantastically shaped spines entirely out of proportion to the test, but which, little by little, with the increase of the number of coronal plates, the addition of primary tubercles, and their proportional decrease in size, assume more and more the structure of the genus to which the young belongs. The original anal plate is gradually lost sight of from the increase in number of the plates covering the anal system, and it is only among the *Temnopleuridæ* that this anal plate remains more or less prominent in the adult. In the *Salenidæ*, of which we know as yet nothing of the development, this embryonic plate remains permanently a prominent structural feature of the apical system.<sup>1</sup>

Among the *Clypeastroids* the changes of form they undergo during growth are most instructive. We have in the young *Fibularinæ* an ovoid test, a small number of coronal plates surmounted by few and large primary tubercles, supporting proportionally equally large primary radioles, simple rectilinear poriferous zones, no petaloid ambulacra—in fact, scarcely one of the features we are accustomed to associate with the *Clypeastroids* is as yet prominently developed. But rapidly, with increasing size, the number of primary tubercles increases, the spines lose their disproportionate size, the pores of the abactinal region become crowded and elongate, and a rudimentary petal is formed. The test becomes more flattened, the coronal plates increase in number, and it would be impossible to recognise in the young *Echinocyamus*, for instance, the adult of the *Cidaridæ*-like or *Echinometra*-like stages of the *Sea-urchin*, had we not traced them step by step. Most interesting, also, is it to follow the migrations of the anal system, which, to a certain extent, may be said to retain the embryonic features of the early stages of all *Echinoderm* embryos, in being placed in more or less close proximity to the actinostome. What has taken place in the growth of the young *Echinocyamus* is practically repeated for all the families of *Clypeastroids*; a young *Echinarachnius*, or *Mellita*, or *Encope*, or a *Clypeaster* proper, resembles at first more an *Echinometra* than a *Clypeastroid*; they all have simple poriferous zones and spines and tubercles out of all proportion to the size of the test.<sup>2</sup>

When we come to the development of the *Spatangoids*, we find their younger stages also differing greatly from the adult. Among the *Nucleolidæ*, for instance, the young stages have as yet no petals, but only simple rectilinear poriferous zones. They are elliptical with a high test, with a single large primary tubercle for each plate, and a simple elliptical actinostome, without any trace of the typical bourrelets and phyllodes so characteristic of this family. Very early, however, this condition of things is changed, the test soon becomes more flattened, the petals begin to form as they do in the *Clypeastroids*, and we can soon trace the rudiments of the peculiar bourrelets characteristic of the family, accompanied by a rapid increase in the number of tubercles and in that of the coronal plates.

Among the *Spatangidæ* some are remarkable in their adult condition for their labiate actinostome, for the great development of the petals, for the presence of fascioles surrounding certain definite areas, for the small size of the tubercles, the general uniformity in the spines of the test, and the specialisation of their anterior and posterior regions. On examining the young stages of this group of *Spatangoids*, not one of these structural features is as yet developed. The actinostome is simple, the poriferous zone has the same simple structure from the actinostome to the apex, the primary tubercles are large, few in number, surrounded by spines which would more readily pass as the spines of *Cidaridæ* than of *Spatangoids*. The fascioles are either very indistinctly indicated, or else the special lines have

<sup>1</sup> The young of the following genera have served as a basis for the preceding analysis of the embryonic stages of the *Desmosticha*: *Cidaris*, *Doricidaris*, *Gonicidaris*, *Arbacia*, *Porocidaris*, *Strongylocentrotus*, *Echinometra*, *Echinus*, *Toxopneustes*, *Hipponoe*, *Temnopleurus*, *Temnechinus*, and *Trigonocidaris*.

<sup>2</sup> Among the *Clypeastroids* I have examined the young of *Echinocyamus*, *Fibularia*, *Mellita*, *Laganum*, *Echinarachnius*, *Encope*, *Clypeaster*, and *Echinanthus*.

not as yet made their appearance; the ambulacral suckers of the anterior zone are as large and prominent as those of the young stages of any of the regular Echini. It is only little by little, with advancing age, that we begin to see signs of the specialisation of the anterior and posterior parts of the test, that we find the characteristic anal or lateral fascioles making their appearance, only with increasing size that the spines lose their Cidarid-like appearance, that the petals begin to be formed, and that the simple actinostome develops a prominent posterior lobe. In the genus *Hemiaster*, the young stages are specially interesting, as long before the appearance of the petals, while the poriferous zone is still simple, the total separation of the bivium and of the trivium of the ambulacral system, so characteristic of the earliest Spatangoids (the *Dysasterinæ*), is very apparent.<sup>1</sup>

From this rapid sketch of the changes of growth in the principal families of the recent Echini we can now indicate the transformations of a more general character through which the groups as a whole pass.

In the first place, while still in the *Pluteus* all the young Echini are remarkable for the small number of coronal plates, for the absence of any separation between the actinal and abactinal systems and the test proper. They all further agree in the large size of the primary spines of the test, whether it be the young of a *Cidarid*, an *Arbacia*, an *Echinus*, a *Clypeaster*, or a Spatangoid. They all in their youngest stages have simple vertical ambulacral zones; beyond this, we find as changes characteristic of some of the *Desmosticha*, the specialisation of the actinal system from the coronal plates, the formation of an anal system, the rapid increase in the number of the coronal plates, with a corresponding increase in the number of the spines and a proportional reduction of their size, the formation of an abactinal ring, and the change of the simple vertical poriferous zone into one composed of independent arcs.

In the Spatangoids and Clypeastroids we find common to both groups the shifting of the anal system to its definite place, the modifications of the abactinal part of the simple ambulacral system in order to become petaloid, and the gradual change of the elliptical ovoid test of the young to the characteristic generic test, accompanied by the rapid increase in the number of the primary tubercles and spines. Finally limited to the Spatangoids are the changes they undergo in the transformation of the simple actinostome to a labiate one, the specialisation of the anterior and posterior parts of the test, and the definite formation of the fascioles.

Comparing this embryonic development with the palæontological one, we find a remarkable similarity in both, and in a general way there seems to be a parallelism in the appearance of the fossil genera and the successive stages of the development of the Echini as we have traced it.

We find that the earlier regular Echini all have more or less a *Cidarid*-like look—that is, they are Echini with few coronal plates, large primary tubercles, with radioles of a corresponding size; that it is only somewhat later that the *Diademopsidæ* make their appearance, which, in their turn, correspond within certain limits to the modifications we have traced in the growth of the young *Diadematiidæ* and *Arbaciadæ*. The separation of the actinal system from the coronal plates has been effected. The poriferous zone has either become undulating, or forms somewhat indefinite open arcs; we find in all the genera of this group a larger number of coronal plates, more numerous primaries, the granules of the *Cidaridæ* replaced by secondaries and miliaries, and traces of a *Hemicidarid*-like stage in the size of the actinal ambulacral tubercles.

Comparing in the same way the palæontological development of the Echinidæ proper, we find that, on the whole, they agree well with the changes of growth we can still follow to-day in their representatives, and that, as we approach nearer the present epoch, the fossil genera more and more assume the structural features which we find developed last among the Echinidæ of the present day. Very much in the same manner as a young *Echinus* develops, they lose, little by little, first their *Cidarid* affinities, which become more and more indefinite, next their *Diadematiid* affinities, if I may so call the young stages to which they are most closely allied, and finally, with the increase in the number of the coronal plates, the great numerical development of the primary tubercles and spines, and that of the secondaries and miliaries which we can trace in the fossil Echini

<sup>1</sup> For this sketch of the embryology of the *Petalosticha* I have examined the young of *Echinolampas*, *Echinoneus*, *Echinocardium*, *Brissopsis*, *Agassizia*, *Spatangus*, *Brissus*, and *Hemiaster*.

of the Tertiaries, we pass insensibly into the generic types characteristic of the present day.

Although we know nothing of the embryology of the *Salenidæ*, yet, like the *Cidaridæ*, they have in a great measure remained a persistent type, the modifications of the group being all in the same direction as those noticed in the other *Desmosticha*; a greater number of coronal plates, the development of secondaries and miliaries combined with a specialisation of the actinal system not found in the *Cidaridæ*.

An examination of the succession of the *Echinoconidæ* shows but little modification from the earliest types; the changes, however, are similar to those undergone by the *Clypeastroids* and *Petalosticha*, though they do not extend to modifications of the poriferous zone, but are mainly changes in the actinostome and in the tuberculation. In fact, the group of *Echinoconidæ* seems to hold somewhat the same relation to the *Clypeastroids* which the *Salenidæ* hold to the *Cidaridæ*, and the earliest genus of the group *Pygaster* has remained, like *Cidarid*, a persistent type to the present day.

The earliest *Clypeastroids* are all forms which resemble the *Fibularina* and the genera following *Echinocyamus* and *Fibularia*; they are mainly characterised by the same changes which an *Echinarachnius* or a *Mellita*, for instance, undergoes as it passes from its *Echinocyamus* stage to the *Laganum* or *Encope* stage. The comparison is somewhat more complicated when we come to the Spatangoids. The comparison of the succession of genera in the different families, as traced in the *Desmosticha* and *Clypeastroids*, is made difficult from the persistency of the types preceding the *Echinoconidæ* and the *Ananchytidæ*, which have remained without important modifications from the time of the lower Cretaceous; previous to that time the modifications of the *Cassidulidæ* are found to agree with the changes which have been observed in the growth of *Echinolampas*. The early genera, like *Pygurus*, have many of the characteristics of the test of the young *Echinolampas*. The development of prominent bourrelets and of the foscelle and petals goes on side by side with that of genera in which the modification of the actinostome, of the test, and of the petals is far less rapid, one group retaining the *Echinoneus* features, the other culminating in the *Echinolampas* of the present day, and having likewise a persistent type, *Echinobrissus*, which has remained with its main structural features unchanged from the Jura to the present day. That is, we find genera of the *Cassidulidæ* which recall the early *Echinoneus* stage of *Echinolampas*, next the *Caratomus* stage, after which the foscelle, bourrelets, and petals of the group become more prominent features of the succeeding genera. Accompanying the persistent type *Echinobrissus*, genera appear in which either the bourrelets or petals have undergone modifications more extensive than those of the same parts in the genera of the *Echinoneus* or *Caratomus* type.

The earliest Spatangoids belong to the *Dysasteridæ*, apparently an aberrant group, but which, from the history of the young *Hemiaster*, we now know to be a strictly embryonic type, which, while it thus has affinities with the true Spatangoids, still retains features of the *Cassidulidæ* in the mode of development of the actinostome and of the petals, as well as of the anal system. The genera following this group, *Holaster* and *Toxaster*, can be well compared, the one to the young stages of *Spatangus* proper before the appearance of the petals, when the ambulacra are flush with the test, and when its test is more or less ovoid, the other to a somewhat more advanced stage, when the petals have made their appearance as semi-petals. In both cases the actinostome has the simple structure characteristic of all the young Spatangoids. The changes we notice in the genera which follow them lead in the one case through very slight modifications of the abactinal system, of the anterior and posterior extremities of the test, to the *Ananchytid*-like Spatangoids of the present day, the *Pourtalesidæ*, the genus *Holaster* itself persisting till well into the middle of the Tertiary period; while on the other side we readily recognise in the Spatanginæ which follow *Toxaster* (a persistent type which has continued as *Palæostoma* to the present day) the genera which correspond to the young stages of such Spatangoids as *Spatangus* and *Brissopsis* of the present day, genera which, on the one hand, lead from *Hemiaster* (itself still represented in the present epoch), through stages such as *Cyclaster*, *Peripneustes*, *Brissus*, and *Schizaster*, and, on the other, through *Micraster* and the like, to the Spatangoids, in which the development of the anal plastron and fasciole performs an important part, while in the former group the development of the peripetalous fasciole and of the lateral

fasciole can be followed. None of the genera of *Petalosticha* belonging to the other groups develop any fasciole in the sense of circumscribing a limited area of the test.

The comparison of the genera of *Echini* which have appeared since the Lias with the young stages of growth of the principal families of *Echini*, shows a most striking coincidence, amounting almost to identity, between the successive fossil genera and the various stages of growth. This identity can, however, not be traced exactly in the way in which it has usually been understood, while there undoubtedly exists in the genera which have appeared one after the other a gradual increase in certain families in the number of forms, and a constant approach in each succeeding formation, in the structure of the genera, to those of the present day. It is only in the accordance between some special points of structure of these genera and the young stages of the *Echini* of the present day that we can trace an agreement which, as we go further back in time, becomes more and more limited. We are either compelled to seek for the origin of many structural features in types of which we have no record, or else we must attempt to find them existing potentially in groups where we had as yet not succeeded in tracing them. The parallelism we have traced does not extend to the structure as a whole. What we find is the appearance among the fossil genera of certain structural features giving to the particular stages we are comparing their characteristic aspect. Thus, in the succession of the fossil genera, when a structural feature has once made its appearance, it may either remain as a persistent structure, or it may become gradually modified in the succeeding genera of the same family, or it may appear in another family, associated with other more marked structural features which completely overshadow it. Take, for instance, among the *Desmosticha* the modifications of the poriferous zone of the actinal and abactinal systems of the coronal plates, of the ambulacral and interambulacral systems, the changes in the relative proportion of the primary tubercles, and the development of the secondaries. These are all structural features which are modified independently one of the other; we may find simultaneous development of these features in parallel lines, but a very different degree of development of any special feature in separate families.

This is as plainly shown in the embryological as in the palæontological development. In the *Cidaridæ* there is the minimum of specialisation in these structural features. In the *Diademopsidæ* there is a greater range in the diversity of the structure of the poriferous zone and of the coronal plates, as well as of the actinal system. There is a still greater range among the *Echinidæ*, while among the *Salenidæ* the modifications, as compared to those of the *Echinidæ* and *Diademopsidæ*, are somewhat limited again, being restricted as far as relates to the poriferous zone and coronal plates, but specialised as far as the actinal system is concerned, and specially important with reference to the structure of the apical system. The special lines in which these modifications take place produce, of course, all possible combinations, yet they give us the key to the sudden appearance, as it were, of structural features of which the relationship must be sought in very distantly related groups. It is to this speciality in the palæontological development that we must trace, for instance, the *Cidarid* affinities of the *Saleniæ*, their papillæ, the existence of few large primary interambulacral tubercles, the structure of their apical system, and their large genital plates; while it is to their affinities with the *Hemicidaridæ* that we must refer the presence of the few larger primary ambulacral tubercles at the base of the ambulacral area, and by their *Diademopsid* and *Echinidian* affinities that we explain the indented imbricated actinal system with the presence of a few genuine miliaries. But all the structural features which characterise the earliest types of the *Desmosticha* can in reality be traced, only in a somewhat rudimentary form, even in the *Cidaridæ*. The slight undulation of the closely packed, nearly vertical poriferous zone is the forerunner of the poriferous zone first separated into vertical arcs and then into independent arcs. The limitation in the number of the rows of granules in the ambulacral zone, and their increase in size, is the first trace of the appearance of the somewhat larger primary ambulacral tubercles of the *Hemicidaridæ* and *Saleniæ*. The existence of the smooth cylindrical spines of the abactinal region of the test naturally leads to similar spines covering the whole test in the other families of the *Desmosticha*. The difference existing in the plates covering the actinal system from those of the coronal plates leads to the great distinction between the structure of the actinal system and of the coronal plates in some of the *Echinidæ*.

Passing to the *Clypeastridæ* and *Petalosticha*, we trace a parallelism of the same kind, and readily follow in the successive genera of fossil *Clypeastroids*, but often in widely separated genera, the precise modifications which the poriferous zone has undergone as it first becomes known to us in *Echinocyamus* and *Fibularia*, and as we find it in the most complicated petaloid stage of the *Clypeastroids* of the present day. We readily trace the changes the test undergoes from its comparatively ovoid and swollen shape to assume first that of the less gibbous forms, next that of the *Laganidæ*, and finally of the flat *Scutellidæ*; while we trace in the *Echinanthidæ* the persistent structural features of some of the earliest *Clypeastroids*, together with an excessive modification of the poriferous zone. Likewise for the *Echinoconidæ* we trace mainly the slight modifications of the poriferous zone and of the coronal plates, and finally, when we come to the *Spatangidæ* we find no difficulty in tracing from the most *Desmostichoid* of the *Spatangoid* genera the modifications of a test in which the ambulacral and interambulacral areas are next up of plates of nearly uniform size, in which the anterior and posterior extremities are barely specialised, to the most typical of the *Ananchytidæ*, in which the anterior and posterior extremities have developed the most opposite and extraordinary structural features. In a similar way we can trace among the fossil genera of different families the gradual development of the actinal plastron from its very earliest appearance as a modification of the posterior interambulacral area of the actinal side, or the growth of the posterior beak into an anal snout, the successive changes of the anal groove, the formation of the actinal labium, or the development of the bourrelets and phylloides from a simple circular actinostome, the gradual deepening of the slight anterior groove of some early *Spatangoid* to form the deeply sunken actinal groove. Equally well we can trace the modifications of the ambulacral system as it passes from the simple poriferous zones of the earlier *Spatangoids* to genera in which the petaliferous portion makes its appearance, and finally becomes the specialised structure of our recent *Spatangoid* genera, such as *Schizaster*, *Moira*, and the like. Finally, we can trace to a certain extent the development of the fascioles on one side from genera like *Hemiaster*, in which the peripetalous fasciole is prominent, to genera like *Brissopsis*, *Brissus*, and the like, of the present day; on the other, perhaps, or both combined, the formation of a lateral and anal fasciole from genera like *Micraster* in *Spatangus* and *Agassizia*. Thus we must, on the same theory of the independent modifications of special structural features, trace the many and complicated affinities which so constantly strike us in making comparative studies, and which render it impossible for us to express the manifold affinities we notice, without taking up separately each special structure. Any attempt to take up a combination of characters, or a system of combinations, is sure to lead us to indefinite problems far beyond our power to grasp.

In the oldest fossil *Clypeastroids* and *Petalosticha*, as well as in the *Desmosticha*, we also find the potential expression of the greater number of the modifications subsequently carried out in genera of later date. The semipetaloid structure of some of the earlier genera of *Spatangoids*, the slight modifications of some of the plates of the actinal side near the actinostome, are the precursors, the one of the highly complicated petaloid ambulacra of the recent *Spatangoids*, the other of the actinal plastron, leading as it does also to the important differences subsequently developed in the anterior and posterior extremities of the test, as well as to the modifications which lead to the existence of a highly labiate actinostome. The appearance of a few miliaries near the actinostome constitutes the first rudimentary bourrelets.

Going back now to the *Palæchinidæ*, the earliest representatives of the *Echini* in palæozoic times, without any attempt to trace the descent of any special type from them, we may perhaps find some clue to the probable modifications of their principal structural features preparatory to their gradual disappearance. In the structure of the coronal plates, the specialisation of the actinal and abactinal systems, the conditions of the ambulacral system, we must compare them to stages in the embryonic development of our recent *Echini* with which we find no analogues in the fossil *Echini* of the Lias and the subsequent formations. In order to make our parallelism, we must go back to a stage in the embryonic history of the young *Echini* in which the distinction to be made between the ambulacral and interambulacral systems is very indefinite, in which the apical system is, it is true, specialised, but in which the actinal system remains practically a part

of the coronal system. But here the comparison ceases, and, although we can trace in the palæontological development of such types as Archæocidaris or Bothriocidaris modifications which would lead us without great difficulty, on the one side to the Cidaridæ, and on the other to the Echinothuriæ and Diadematiidæ of the present day, we cannot fail to see most definite indications in some of the structural features of the Palæchinidæ of characteristics which we have been accustomed to associate with higher groups. The minute tuberculation, for instance, of the Clypeastroids and Spatangoids, already existing in the Melonitidæ, the genital ring, and anal system, are quite as much Echinid as Cidarid. The polyporus genera of the group represent to a certain extent the polypteri of the regular Echini, and the lapping of the actinal plates of the Cidaridæ and of the coronal plates in some of the Diadematiidæ, as well as the existence of such genera as Tetracidaris, of four interambulacral plates in *Astropyga*, and of a large number of ambulacral plates in some of the recent Echinometradæ, all these are Palæchinid characters which we can explain on the theory of the independent development of the structural features of which they are modifications. We should, however, remember, that the existence of a large number of coronal plates, especially interambulacral plates, in the Palæchinidæ, is a mere vegetative character, which they hold in common with all the Crinoids,—a character which is reduced to a minimum among the Holothurians, and still persists in full force among the Pentacrinis of the present day, as well as the Astrophytidæ and Echinidæ.

It would lead me too far to institute the same comparison between the embryonic stages of the different orders of Echinoderms and their earliest fossil representatives. We may, however, in a very general way, state that we know the earliest embryonic stages of the order of Echinoderms of to-day, which, with the exception of the Blastoidæ and Cystideans, are identical with the fossil orders, and that as far as we know they all begin at a stage where it would be impossible to distinguish a Sea-urchin from a Star-fish, or an Ophiuran, or a Crinoid, or an Holothurian,—a stage in which the test, calyx, abactinal and ambulacral systems are reduced to a minimum. From this identical origin there is developed at the present day, in a comparatively short period of time, either a Star-fish, a Sea-urchin, or a Crinoid; and if we have been able successfully to compare, in the development of typical structures, the embryonic stages of the young Echini with their development in the fossil genera, we may fairly assume that the same process is applicable when instituting the comparison within the different limits of the orders, but with the same restrictions. That is, if we wish to form some idea of the probable course of transformations which the earliest Echinoderms have undergone to lead us to those of the present day, we are justified in seeking for our earliest representatives of the orders such Echinoderms as resemble the early stages of our embryos, and in following, for them as for the Echini, the modifications of typical structures. These we shall have every reason to expect to find repeated in the fossils of later periods, and, going back a step further we may perhaps get an indefinite glimpse of that first Echinodermal stage which should combine the structural features common to all the earliest stages of our Echinoderm embryos.

And yet, among the fossil Echinoderms of the oldest periods, we have not as yet discovered this earliest type from which we could derive either the Star-fishes, Ophiurans, Sea-urchins, or Holothurians. With the exception of the latter, which we can leave out of the question at present, we find all the orders of Echinoderms appearing at the same time. But while this is the case, one of the groups attained in these earliest days a prominence which it gradually loses with the corresponding development of the Star-fishes, Ophiurans, and Sea-urchins, it has steadily declined in importance; it is a type of Crinoids, the Cystideans which culminated during Palæozoic times, and completely disappeared long before the present day. If we compare the early types of Cystideans to the typical embryonic Echinodermal type of the present day, we find they have a general resemblance, and that the Cystideans and Blastoids represent among the fossil Echinoderms the nearest approach we have yet discovered to this imaginary prototype of Echinoderms.

This may not seem a very satisfactory result to have attained. It certainly has been shown to be an impossibility to trace in the palæontological succession of the Echini anything like a sequence of genera. No direct filiation can be shown to exist, and yet the very existence of persistent types, not only among Echinoderms, but in every group of marine animals, genera which have

continued to exist without interruption from the earliest epochs at which they occur to the present day, would prove conclusively that at any rate some groups among the marine animals of the present day are the direct descendants of those of the earliest geological periods. When we come to types which have not continued as long, but yet which have extended through two or three great periods, we must likewise accord to their latest representatives a direct descent from the older. The very fact that the ocean basins date back to the earliest geological periods, and have afforded to the marine animals the conditions most favourable to an unbroken continuity under slightly varying circumstances, probably accounts for the great range in time during which many genera of Echini have existed. If we examine the interlacing in the succession of the genera characteristic of later geological epochs, we find it an impossibility to deny their continuity from the time of the Lias to the present day. The Cidaris of the Lias and the Rhabdocidaris of the Jura are the ancestors of the Cidaris of to-day. The Salenidæ of the lower Chalk are those of the Salenidæ of to-day. *Acrosalenia* extends from the Lias to the lower Cretaceous, with a number of recent genera, which begin at the Eocene. The *Pygaster* of to-day dates back to the Lias; *Echinocyamus* and *Fibularia* commence with the Chalk. *Pyrina* extends from the lower Jura through the Eocene. The *Echinobrissus* of to-day dates back to the Jura. *Holaster* lived from the lower Chalk to the Miocene, and the *Hemiaster* of to-day cannot be distinguished from the *Hemiaster* of the lower Cretaceous.

Such descent we can trace, and trace as confidently as we trace a part of the population of North America of to-day as the descendants of some portion of the population of the beginning of this century. [But we can go no further with confidence, and bold indeed would be who would attempt even in a single State to trace the genealogy of the inhabitants from those of ten years before. We had better acknowledge our inability to go beyond a certain point; anything beyond the general parallelism I have attempted to trace, which in no way invalidates the other proposition, we must recognise as hopeless.

But in spite of the limits which have been assigned to this general parallelism, it still remains an all-essential factor in elucidating the history of palæontological development, and its importance has but recently been fully appreciated. For, while the fossil remains may give us a strong presumptive evidence of the gradual passage of one type to another, we can only imagine this modification to take place by a process similar to that which brings about the modifications due to different stages of growth,—the former taking place in what may practically be considered as infinite time when compared to the short life history which has given us as it were a *résumé* of the palæontological development. We may well pause to reflect that in the two modes of development we find the same periods of rapid modifications occurring at certain stages of growth or of historic development, repeating in a different direction the same phases. Does it then pass the limits of analogy to assume that the changes we see taking place under our own eyes in a comparatively short space of time,—changes which extend from stages representing perhaps the original type of the group to their most complicated structures,—may, perhaps, in the larger field of palæontological development, not have required the infinite time we are in the habit of asking for them?

Palæontologists have not been slow in following out this suggestive track, and those who have been anatomists and embryologists besides have not only entered into most interesting speculations regarding the origin of certain groups, but they have carried on the process still further, and have given us genealogical trees where we may, in the twigs and branches and main limbs and trunk, trace the complete filiation of a group as we know it to-day, and as it must theoretically have existed at various times to its very beginning. While we cannot but admire the boldness and ingenuity of these speculations upon genetic connection so recklessly launched during the last fifteen years, we find that with but few exceptions there is little to recommend in reconstructions which shoot so wide of the facts as far as they are known, and seem so readily to ignore them. The moment we leave out of sight the actual succession of the fossils and the ascertainable facts of postembryonic development, to reconstruct our genealogy, we are building in the air. Ordinarily, the twigs of any genealogical tree have only a semblance of truth; they lead us to branchlets having but a slight trace of probability, to branches where the imagination plays an important part, to main

limbs where it is finally allowed full play, in order to solve with the trunk, to the satisfaction of the writer at least, the riddle of the origin of the group. It seems hardly credible that a school which boasts for its very creed a belief in nothing which is not warranted by common sense should descend to such trifling.

The time for genealogical trees is passed; its utility can, perhaps, best be shown by a simple calculation, which will point out at a glance what these scientific arboriculturists are attempting. Let us take, for instance, the ten most characteristic features of Echini. The number of possible combinations which can be produced from them is so great that it would take no less than twenty years, at the rate of one new combination a minute for ten hours a day, to pass them in review. Remembering now that each one of these points of structure is itself undergoing constant modifications, we may get some idea of the nature of the problem we are attempting to solve, when seeking to trace the genealogy as understood by the makers of genealogical trees. On the other hand, in spite of the millions of possible combinations which these ten characters may assume when affecting not simply a single combination, but all the combinations which might arise from their extending over several hundred species, we yet find that the combinations which actually exist—those which leave their traces as fossils—fall immensely short of the possible number. We have, as I have stated, not more than twenty-three hundred species actually representing for the Echini the results of these endless combinations. Is it astonishing, therefore, that we should fail to discover the sequence of the genera, even if the genera, as is so often the case, represent, as it were, fixed embryonic stages of some Sea-urchin of the present day? In fact, does not the very history of the fossils themselves show that we cannot expect this? Each fossil species, during its development, must have passed through stages analogous to those gone through by the Echini of the present day. Each one of these stages at every moment represents one of the possible combinations, and those which are actually preserved correspond only to the particular period and the special combination which any Sea-urchin has reached. These stages are the true missing links, which we can no more expect to find preserved than we can expect to find a record of the actual embryonic development of the species of the present day without direct observation at the time. The actual number of species in any one group must always fall far short of the possible number, and for this reason it is out of the question for us to attempt the solution of the problem of derivation, or to hope for any solution beyond one within the most indefinite limits of correctness. If, when we take one of the most limited of the groups of the animal kingdom, we find ourselves engaged in a hopeless task, what must be the prospect should we attack the problem of other classes or groups of the animal kingdom, where the species run into the thousands, while they number only tens in the case we have attempted to follow out? Shall we say "ignorabimus" or "impavidi progrediamus" valiantly to chase a phantom we can never hope to seize?

#### NOTES

THE Second International Geological Congress will be held at Bologna in September, 1881. It is proposed to award a prize of 5,000 francs for the best international scale of colours and conventional signs for the graphic representation of formations on geological maps and sections. Each scale should be accompanied by an explanatory memoir and a sufficient number of maps and sections relative to regions of different geological characters; for the memoirs the French language is recommended. The names of the competitors should be inclosed in sealed envelopes, on which should be a motto. The scales and memoirs should be addressed, before the end of May, 1881, to the President of the Committee, Signor J. Capellini, 65, Via Zamboni, Bologna.

"THE Official Guide and Handbook to Swansea and its District," prepared at the request of the Local Committee by Mr. S. C. Gamwell, is a really useful little work, which must prove of permanent value as a guide to Swansea. It contains much carefully compiled information on the History and Antiquities of Swansea, its literary and scientific, educational

and other institutions, industries, places of interest in and around the town, geology of the district, palæontology, and natural history. A considerable amount of space is appropriately devoted to the scientific aspects of the district, the information given is very full, and we believe trustworthy. An excellent map is prefixed, and the work as a whole is creditable to the Local Committee and to the author.

ON Sunday a statue of Denis Papin was unveiled at Blois, where he was born in 1647.

THE late Mungo Ponton, W.S., Fellow of the Royal Society of Edinburgh, whose death was recently announced, was known as the discoverer of the peculiar effect of light on gelatine when treated with the bichromates, which was afterwards practically applied in the autotype process. He obtained the silver medal from the Royal Society of Edinburgh in 1838 for "model and description of improved electric telegraph." He was the first who employed the photographic method for registering automatically the fluctuations in thermometers and other instruments, and for this service he received also the silver medal of the same society in 1845. Several papers of his appeared in the new *Philosophical Journal* and in the *Quarterly Journal of Science*.

THE Fifth Annual General Meeting of the Mineralogical Society of Great Britain and Ireland was held at Swansea on August 27, Mr. Jas. S. Merry, F.C.S., in the chair. A favourable report was presented by the Council and adopted by the meeting. The election of the following new members was announced:—Dr. Jas. Hector, F.R.S., of New Zealand, Mr. Thos. Stewart of Glasgow, Rev. R. Graham, LL.D., of Errol, Perthshire; Mr. Jos. Gill of Leadhills, and Rev. Geo. Gordon, LL.D., of Elgin. The following papers were read and discussed:—"On the Chemical Formula of Epidote," by M. l'Abbé Renard; "On Certain Crystallised Furnace-products," by Wm. Terrill, F.C.S.; "On the Serpentine and Hornblende, and Schistose Rocks of Porthalla in Cornwall," by II. Collins.

ON the evening of July 20, about half-past eight o'clock, a remarkable meteor, said to have resembled a comet, apparently about twenty yards in length, was observed at Vizimgaum and other places in India, traversing the sky from south to north, and remaining visible for about three-quarters of a minute, during which time the whole sky and country were brilliantly illuminated. The meteor then burst, and some time afterwards a loud sound like distant thunder, which lasted two minutes, was heard.

THE rainfall in Southern China appears to have been abnormally large in the early part of this summer, for we learn from the *Daily Press* of Hongkong that the rainfall in that colony during the month of June was no less than 28'06 inches, compared with 11'32 inches in June, 1879, 15'36 inches in June, 1878, and 9'37 inches in June, 1877. It is stated that so large a rainfall as we have mentioned has never before been registered in Hongkong.

THE Government printer at Brisbane has published three valuable Reports, by Mr. Robert L. Jack, the Geological Surveyor of Northern Queensland, who during the last few years has done excellent service in the cause of both geography and geology in Cape York Peninsula. The first of these Reports deals with the geology and mineral resources of the district between Charters Towers gold-fields and the coast, and is illustrated with a map and several woodcuts taken from photographs, while the second is a preliminary Report on the geological features of part of the coast range between the Dalrymple and Charters Towers roads. The third Report treats of the important Bowen River coal-field. This is accompanied by a map and some large and interesting woodcuts.