


quent want of sharpness on one side, but the definition in the centre of the field is not injured.

To give some idea of the stars that can be seen and the value that may be given to photographic evidence of the existence or non-existence of faint stars, I give some particulars of this photograph. Without a magnifier 124 stars are to be seen within a radius of twenty minutes of arc from the nucleus.

I have traced these (see Fig. 1) so that they can be identified in the telescope; some of them may not be less than 13th magnitude, possibly fainter; the bright stars marked B, C, D, and E being shown in Argelander's maps of the Northern Heavens. B and C are at the present time about the same brightness as the new star, and can be well used to watch any variation in its light (when first seen by me on September 3 the new star was very much brighter than B or C, almost as bright as a star I have called A in my note-book that is just beyond the smaller nebula).

Using a magnifier to detect any fainter stars I find six near the nucleus: these I have shown as black dots on Fig. 2, using a



circle to show the stars near the nucleus that appear on Fig. 1, and a cross (X) to indicate the place of the new star. At this particular place there is not the slightest indication of any difference in the regular shading of the deposited silver from the denser part of the nucleus to the faint edge. The six stars indicated are extremely faint in the photograph and difficult to see, but I have no doubt of their real existence; from a comparison with other photographs I estimate them of about 15th magnitude, perhaps fainter. It may be that some of these may be identified at Birr Castle. From the absence of scale and orientation of the sketch given by Lord Rosse on p. 465 comparisons cannot be made, but a reference to the note-books would enable this to be done.

A. A. COMMON

DURING last week I examined on three evenings the spectrum of this star apparently in the nebula. It appears to be continuous, extending from about D, as far as, or perhaps a little past F. Both Mr. Percy Smith and I are able to confirm Lord Rosse's conviction of the existence of a bright line or band. We compared its position with spark spectra, and feel satisfied that its position is not far from the bright line of the spark in air near to, and on the more refrangible side of D. The slit was of course necessarily wide, and the spectrum faint, so that this must only be considered as approximate.

GEO. M. SEABROKE

Temple Observatory, Rugby, September 29

IN the first evenings of September I observed the new nucleus of the nebula in Andromeda: I find it of the 8th magnitude. With a little Maclean's star spectroscope applied to the 10-inch refractor the spectrum of the nucleus is continuous, with suspected brighter bands. On the nights of Sept. 14 to 16, with 340 and 470 enlargements, I found to the east of the nucleus, and 15" to 18" distant from it, a faint object, probably a second nucleus, of 12th to 13th magnitudes.

A. RICCO

Palermo Observatory

The Proposed Change in the Astronomical Day

IN your account of the proceedings at the recent meeting of the Astronomische Gesellschaft at Geneva (*NATURE*, vol. xxxii. p. 517) Dr. Struve is reported to have stated "that in the

Royal Astronomical Society the majority were in favour of the universal day." There appears to be some mistake here: the Royal Astronomical Society as a body has not expressed any opinion on the subject. And, judging from the individual expressions of opinion which have been published, I should imagine that here, as at Geneva, the majority of real workers in our science (with the probable exception of those engaged on solar work) would be opposed to the proposed change. But how the majority of the Fellows of the Royal Astronomical Society could vote on the question it is impossible to say. My desire that a wrong impression on this subject, arising from a statement reported to have been made by such a high authority as Dr. Struve, should not be spread abroad, must be my excuse for trespassing thus far on your space.

A. M. D. DOWNING

Royal Observatory, Greenwich, S.E., September 26

A Tertiary Rainbow

PROF. TAIT remarks, in his recently-published work on "Light," that rainbows due to three or more internal reflections "are too feeble to be observed." It may therefore be worth recording that a tertiary bow was clearly visible from Thandi Hill, Punjab, one evening last week (August 17). The bow extended over an arc greater than a semicircle, but was broken in two places. The colours were as distinct as in many an ordinary bow.

The condition of the sky was specially favourable for seeing a tertiary bow. The sun was low, and on nearly the same level with it there were several horizontal layers of cloud of considerable extent, whose nearer, unilluminated sides were therefore dark enough to serve as a good background for the bow. There was also a cloud in front of the sun itself, partially reducing its brightness.

T. C. LEWIS

August 25

A White Swallow

ON August 3 I saw a white swallow flying among its fellows over a mill-pond at Garioch's Ford, Auchterless, Aberdeenshire. When I repassed on the following day it was still there, and it appeared to my brother and to me to be *entirely* white: otherwise I should suggest that the one seen in Westmoreland on September 4 (*NATURE*, No. 830, p. 500) might be the same bird on its southward pilgrimage. If it is true that the albino bird is never courted or paired ("Descent of Man," chap. xiv.) we are not likely ever to see many white swallows.

Mirfield, Yorks, September 28

ALEX. ANDERSON

THE enclosed paragraph from Yarmouth, in the *Norfolk News* of this day, will have interest for your correspondent at Milnethorpe.

HUBERT AIRY

Stoke House, Woodbridge, September 26

Rara Avis.—A cream-coloured specimen of the swallow (*Hirundo urbica*) was shot on Caister Road, on Monday morning last, by Mr. A. Patterson. It is now in the hands of Mr. B. Dye of Row 60 for preservation.

DURING the summer of 1883 Mr. Cooper, of Bromwich, observed a white swallow throughout the season at a place within the city on the banks of the Severn.

J. LL. BOZWARD

Worcester, September 28

THE ANNUAL CONGRESS OF THE SANITARY INSTITUTE OF GREAT BRITAIN

THE subjects dealt with by the Sanitary Institute of Great Britain at its annual meetings cover a wide field, and the Leicester gathering of this year, under the presidency of Prof. de Chaumont, F.R.S., forms no exception to the rule. The first aim of the Institute is, through its various agencies, to assist and indeed to lead in the improvement of public health, and the President did well to prove, by mortality statistics, how great a saving of life can be effected by the adoption of efficient sanitary measures, and how remunerative expenditure in this direction proves itself to be. The result of the sanitation carried out in the Army, and which is so much due