

Trustees felt themselves justified in making the transfer. Although this collection is now at the University, its ownership remains with the Trustees.

9. There has been no change in the Board during the year, by death or otherwise.

10. Annexed to this Report are the following Appendices:—

- I.—Annual Balance-sheet.
- II.—Attendance of Visitors.
- III.—Attendance of the Trustees.
- IV.—Work done by Taxidermist and Articulator.
- V.—Specimens collected.
- VI.—Specimens purchased.
- VII.—Exchanges.
- VIII.—Donations.
- IX.—Books acquired.
- X.—Duplicate Books.
- XI.—Mr. Ramsay's Report.

(Signed) ALFRED STEPHEN,
Crown Trustee and Chairman

UNIVERSITY AND EDUCATIONAL INTELLIGENCE

CAMBRIDGE.—The Special Board for Medicine have presented to the Vice-Chancellor the following Report with a view to its communication to the Senate:—"The Board have considered the requirements of the Previous Examination from the point of view of its suitability as a preliminary examination for students entering on the study of medicine, and have come to the conclusion that in the interests of mental training these requirements may with advantage be modified. They would desire to see introduced an adequate examination in the elementary mechanical principles of Physics, meaning thereby—the fundamental notions of matter, motion, and energy, and the simple laws which govern their relations; the physical properties of matter in the solid, liquid, and gaseous states; and the application of these properties and laws in the case of simple instruments and machines. An examination in these principles need not involve any but the most elementary mathematics, yet it could be made to exercise the student in clearness of conception, in accuracy of statement, and in soundness of reasoning. These qualities are in a special degree essential to students of medicine, but from our Report of November 11, 1885, it would appear that in these respects the preliminary training of many who propose to become students of medicine has not been satisfactory. The subject we propose is already well taught and appreciated in many good schools, and it appears to us extremely desirable that the University should encourage all schools to improve themselves in this direction by including the subject in its Previous Examination. It is not for the Board to say whether the subject should form part of the Previous Examination proper (though many considerations might be urged for this plan), or be required as an additional subject in place of the present examination in Elementary Mechanics. They are, however, persuaded that, if introduced in some form, the examination would be for all students at least of equal value to the present examination in additional subjects, and for students whose work at the University is to consist largely in the study of nature it would be of considerably greater value."

Mr. H. D. Rolleston, of St. John's College, has been appointed Assistant Demonstrator of Physiology, in succession to Mr. Green. Mr. Rolleston was placed in the First Class in the Natural Sciences Tripos, Part I., in the Easter Term, 1885.

SCIENTIFIC SERIALS

Annalen der Physik und Chemie, No. 12, December 1885.—J. Fink, on the influence of pressure on the electric resistance of electrolytes. Cailletet's apparatus was used for producing compression, Kohlrausch's induction apparatus for the electric measurements. A solution of hydrochloric acid (5.02 per cent.), having a resistance of 7.490 Siemens' units at 1 atmo., fell to 7.335 at 200, and to 7.126 at 500 atmo. A weaker solution (0.98 per cent.) showed a diminution of 7.39 per cent. in its resistance at 500 atmo. A similar solution of zinc sulphate showed a diminution of 11.74 per cent. The diminution is

proportional up to 300 atmo.—E. Edlund, on the transition-resistance in the voltaic arc. The conclusion is against the existence of such a resistance.—K. Wesendonck, on the fluorescence of naphthalin-red.—H. W. Vogel, on the relation between absorption by colouring matters and their sensitising action on bromide of silver.—G. Kötschau, studies on fluid motions. Some very extraordinary figures are produced by careful introduction of a coloured liquid into an uncoloured one.—F. Himstedt, a determination of the ohm. This paper describes the method, depending on a knowledge of the coefficient of mutual induction of two coils, which has already been discussed by Lord Rayleigh, and which is similar to that of Roiti. The final result gives as equivalent to the ohm a column of mercury of 1 square millimetre section and 105.98 centimetres length.—W. B. Brace, on the magnetic rotation of the plane of polarisation, and some special cases of refraction. It is shown that there may be in a calc-spar crystal three rays which suffer no double refraction. Experiments are also described concerning prisms of heavy glass in a magnetic field.—G. Stern, position of the commutator in electro-dynamic machines. A discussion of Clausius' formulæ with respect to the relation of the current to the angle of lead.—E. Mach and J. Wenzel, a contribution to the mechanics of explosions.—K. L. Bauer, apparatus for demonstrating that electricity resides only on the surface of a conductor. This is a modification of Biot's apparatus, consisting of two concentric hemispheres, and convenient means of insulating and discharging.

Journal of the Russian Chemical and Physical Society, vol. xvii. fasc. 7.—On the part played by contact actions in the phenomena of dissociation, by D. Konovaloff.—Thermic data for some combinations of the aromatic series, by E. Werner, being numerical data as to the heat of neutralisation of saligenin and oxybenzoic aldehydes and acids, and mellic acid.—On the oxidation of oleic and elaidic acids by permanganate of potassium, by A. Saytzeff.—Notes by MM. Albitzky, Nikolsky, and Ustinoff.—On the motion of a solid body having cavities filled with a homogeneous liquid, by M. Joukowsky, being the second part of a mathematical inquiry into ellipsoidal, cylindrical, and such other cavities as have the shape of a rotation-body, and also several cavities connected together.—On the collision of absolutely solid bodies, by M. Schiller, second part, being a further mathematical development of the theory, together with answers to Prof. Joukowsky's observations.—On the influence of an electric current on the resistance of selenium and its sensibility to light, by N. Hesehus, being an explanation of the experiments of Fritts by the theory of allotropic dissociation.

SOCIETIES AND ACADEMIES

LONDON

Royal Society, December 17, 1885.—"On the Formation of Vortex-Rings by Drops falling into Liquids." By Prof. J. J. Thomson, M.A., F.R.S., and H. F. Newall, M.A.

When a drop of ink falls into water from not too great a height, it descends through the water as a ring, in which there is considerable rotation about the circular axis passing through the centres of its cross-sections; as the ring travels downwards, inequalities appear, and the ring breaks up into a number of smaller rings, which in turn may again subdivide.

It is shown that capillarity plays no essential part in the formation of the rings; in fact, it may be said that, with very few exceptions, rings are formed only when a liquid is dropped into one with which it can thoroughly mix. There are very many cases in which rings are formed when there is no possibility of capillary action, such as when the liquid into which the drop falls is the same as the drop itself.

The drops were observed by instantaneous illumination; and it was seen that the drop enters the liquid as a sphere, becomes flattened as it descends, and finally breaks into a ring more than half an inch below the surface.

When a sphere moves through a liquid, the tangential velocity of the liquid is different from that of the sphere. If the sphere be a liquid drop, there is no absolute discontinuity in the motion, but only a very rapid change, so that there is a finite alteration in a very small distance. This is equivalent to a vortex-film covering the sphere, the lines of vortex-motion being horizontal circles. If the liquid be viscous, the vorticity will diffuse inwards and outwards. The drop, as it falls, becomes flattened, on account of the resistance to its fall; and if by