

Meteor-Showers

	R.A.	Decl.
Near ν Herculis	236	47° N.
η Ophiuchi	254	21 S.
ζ Draconis	262	64 N.
α Aquilæ	298	5 N.

GEOGRAPHICAL NOTES

WE are glad to know that General R. Strachey, R.E., F.R.S., has agreed to accept the presidency of the Royal Geographical Society, in succession to Lord Aberdare. The Society's honours for the present year will be conferred as follows:—The Founder's Medal to Lieut.-Colonel T. H. Holdich, R.E., for the eminent services he has rendered to geography in Afghanistan; the Patron's Medal to the Rev. George Grenfell, for the extensive explorations he has carried out during his thirteen years' residence in Africa, partly in the Cameroons country and more recently in the Congo region; the Murchison Grant to Mr. George Bourne, second in command and sole survivor of the Landsborough Expedition, which crossed the continent of Australia in 1861; the Back Premium to Sarat Chandra Das, for his researches in Tibet; the Gill Memorial to Mr. J. F. Needham, for his explorations in the Lohit Valley of the Brahmaputra. The following have been made Honorary Corresponding Members:—H.R.H. Krom Mun Damrong Rajah Nubhar, Director-General of Surveys and Minister of Public Instruction, Siam; Dr. Alfred Kirchhoff, Professor of Geography at Hallé University, and President of the Hallé Geographical Society; and Dr. E. Naumann, late Director of the Geographical and Topographical Survey of Japan.

THE paper read at the Royal Geographical Society on Monday was by General J. T. Walker, F.R.S., on the Lu River of Tibet, the Lu-Kiang, or Lu-tse-Kiang of the Chinese. This river is generally held to be the source of the Salwin, but General Walker adduced many reasons for maintaining that it is more probably that of the Irrawadi. In the course of an able paper, the result of much research, General Walker gave a most useful summary of exploration in this highly interesting hydrographical region; more particularly insisting on the value of the work of the late Abbé Krick, who ascended the Lohit in 1852, but of whom little is known in this country. General Walker made out a strong case for his position, but the leading conclusion of his valuable paper is that further exploration in this remarkable region is urgently demanded. Probably no region of the earth would yield more valuable results to scientific geography.

ACCORDING to the latest news, Mr. Stanley is well up to time in his ascent of the Congo with the Emin Pasha Expedition. He is at present on his march across country from Matadi, at the lower end of the Livingstone Falls, to Leopoldville, on Stanley Pool. It is hoped that when he reaches the Pool he will find sufficient vessels in readiness to convey his large following up the river without delay. So far the Expedition has been exceedingly fortunate.

THE statement that Baron Nordenskjöld will undertake an Antarctic expedition at the expense of the King of Sweden and Mr. Oscar Dickson is, to say the least, premature. We are informed by Mr. Dickson that Baron Nordenskjöld is "willing" to undertake such an expedition, but that if he does so neither the King of Sweden nor Mr. Dickson will find the money. No doubt Baron Nordenskjöld would be an excellent leader for such an expedition, and as Committees have been formed both in this country and in Australia to promote Antarctic exploration, would it not be wise in them to unite their forces, and place themselves in communication with the Baron? Those who are competent to give an opinion on the subject maintain that an Antarctic expedition is much less risky than one to the other Pole. There would be no difficulty in a party wintering on some part of the Antarctic continent; a vessel could cruise round the verge of the ice during the winter and watch a favourable opening, of which immediate notice could be given to the exploring party, while a third vessel could leave New Zealand at a suitable time with additional supplies. No doubt the subject will again be brought up at the next meeting of the British Association, when it is hoped a strong and active Committee will be appointed. Baron Nordenskjöld will be among the distinguished foreigners invited to the meeting, and we hope he will accept the invitation.

THE Germans continue to show great activity in the exploration of their portion of New Guinea. Freiherr von Schleinitz has recently accomplished a running survey of Huon Gulf, and besides establishing the direction of the coast-line and the positions of reefs, has laid down eight hitherto unknown harbours and discovered nine new rivers. Some of them, especially the Markham River, would form excellent routes for the exploration of the interior; the broad valley of the latter extends for miles between high ranges of mountains. The south coast of Huon Gulf consists exclusively of primitive and metamorphic rocks, with older sedimentary rocks and volcanic formations. At a later date a further survey was made of the coast from Astrolabe Bay to the mouth of the Empress Augusta River, and led to the discovery of a series of bays, harbours, islands, and rivers.

M. GRIMAILO, in company with his brother, an engineer, and six Cossacks, has set out for a further exploration of the Pamir.

DR. LABONNE left Cherbourg a few days ago for a further exploration of the geysers and glaciers of Iceland.

PRELIMINARY NOTE ON THE FOSSIL REMAINS OF A CHELONIAN REPTILE, CERATOCHELYS STHENURUS, FROM LORD HOWE'S ISLAND, AUSTRALIA¹

THE interesting remains of which I propose to give a brief notice in the present communication are contained in a friable sandstone (apparently formed of concreted blown sand), and they have a very recent appearance. The age of the deposit in which they are found is unknown, but it is probably Quaternary. The specimens have been for some years in the palæontological collection of the British Museum, and, for the most part, they have not yet been submitted to careful examination. But I learn that the greater number of them were long since rightly determined to be Chelonian by Mr. Davis, and set aside as such.

Several of the most important of these numerous and, in general, very fragmentary bones were originally found imbedded close together in the same block of sandstone. They consist of a great part of a pelvis, a caudal vertebra, and an imperfect skull. Of the pelvis, a right ischium and a pubis are imbedded in the rock, while an imperfect right ilium, which fits well on to the ischium, is separate; all these bones are unmistakably Chelonian. The caudal vertebra has remarkable peculiarities. It resembles an ordinary Chelonian caudal vertebra from the anterior half of the tail, in its general characters; but it is strongly opisthocœlous, the centrum having a deep cup behind and a correspondingly curved articular head in front. From the posterior part of the ventral face two stout processes diverge, and present terminal rounded facets for the rami of the large chevron bone which must have articulated with them. As a general rule, the caudal vertebræ of Chelonia are procœlous—but *Chelydra* and *Gypochelys* (perhaps also *Staurotyphus* and *Platysternum*) form well-known exceptions,² in so far as the vertebræ behind the third and fourth are strongly opisthocœlous. In fact, the vertebra in question closely resembles the sixth or seventh of *Chelydra* or of *Gypochelys* (see Figs. 1 and 2). In the first, however, the transverse processes are very much stronger, and the pentagonal platform into which the upper surface of the neural arch expands, in place of a neural spine, is as long as the vertebra, instead of being only about half as long. The stout pre-zygapophysis of the right side is broken off, leaving only the base visible in the fossil.

¹ Paper read at the Royal Society, by Prof. Thomas H. Huxley, F.R.S., on March 31.

² The opisthocœlous character of most of the caudal vertebræ of *Chelydra* was first pointed out by Von Meyer in his description of the Eningen *Chelydra*. Baur ("Osteologische Notizen," *Zool. Anzeiger*, No. 238, 1886) has gone fully into the question, and has pointed out the exceptional nature of their structure among the Chelonia. Since the above paragraph was written, Dr. Günther has kindly enabled me to examine a spirit specimen and a skeleton of *Platysternum*. The caudal vertebræ resemble those of *Chelydra*, except that the last nine are procœlous, while that between these and the more anterior opisthocœlous vertebræ is nearly flat at the ends. In this, as in other respects, *Platysternum* presents characters intermediate between *Chelydra* and the ordinary *Emyda*. Prof. Cope ("Vertebrata of the Tertiary Formations of the West," 1883, p. 111) ascribes opisthocœlous caudal vertebræ to the *Bœnida*, but no figures or descriptions of such vertebræ are given. Of the opisthocœlous Chelonian vertebræ figured in Plate XXIV. of the "Report of Extinct Vertebrata obtained in New Mexico" (1887), it is expressly stated that their "correct reference cannot now be made" (p. 43).

Two other caudal vertebræ, having the same structural features, occur among the detached remains, and belong, like the first, to the second fourth of the tail. Another tolerably complete vertebra, with a considerably longer centrum, corresponds very closely with a caudal vertebra of *Gypochelys* from the third fourth of the tail. In this, as in one of the foregoing vertebræ, the chevron bones are ankylosed with the centrum. I conceive, then, that there can be no doubt that the pelvic bones and these caudal vertebræ belonged to a Chelydroid Chelonian, of about the size of the largest "snapping turtles" which are met with in North America at the present day.

Primâ facie, the skull found in the same block might also be expected to be that of a Chelydroid; and, in fact, it is so. I do not base this interpretation on the Chelonian character of the

upper jaw, as there are various extinct Saurian reptiles which closely approximate to Chelonia in this part of their structure. The diagnostic characters lie in the back part of the skull; and especially in the auditory region, which is altogether Chelonian. Not only so, but when this fragmentary skull is compared with that of *Chelydra*, the correspondence between the two is singularly exact (Figs. 3 and 4). In two respects, however, the fossil differs from *Chelydra* and *Gypochelys*.

(1) The roof over the temporal fossa formed by the parietal, post-frontal, and other bones, which leaves the auditory region uncovered in the recent genera,¹ extends back, beyond the occiput, in the fossil, and sends down a broad vertical rim from its margin.

(2) The upper surface of the cranial shield is, at most, rugose




FIG. 1.—Caudal vertebra of *Ceratochelys*. *N*, platform on the neural arch; *p3*, pre-zygapophysis mutilated; *tr*, broken transverse process; *Chv'*, processes for the chevron bone; *Chv*, chevron bone.




FIG. 3.




FIG. 2.—Caudal vertebra of *Chelydra*. Letters as in Fig. 1.




FIG. 4.

FIGS. 3, 4.—Skulls of *Ceratochelys* (Fig. 3) and *Chelydra* (Fig. 4); the latter of the natural size, the former much reduced. The portion of the skull of *Chelydra* which corresponds with the fossil is shaded.

in the recent *Chelydridæ*; in the fossil, three strong conical processes, like horn-cores, of which the middle is the longest, are developed from its posterior and lateral region.¹

This skull is described and figured in the Philosophical Transactions for 1886 (Plate 30, Fig. 1) by Sir R. Owen, under the generic or sub-generic name of *Meiolania*, and is said to belong to a Saurian reptile closely allied to the "*Megalania prisca*" described in earlier communications. But the skull is assuredly that of the Chelydroid Chelonian to which the pelvis and caudal vertebra belong. What *Megalania prisca* may be I do not pretend to say; but the remains which I have described can have nothing to do with any Saurian reptiles; and I propose to confer


¹ It is possible that these may be dermal bones coherent with the proper cranial shield.

on the genus of Chelonia to which they belong the name of *Ceratochelys*.

The singular osseous caudal sheaths described by Sir R. Owen, in the same memoir, also appertain to *Ceratochelys*. They formed part of the series of remains sent to the British Museum along with the foregoing, in which none but Chelonian bones have yet been discovered; and the remains of vertebræ left in these sheaths are similar to the caudal vertebræ of the terminal fourth of the tail in the *Chelydridæ*. The snapping turtles are noted for the length and strength of the tail and for the strong, laterally-compressed, acuminated "scales" which form a crest along the median dorsal line, while others, less strongly keeled, lie at the sides of the tail. In many Chelonia, the extremity of

² The "roof" extends much further back in *Platysternum*.

the tail is enveloped in a continuous sheath. These and other scale-like structures in the Chelonia are usually spoken of as if they were entirely epidermal. But, a day or two ago, Dr. Günther informed me that in the Australian tortoise, *Manouria*, the great imbricated scales of the limbs contain bony scutes; and that similar scutes are to be found in *Testudo graeca*. This, of course, suggested the examination of the caudal scales of *Chelydra* and *Gypochelys*; and, having been enabled by Dr. Günther's kindness to examine the caudal scales of a good-sized specimen of the latter, I have found that those of the crest contain bony scutes.¹ The bony scute corresponds very closely in form with the whole "scale," but the recurved apex of the latter is formed only by epidermal substance (Figs. 5 and 6).



FIGS. 5, 6.—Sectional views of a scute of the tail-armour of *Ceratochelys* (Fig. 5), and of one of the crest plates of *Gypochelys*, both of the natural size.

The living *Chelydra*, therefore, has a caudal armature which, in principle, is similar to that of *Ceratochelys*, but the osseous elements are relatively atrophied. There is exactly the same relation between the armour of species of living *Crocodyles* and *Alligators*, on the one hand, and those of *Facare* and *Caiman* and the extinct *Teleosauria*, on the other. In the former, the epidermal scales remain well developed on the ventral side of the body, while the corresponding osseous scutes, fully developed in *Facare*, *Caiman*, and *Teleosauria*, have vanished.

Among the detached fragments to which I have referred, there are remains of ribs, with their costal plates; marginal and other plates of the carapace; parts of the plastron; part of a scapula; sundry limb bones; and several of the cranial processes called "horn-cores." They all agree, so far as they can be compared, with the determination already arrived at; which, to sum it up in a few words, is that the remains of crania and caudal sheaths from Australia, hitherto referred to Saurian reptiles, under the names of *Megalania* and *Meiolania*, appertain to a hitherto unknown species of Chelonian, *Ceratochelys sthenurus*, closely allied to the living *Chelydra*, *Gypochelys*, and *Platysternum*.

The evidence of this fact offered in the present note appears to me to be conclusive, but it may be desirable hereafter to figure the parts mentioned and to describe them at length.

The interest which attaches to the discovery of this singular Chelonian arises partly from the fact that the group of Chelonia to which it belongs is wholly unrepresented in the fauna of Australia, as at present known. *Platysternum* is usually said to be found in China. Dr. Günther, however, informs me that Upper Burmah is its proper habitat; otherwise, North America, east of the Rocky Mountains, is the nearest region in which the *Chelydridæ* are to be found. But *Chelydridæ*, and, indeed, species of the genus *Chelydra*, occur in Upper Miocene (Eningen) and in Eocene formations in Europe. Moreover, *Platy-chelys*, of the Upper Jurassic series of Bavaria and Switzerland, is regarded by Rüttimeyer as an early form of the group.

Lord Howe's Island is about 200 miles from the nearest Australian mainland, and something like 400 miles, as the crow flies, from the Darling Downs, in which the caudal armour, which has been ascribed to *Megalania*, was found. The discovery of *Ceratochelys*, therefore, has an interesting bearing on

¹ The fact is noted by Rüttimeyer (Lang and Rüttimeyer, "Die Fossilen Schildkröten von Solothurn," *Denkschriften der Allg. Schweiz. Gesellschaft*, vol. xxii.). The armature of the tail in *Platysternum* is for the most part arranged in zones, of four plates in each zone; but I have not yet been able to find any bone in them.

the question of the former extension of Australia to the eastward, on the one hand; and of the possible derivation of such forms as *Ceratochelys* from Asia, on the other hand. An elevation of the sea-bottom of 6000 feet would place Norfolk Island and Lord Howe's Island on a peninsula extending from the region of the present Barrier Reef to New Zealand; and the floræ and faunæ of those islands are known to have special affinities with those of New Zealand, and none with those of Australia.

Speculations respecting the origin of the Chelonian carapace are suggested by the discovery of osseous scutes in the vertebral region of the tail, and their coalescence in *Ceratochelys* to form a sort of caudal carapace, ridged in a manner resembling that of *Chelydra* and *Platy-chelys*. But the consideration of these points would take me beyond the limits of the present note.

THE WORK OF THE IMPERIAL INSTITUTE¹

I.

THE Colonial and Indian Exhibition, which owes not only its conception, but also its brilliantly successful realisation to your Royal Highness, will be pre-eminently remarkable in times to come, for having achieved many results of vital importance and highest benefit to Her Majesty's subjects in all parts of her vast realms.

The collection of all that is commercially valuable and scientifically interesting of the natural products of the great Indian Empire and of the Colonies in one Exhibition, embracing as it also did very comprehensive illustrations of the development of commerce, of the arts, and of certain industries, in the many countries beyond the seas which combine with the United Kingdom to constitute our vast Empire, afforded those at home an opportunity, surpassing all previous conception, of studying and comparing the natural history and resources of those distant lands, of which, attached though we might be individually to one or more of them by ties of friendship or of interest, the knowledge of many of us was of a very vague or partial character.

To the Colonists who visited us last year, the Exhibition has been of inestimable value, in affording them a most favourable and appropriate opportunity of becoming acquainted or renewing their old friendship with the mother country, and of examining the progress there made in industrial, educational, and commercial development; in leading to the cultivation of intimacy between Colonists from different sections of the Queen's Dominions; and in affording them invaluable opportunities of comparing the resources and state of development of their respective countries with those of other parts of Europe. No more convincing illustrations than were provided by this great Exhibition could have been conceived of the importance to the home country, to each Colony, and to India, of fostering intimate relationship and unity of action. No more encouraging proof could have been afforded of the desire of all classes of Her Majesty's subjects at home to cultivate a knowledge of those far-off countries which the enterprise and perseverance of the British, and men of British offspring, have converted into prosperous and important dominions, chiefly during the period of the Queen's reign, than was furnished by the interest which the thousands upon thousands, who came from all parts, displayed in the study of the instructive collections in the galleries at South Kensington.

It was the success of the Exhibition which led to the definite formulation of the suggestion first made by your Royal Highness in a letter addressed by you in the autumn of 1884 to the Agents-General of the Colonial Governments, that a permanent representation of the resources of the Colonies and India, and of their continually progressing development, might, with great benefit to the Empire at large, be established in this country. That the realisation of this idea upon a sufficiently comprehensive basis might constitute a worthy memorial of the accomplishment of fifty years of a wise and prosperous reign; a memorial not personal in its character excepting so far as it constituted an emblem of the love and loyalty of Her Majesty's subjects, but tending, as she would most desire, to serve the interests of the entire Empire; this had only to be pointed out by your Royal Highness to be heartily concurred in by the official representatives of the Colonies and India, who

¹ Lecture (abridged) delivered at the Royal Institution, on Friday, April 22, by Sir Frederick Abel, C.B., F.R.S.; H.R.H. the Prince of Wales, K.G., F.R.S., Vice-Patron, in the Chair.