

was read by Mr. George C. Thomson on "Smoke," and in the discussions which followed, Mr. W. R. W. Smith, Chairman of the Health Committee of the Glasgow Corporation, urged upon the members present the desirability of doing all in their power to secure that at the forthcoming International Exhibition in Glasgow each of the boilers be supplied with a separate chimney, so that a series of exhaustive trials may be made with mechanical stokers, &c., and other means for the purpose of showing what might be done in the way of smoke prevention.

With reference to the subject of testing, the Committee are of opinion that arrangements should be made as soon as possible for obtaining the use of three testing-rooms for testing stoves, grates, and ranges, the rooms being conveniently accessible for such articles, and having gas connections under command. The tests made in these rooms, under the same conditions of chimney and cubic capacity, would then become of greater comparative value than tests made in independent rooms.

Arrangements will be made as soon as practicable for procuring such accommodation for testing, and also for providing the necessary instruments used for testing; and as the system develops, attention will be given to the establishment of a chemical laboratory, the analysis of gases, and testing-rooms for testing-apparatus incidental to the work of the Institution.

SCIENTIFIC SERIALS.

American Journal of Science, December 1887.—On the destruction of the passivity of iron in nitric acid by magnetization, by Edward L. Nichols and W. S. Franklin. From the experiments described in this paper, which was originally read before the Kansas Academy of Science, November 1885, it appears that the action of the magnet tends to lower the temperature of transition to the active state, and that the intensity of the magnetic field necessary to convert passive into active iron at a given temperature increases rapidly with the concentration of the acid. An account is promised of further researches offering a satisfactory explanation of the manner in which the chemical behaviour of iron is modified, and its passivity destroyed in the magnetic field.—On a method of making the wave-length of sodium light the actual and practical standard of length, by Albert A. Michelson and Edward W. Morley. The preliminary experiments recently carried out according to the method here proposed seem to confirm the anticipation that it would furnish results more accurate than any of those hitherto suggested. The apparatus for observing the interference phenomena is the same as that used in the experiments on the relative motion of the earth and the luminiferous ether.—The work of the International Congress of Geologists, by G. K. Gilbert. This is a reprint of an address delivered before the Section of Geology and Geography of the American Association for the Advancement of Science at the New York meeting, August 10, 1887. It deals largely with a revised system of geological terminology, the substance of which has already been published. The question of geological coloured maps is also considered, and practical suggestions made for their greater efficiency and economy.—On the existence of certain elements together with the discovery of platinum in the sun: contributions from the physical laboratory of Harvard University, by C. C. Hutchins and E. L. Holden. These investigations, carried on with Prof. Rowland's magnificent diffraction grating, deal with cadmium, lead, tin, silver, potassium, and several other elements, including platinum, the presence of which in the solar atmosphere is here for the first time determined. Between 4250 and 4950 were found sixty-four lines of platinum, sixteen of which agree with the solar lines.—The flora of the coast islands of California in relation to recent changes of physical geography, by Joseph Le Conte. A careful study of these insular groups, at present from 20 to 30 miles distant from the coast, shows that they at one time formed part of the mainland, from which they were undoubtedly separated during the Quaternary period. That they still formed part of the continent during later Pliocene times is shown by the remains of the mammoth found on Santa Rosa, one of the largest and furthest off of the whole group.—A new instrument for the measurement of radiation, by C. C. Hutchins. The instrument here described and illustrated presents great advantages over the thermopile as an accurate measurer of radiations. It is much more sensitive and requires

no longer time to return to zero than for the galvanometer needle to come to rest. A lighted match at 6 feet drives the needle round to its stop.—Mineralogical notes, by George F. Kunz. Descriptions with analyses are given of a rhodochrosite from Colorado, of crystals of hollow quartz from Arizona, of hydrophane from Colorado, and of a remarkable silver nugget weighing 606 ounces from the Greenwood mines of Michoacan, Mexico.

January.—The speed of propagation of the Charleston earthquake, by Prof. Simon Newcomb and Captain C. E. Dutton. A careful comparative study of the reports from all parts of the disturbed area shows a general average speed of $3'214 \pm 0'072$ miles, or 5171 ± 116 metres per second.—History of the changes in the Mount Loa crater, Hawaii; Part I, Kilauea, by James D. Dana. The first paper embraces the whole period from 1823 to 1886, during which there appear to have been at least eight discharges from Kilauea. The general dynamical conclusions are that the cycle of movement is simply (1) a rising in level of the liquid lavas, and of the bottom of the crater; (2) a discharge of the accumulated lavas down to some level in the conduit determined by the outbreak; (3) a down-plunge of more or less of the floor of the region undetermined by the discharge. It is further shown that Kilauea is a true basalt volcano in its normal state, the rock material being dolerite or basalt, and the heat sufficing for the perfect mobility of the lavas.—The analysis and composition of tourmaline, by R. B. Riggs. The methods of analysis are described, with results for various specimens from different parts of North America and Brazil. The general inference is that there are three types, lithia, iron, and magnesia tourmaline, with an indefinite number of intermediate varieties, iron appearing to be the connecting link between the whole series. The special formulas of the three distinct types are:—

- (1) Lithia : $12\text{SiO}_2, 3\text{B}_2\text{O}_3, 4\text{H}_2\text{O}, 8\text{Al}_2\text{O}_3, 2(\text{NaLi})_2\text{O}$.
- (2) Iron : $12\text{SiO}_2, 3\text{B}_2\text{O}_3, 4\text{H}_2\text{O}, 7\text{Al}_2\text{O}_3, 4\text{FeO}, \text{Na}_2\text{O}$.
- (3) Magnesia : $12\text{SiO}_2, 3\text{B}_2\text{O}_3, 4\text{H}_2\text{O}, 5\text{Al}_2\text{O}_3, \frac{2}{3}\text{MgO}, \frac{2}{3}\text{Na}_2\text{O}$.

—On the different types of the Devonian system in North America, by Henry S. Williams. It is shown that in North America the Devonian system offers at least four distinct types in four corresponding areas, blending somewhat at their borders, but in their central parts presenting marked peculiarities. The four areas are: (1) Eastern Border, mainly in Northern New England; (2) Eastern Continental, including New York, thence southwards to West Virginia and north-westwards to Canada West and Michigan; (3) Interior Continental, chiefly Iowa and Missouri, extending northwards probably to the Mackenzie basin; (4) Western Continental, in Nevada and conterminous States.—On the law of double refraction in Iceland spar, by Charles S. Hastings. The general inference from these researches is that Huyghens' law of double refraction in uniaxial crystals is probably true to less than 1 part in 500,000, and consequently that there is no known method by which any error in it can be detected by observation alone.—In the Appendix, Mr. O. C. Marsh describes a new genus of *Sauropoda* and other new Dinosaurs from the Potomac formation; also a new fossil Sirenian from California.

SOCIETIES AND ACADEMIES.

LONDON.

Royal Society, January 19.—"Notes on the Spectrum of the Aurora." By J. Norman Lockyer, F.R.S.

I exhibited to the Society on November 17, 1887, a tabular statement showing the bright lines seen in the spectra of various celestial bodies, and I also gave those recorded in the spectrum of the aurora, showing many remarkable coincidences.

I now find that the connection is closest between the auroral spectrum and that of stars III.a, and, in anticipation of a subsequent communication of details, I send on the accompanying table, showing the origin of Dunér's bands, so far as I have at present made them out, and their connection with the spectrum in question.

The individual observations which I have used in the table are those collected by Mr. Capron and Mr. Backhouse (*NATURE*, vol. vii. pp. 182 and 463).