Cloud Electric Potential.

UNDER the above heading, in NATURE of July 19 (p. 269), which has just come into my hands, Mr. E. Douglas Archibald criticizes a statement of mine in Part III. of "Deschanel," respecting electrified drops of water in a cloud. The following is the statement:—

"The coalescence of small drops to form large ones, though it increases the electrical density on the surfaces of the drops, does not increase the total quantity, and therefore cannot directly influence the observed potential."

At the word "therefore" I give a reference to a previous section, in which it is shown that the potential at a point is the sum of all the quotients q/r , q denoting an element of the electricity to which the potential is due, and r the distance of this element from the point in question. Since the coalescence of drops is without effect on the value of each q and its corresponding r , it cannot affect any one of the quotients q/r , whose sum constitutes the potential.

Mr. Archibald's criticism is:—

"Surely this entirely omits the fact that the capacity of a sphere is equal to its radius, and thus, in the case of eight equal spheres coalescing into one, not merely would the density be doubled, but the potential of the same quantity would be increased four times."

This criticism rests on two false assumptions:—

First, that the potential of a drop depends on its own charge only, and can therefore be computed by dividing its charge by its radius.

Secondly, that the potential of the drops (which on this supposition would be very different from the potential at a point midway between two drops) can be identified with "the observed potential."

J. D. EVERETT.

Cushendall, Co. Antrim, August 3.

THE ABSORPTION SPECTRA OF CRYSTALS.

ALL who are interested in the difficult work now going forward in so many chemical laboratories, in connection with the nature and constitution of those most complex mixtures known as "rare earths," and who recognize the extremely important influence which the solution of this subject must exert upon the very basis of our modern chemistry, will gladly welcome a new and exquisite means of investigation which M. Becquerel has recently brought to light.

As the reward of a most exhaustive study of the changes which are brought about in a beam of light by its passage through a crystal, M. Becquerel has discovered the key by means of which he is enabled to interpret the subtle indications which the issuing rays afford as to the nature of the molecules among which they have been threading their way. It appears at first sight more than wonderful that these delicate indications can have led to precisely the same weighty conclusions as those arrived at from the renowned physico-chemical researches of Auer von Welsbach, Lecoq de Boisbaudran, Demarçay, Sorêt, Crookes, and Krüss and Nilson. Yet such indeed is the case, and it even appears likely that the new method may be carried still further into the region beyond that which has up to the present been reached by these experimenters.

In order to explain the nature of this discovery, it will be necessary to describe the experimental steps which have led M. Becquerel towards it. In the year 1866 Bunsen found what now appears to be the germ of a great principle—that when a crystal of the sulphate of the substance didymium, now known to be a most complex mixture, was traversed by a beam of plane-polarized light vibrating at an angle of 20° to the horizontal diagonal of the crystal, the absorption spectrum was slightly different from that which was obtained when the ray was polarized in a plane at right angles. This observation did not attract much attention at the time, it being considered merely as a curious manifestation of the phenomenon of pleochroism.

Sorby, however, in 1869 again reopened the question, having found that in zircons the ordinary and extra-

ordinary rays presented different bands of absorption. Since that time Becquerel himself has shown that the same applies to all birefractive crystals which give absorption spectra.

With so much premised, we are now in a position to consider the main results of this more recent investigation. They may be very briefly summarized as follows:—

(1) The bands in the absorption spectra of all crystals have fixed positions: the intensity alone varies with the direction of propagation of the light.

(2) In most crystals, the principal directions of absorption coincide with the directions of optical elasticity.

(3) In certain crystals the directions appear to be very different for different bands, but they always remain subject to the conditions imposed by the crystalline symmetry; thus in monoclinic crystals one of the principal axes of absorption always coincides with the axis of symmetry, and the other two rectangular axes are situated in the plane of symmetry.

Hence it appears to be a fact that the absorption of luminous radiations of fixed wave-length admits of three directions of symmetry. These directions appear generally to coincide with the principal directions of optical elasticity, with the exception of certain remarkable anomalies in particular crystals. Here, however, is the whole gist of the matter. Why these anomalies? Just as from a consideration of the deviations from Boyle's law physicists have learned how to measure the size of those wonderfully minute entities familiar to us as molecules, so has M. Becquerel extracted a most important principle out of the anomalies to the law of absorption in crystals.

It appears probable that absorption may be due to a physical phenomenon dependent upon the intermolecular movements. The intimate relation between phosphorescence and absorption, notably in the compounds of uranium and certain of the rare earths, appears to show that in solids and liquids the radiations absorbed are those which vibrate in unison with the molecular movements. This conception is in fact nothing more than an extension to solids and liquids of the well-verified law of the absorption by incandescent vapours.

As the molecular elasticity varies in different directions in crystals not isotropic—that is, not belonging to the cubic system—so will the absorption vary; and if, in two isomorphous substances, the directions of molecular elasticity do not exactly correspond, the directions of different absorption in the two substances will vary in like manner. Now it is quite true that many crystals of isomorphous substances—that is to say, substances of analogous chemical constitution crystallizing in similar forms—have their optic axes unequally inclined.

If we crystallize two such substances together, in gradually increasing proportions of one of them, we find that the angle between the optic axes in the mixed crystals diminishes progressively until it reaches zero, after which the two axes again diverge in a plane perpendicular to their original plane. Thus can we cause the influence of each in turn to preponderate.

Each chemical substance therefore exerts its own influence, and the molecules retain the optical properties which they manifest when the substance crystallizes alone. Hence the propagation of luminous waves is the resultant of the actions which each of the molecules composing the crystal exerts upon the luminous vibrations. If the directions of absorption do not coincide with the axes of optical elasticity, it indicates the presence of molecules of different substances in the crystal. From these considerations it will be evident that the anomalies are probably due to the coexistence in the same crystal of different matters, geometrically isomorphous, but optically unlike, and which from the absorption point of view behave as if each were alone. The use of the spectroscope will therefore enable us to recognize the individuality of differently

absorbing molecules in the molecular groupings, which other optical methods cannot indicate; for the absorption due to one molecule is independent of that of a neighbouring molecule, while the phenomena of refraction only show resultant effects.

Further, as experiment shows that in most crystalline substances the principal directions of absorption coincide with the principal directions of optical elasticity, and as it is probably right to assume that each molecule is subject to the same laws as the whole of the crystal, there is no reason to suppose that the directions of symmetry should be different in the molecule and in the crystal, provided the latter presents no optical anomaly. One can therefore assume that the principal directions of absorption in the molecules themselves coincide with their axes of optical elasticity, and that in mixed crystals the anomalous directions of absorption indicate the direction of the optic axes of the different absorbing substances. If this is really the cause of the anomalies in the direction of certain bands, each group of anomalous bands ought to belong to different substances, of which the existence in the crystal is thus revealed.

To prove the truth of this beautiful theory, M. Becquerel points out the significant fact that among the substances which he finds to be characterized by anomalous bands, several have been separated chemically into their components.

We have, therefore, in the observation of anomalous directions of absorption a new method of spectral analysis, a method of determining in a crystal the presence of isomorphous substances, optically dissimilar. We can even go further still, and recognize the existence of different substances, although they may not manifest anomalous directions of absorption. For, suppose the same bands are noticed to occur in the spectra of several crystals; if in one of these crystals two bands become maxima or minima at the same time for the same direction of vibration, and if in another crystal one of them disappears for the direction which renders the other a maximum, one may conclude that the bands are due to two different molecules.

This new method of analysis appears to be specially suitable for use in unravelling the mystery of the constitution of the rare earths. If, as seems now almost certain, they consist of the oxides of a large number of elementary substances, the difficulty experienced in separating them points to the fact that these constituent oxides must resemble each other closely. It is therefore most probable that their salts will be isomorphous, and the crystals of these salts may consequently be expected to give absorption spectra of great interest in the light of the foregoing theory. M. Becquerel has therefore subjected the crystalline salts of didymium to the test of experiment, with the important result that several substances have been detected which chemists have recently isolated chemically; and also new substances have been identified as constituents, of which chemical methods have not as yet revealed the presence.

It will be remembered that Auer von Welsbach, by fractional crystallization of the double nitrates of didymium and ammonium, obtained two solutions—one possessing a green colour, showing almost exclusively the three bands $\lambda = 482$, 469, and 445, and which he termed praseodymium; the other a red solution, giving the other bands of the didymium mixture except $\lambda = 475.5$, which received the name neodymium. The study of the absorption spectrum of crystals of sulphate of didymium now shows that the two groups $\lambda = 483.6$ –482.2 and $\lambda = 471.5$ –470, which have anomalous directions to a remarkable extent, are characteristic of praseodymium, while most of the bands of neodymium have directions quite different. Again, on examining these same groups belonging to praseodymium in the crystals of double nitrate of didymium and potassium,

it is noticed that the bands which appear to have the same principal directions in the sulphate have in the double nitrate directions quite different, characterizing two distinct substances. Later experiments by Demarçay have indeed shown the possibility of chemically isolating two constituents—one exhibiting the band $\lambda = 469$, the other giving the bands of praseodymium.

Hence the new method proves a most valuable test of the accuracy of chemical work. In multiplying the observations, M. Becquerel concludes that didymium is, as expected, a mixture of a large number of substances, chemically different; among the identified constituents are almost all that have been already chemically isolated, and very probably others, notably one substance which is characterized by the band $\lambda = 571.7$.

A remarkable confirmation of this new law of crystal absorption was obtained in the following way. When a crystal of the sulphate or nitrate of didymium is dissolved in water, the spectrum of absorption of the solution presents curious differences from that of the crystal. Certain bands have remained permanent, but others are displaced, and some have entirely disappeared. This is readily explained if one admits that the crystal consisted of a mixture of compounds unequally acted upon by water. The most interesting fact, however, is that the bands which manifest these variations are precisely those which in the crystal present the anomalies.

In conclusion, we see that by the employment of this new method of analysis we are enabled, without destroying the crystal, as is necessary in chemical analysis, to recognize the presence of different chemical molecules; and as we obtain three distinct spectra from the three directions of optical elasticity, the method is one of extreme sensibility. Every investigator likes to see his work confirmed, and in this most difficult field of the rare earths we cannot have too many confirmations. The more points of the compass from which we approach it the better, for we are sure then of surrounding and finally of grasping the truth itself, in all its grand simplicity.

A. E. TUTTON.

THE NEW VEGETATION OF KRAKATÃO.

THE great volcanic eruption of Krakatão in August 1883 will be fresh in most memories. It was at one time reported that the island had totally disappeared, but this was not so. Previous to the eruption, however, it was covered with a luxuriant vegetation, no trace of which existed after the event.

Dr. M. Treub, the Director of the Botanic Garden at Buitenzorg, Java, when at Kew last year informed the writer that he had visited the island the previous year, and intended publishing the results of his botanical investigations. This he has now done,¹ and as the derivation of insular floras is a subject of great interest to many persons, the substance of Dr. Treub's communication deserves a place in NATURE.

The existing portion of Krakatão Island is about three miles across, and has the form of a mountain rising out of the sea. On one side it is nearly perpendicular almost to the summit of the peak, which has an altitude of about 2500 feet, and on the other it presents a steep slope. It is situated ten miles distant from the Island of Sibesie, the nearest point where there is terrestrial vegetation; twenty miles from Sumatra, and twenty-one miles from Java. Verlaten and Lang Islands, lying much nearer Krakatão, were utterly desolated and denuded of their vegetation by the great catastrophe, and were still absolutely bare in 1886.

With regard to the total destruction of vegetable life in the island, Dr. Treub asserts that there can be no doubt:

¹ *Annales du Jardin Botanique de Buitenzorg*, vii. pp. 213–23, with a sketch map.