Temperature.	Pressure.	Vapour-Density.	Method.	Observer.	Remarks.
° C.					
440	1 atm.	$\left\{ \begin{array}{l} 11.42 \\ 11.37 \end{array} \right\} 11.39$	Dumas	Deville and Troost	
440		$\left\{ \begin{array}{l} 10.75 \\ 10.97 \end{array} \right\} 10.86$	"	Friedel and Crafts	In atm. of nitrogen
321.6	0.23	11.41	"	"	
325.2	0.19	12.47			
356.9	0.12	12.04			
357.0	0.50	11.85			In atm. of chlorine
442.2	0.27	11.66			
442.2	0.14	11.30			
440		11.14	V. Meyer	V. Meyer	
619		11.01	"	"	
Very high		$\left\{ \begin{array}{l} \text{Molecular weight} \\ = 136.1; \text{FeCl}_3 = 162.5 \end{array} \right\}$	"	A. Scott	
448		10.49	"	Grünewald and Mayer	Dissociation above 518°, with liberation of chlorine.
518		9.57			
606		8.38			
750		5.40			
1050		5.21			
1300		5.13			
448 } 518 }		Same as above			In atm. of chlorine

Yellow heat	$\left\{ \begin{array}{l} 6.67 \\ 6.38 \end{array} \right\} 6.52$	V. Meyer	V. Meyer	
1300-1400	4.34	"	Nilson and Pettersson	
1400-1500	4.29

It has been pointed out by Messrs. Friedel and Crafts that the method of Dumas has this advantage over that of Victor Meyer, that the pressure of the vapour can be accurately ascertained, whereas in Meyer's apparatus diffusion takes place to an unknown extent, so that the actual pressure of the vapour is indeterminable.

It appears to me that the following conclusions, which are in general accordance with the views of Friedel and Crafts, may be drawn from the results so far obtained:—

Aluminium Compounds.—The evidence in favour of the existence of the molecules Al_2R_6 is overwhelming, but it is probable that at high temperatures they undergo dissociation, thus $\text{Al}_2\text{R}_6 = 2\text{AlR}_3$.

Gallium Trichloride.—The results point to the existence of Ga_2Cl_6 at low temperatures, and GaCl_3 at high, the more complex molecules undergoing dissociation.

Gallium Dichloride.—Most probably GaCl_2 .

Indium Chlorides.—Formulae probably InCl_3 , InCl_2 , InCl ; very little indication of the existence of the more complex molecules, but the results are all by V. Meyer's method.

Chromium Chlorides.—The trichloride CrCl_3 undoubtedly exists; little or no evidence in favour of Cr_2Cl_6 . The dichloride, even at the highest temperature reached, appears to consist largely of molecules of the formula Cr_2Cl_4 .

Iron Chlorides.—All the results by Dumas's method, and those of V. Meyer by his own, point to the stability of the molecule Fe_2Cl_6 up to 500° or 600°. The values obtained by Grünewald and Mayer probably point to dissociation into the simpler molecules FeCl_3 , but since they observed the liberation of free chlorine above 518°, and as the results above 750° are lower even than that calculated for the simple molecule FeCl_3 , there is clearly necessity for caution in drawing deductions from the experiments.

The values obtained for ferrous chloride indicate, as far as they go, a gradual dissociation of Fe_2Cl_4 into FeCl_2 .

SYDNEY YOUNG.

University College, Bristol, November 6.

THE UTILITY OF SPECIFIC CHARACTERS.

THE question of the utility or inutility of specific characters is one which is of considerable importance in the philosophy of biology on account of its connection with the action of natural selection; and it is one which is of special interest at the present time, because of the attention which has been drawn to it by Dr. Romanes's essay on physiological selection, by the Presidential address to the Biological Section at the recent meeting of the British Association, and by various letters and articles in NATURE and elsewhere. This is a matter upon which a biologist who is practically acquainted with species can alone express an authoritative opinion. It is only the naturalist who has an intimate knowledge of the characteristics and the habits of species who can judge accurately of the relations between such structural features and the animal's habits and surroundings, and who can appreciate the fact that many structures or variations of structure may be of importance, although their precise functions and relations to environment may not yet be known.

The more minutely a group of organisms is studied, the more the object or utilitarian significance of the specific characters becomes evident. In the Tunicata, the class of animals I happen to have paid most attention to of late years, I am convinced of the practical importance or usefulness of the recognized specific modifications:—such as the condition of the muscular system, the arrangement of the vessels in the branchial sac, the number and arrangement of the tentacles, and so on—these structures being all related to most important functions, such as respiration and the regulation of the food-supply. Even in the case of such apparently trivial characters as the shapes and

distribution of the minute spicules throughout the colonies of *Leptoclinum* and some other Compound Ascidiæ, I know from experience that they affect the hardness and roughness, as well as the colour, of the colony, and so may be of considerable importance in repelling enemies and in keeping the colony free from injurious parasites. As a matter of observation, I find that the colonies of Didemnidæ (which are provided with calcareous spicules) are much freer from both external and internal parasites than are the softer-tested Compound Ascidiæ.

During the last few years I have had occasion to study closely the fauna of the sea-shore at different parts of our coast. I have spent many hours on the rocks at Puffin Island and elsewhere at extreme low water, watching the animals in the pools and under the ledges in their natural conditions. Such work impresses very forcibly upon the observer the reality and importance of such fundamentals of evolution as variation, over-crowding, and struggle for existence, the action of natural selection, the benefit of protective colouring, the completeness and the advantage of mimicry, the benefit of spicules, shells, various shapes, &c., the purposes and origins of peculiar habits, the complicated relations between the animals and their environment, and finally the utility of specific characteristics.

Of all the regions of sea and land, so far as my limited experience goes, by far the most prolific of animal life is that region of the shore which is known as the upper edge of the Laminarian zone. It lies just beyond the ordinary beach, and is only exposed at the lowest spring tides. There, amongst the tangled masses of *Laminaria*, especially if there are large irregular stones with many pools and crevices between, marine invertebrate life is to be seen in very great profusion; in a favourable locality, such as Puffin Island, all the chief groups of marine animals being abundantly represented. There, competition is probably very keen amongst allied forms of animals, and the conditions necessary for natural selection to take place, and the results of that process, may most advantageously be studied.

I have lately been noting which animals in this region of the shore appeared to be the most conspicuous by their colour or shape or other peculiarities on various occasions and in different localities, with the object of seeing how far the want of protective colouring or attempt at concealment can be accounted for; and the result is that, so far as I have observed, all the most brightly-coloured or otherwise very noticeable species are provided with some defensive or offensive contrivance which appears to protect them from enemies. Amongst these conspicuous forms are: the white calcareous and some of the scarlet and other brightly-coloured siliceous Sponges (which are well protected by their numerous pointed spicules), the gleaming white Compound Ascidiæ (provided with sharp-pointed calcareous spicules), some of the Annelids (protected by their setæ, elytra, &c.), some of the Polyzoa, such as the Escharidæ and other incrusting forms (covered by calcareous ectocysts, often provided with spines and other projections), and a few erect forms such as *Bugula* (which are protected by the presence of numerous sharp-beaked snapping avicularia). The bright orange-coloured *Bugula turbinata* is certainly one of the most conspicuous animals on certain parts of the shore about low-water mark at Puffin Island. Of course such protective characters as these animals possess are not all necessarily specific, but may be generic, or characteristic of still larger groups, and I am not now citing them as instances of useful specific characters, although I do not doubt they would prove such, if the details were properly worked out.

As examples of these last (useful specific characters), I would point to the distinctive features of the species of Ascidiæ, where even the very varied external shapes may

be regarded as useful modifications, since they allow of, or correspond to, particular forms of the muscular mantle and the branchial sac and the other viscera within the test, and of course the shapes of the mantle and branchial sac are of functional importance. It is important to note that the one external feature which it is difficult to see any use in—viz. the number of lobes surrounding the branchial and atrial apertures—is not a specific character, but is distinctive of genera, and even more so of families and sub-families.

On passing to the interior of the body in the Ascidiæ, we find that the best-marked characters are taken from the condition of the mantle and of the branchial sac, and its neighbouring structures the tentacles and the dorsal lamina. Now, these are all organs with most important functions to perform in regard to respiration, nutrition, the circulation of water through the body, and the collection and agglutination of food-particles. And all the structural modifications found are such as must evidently be of actual use to the possessors. The current of water passing through the body of an Ascidian—in at the branchial aperture, through the stigmata in the walls of the branchial sac into the atrial cavity, and from that out by the atrial aperture—is of primary importance, since it serves the following purposes: (1) it conveys oxygen into the body for respiratory purposes, (2) it brings the food-matters into the body, (3) it removes waste matters from the body, and (4) it conveys to the exterior the ova and spermatozoa. This current of water is caused and guided by (a) the shape of the mantle and the arrangement of the sphincters and other muscles, and (b) the cilia covering certain of the vessels and other parts of the wall of the branchial sac. Hence modifications of the form of the mantle and of its muscles, and of the vessels, bars, papillæ, &c., forming the wall of the branchial sac (which are precisely the characters made use of in distinguishing the species), must surely be of functional importance, or, in other words, are useful modifications, such as would be produced by the action of natural selection.

It is scarcely necessary to call attention to such important adaptive characters as the arrangement of the blood-vessels and water-passages in the walls of the branchial sac, but it may be pointed out that even such trivial structures as the spine-like scales lining the branchial siphon in some Cynthiidæ may well be more or less useful, according to their shape and size, in keeping out small unwelcome intruders, such as the young of the parasitic Copepoda, sometimes found in the branchial sacs of some Ascidiæ.

Another point in which species of Ascidiæ differ is the condition of the tentacles round the entrance to the branchial sac—i.e. their number, shape, branches, and arrangement. These organs probably perform various functions: they break up and distribute the currents of water, they intercept and guide the food-particles, they probably act as sensory organs, and they form a more or less perfect grid for preventing large objects from entering the branchial sac. Hence there can be little or no doubt that in this case also the various modifications are really useful.¹

These few instances are, perhaps, sufficient to show that, in the Tunicata at least, specific characters are of actual importance to their possessors, and are adaptive modifications such as would be produced by the action of natural selection; and I fancy that the same will be found to be the case in other groups of animals, if those biologists who are intimately acquainted not only with the characters of the various species, but also with their habits in a condition of nature, and the environment generally, would turn their attention to the matter.

W. A. HERDMAN.

¹ I am working up the matter in further detail for the Report upon the Tunicata in the second volume of the "Fauna of Liverpool Bay."