

comprising several interesting points of construction and detail, is at work at the Paris Exhibition.

Mr. Deboutteville, in his paper, first reviews the gas-engines hitherto proposed or employed. He carries back his researches nearly a century, when the first gas-engine was proposed by Barber, and completes them with a description of the Simplex engine, which he brought out with Mr. Malandin in 1884. This engine is founded on principles laid down by Mr. Beau de Rochas—that, to realize the best results from the elastic force of gas, the cylinders should have the greatest capacity with the smallest circumferential surface, the speed should be as high as possible, the cut-off should be as early and the initial pressure as high as possible. In the author's engine the ignition is effected by a practically continuous electrical spark; the air and gas are mixed in an external receptacle fixed on the cover of the slide, and are drawn in through channels of varied forms so as intimately to mix them. The governors described act on the principle of totally cutting off the supply of gas for one or more strokes whenever an increase of speed occurs. From the tests made with this engine the consumption of gas is low, and it appears to compare favourably with good steam-engines as regards economy of application.

Besides the reading and discussion of papers, the members of the Institution visited the Exhibition, and various works which were opened for this purpose.

SCIENTIFIC SERIALS.

THE numbers of the *Journal of Botany* for June and July are chiefly devoted to articles interesting to students of systematic or geographical botany, especially that of our own islands; the latter number contains a biographical sketch of the late Prof. Reichenbach, by the editor.

THE most interesting article in the *Botanical Gazette* for May is the commencement of a detailed paper by Mr. C. Robertson, on the relations between insects and flowers in regard to American plants. The number for June contains original articles by Mr. H. L. Bolley, on sub-epidermal rusts, and by Mr. J. N. Rose, on the Achenia of *Coreopsis*.

THE *American Meteorological Journal* for May contains abstracts of the papers read at the meeting of the New England Meteorological Society on April 16:—In a paper on lightning and the electricity of the atmosphere, Mr. McAdie gave an account of some kite experiments at the Blue Hill Observatory, near Boston, in which the potential was determined at various heights. He also referred to the observations on the character of lightning at the top of Mount Washington during thunderstorms, and to the effect of the electrification of the air upon water, dust, and other particles in it, and to the possibility of foretelling the moment of a flash of lightning.—Prof. W. M. Davis made a report upon the investigation of the sea-breeze, undertaken in 1887, from observations at 100 stations. One fact shown was that the diurnal range of temperature, which is diminished on the coast by the action of the sea-breeze, is not lessened at the inland stations.—Mr. E. B. Weston read a paper on the practical value of self-recording rain-gauges, referring to the importance of knowing the hourly falls when constructing drainage systems.—Prof. H. A. Hazen continues the discussion upon anemometer comparisons, and upon the question of the probable effect of the momentum of heavy cups, when placed on a whirling machine. He considers that the Robinson anemometer is by far the best instrument ever devised for variable winds.—Lieut. Finley discusses the frequency of tornadoes in Illinois for fifty-four years, ending with 1888. The total number of storms was 141. The month of greatest frequency was May, no month being free from storms. The prevailing direction of movement was north-east.—Prof. Harrington communicates the instructions issued by the Chief Signal Officer for the preparation of forecasts and for their verification. The instructions contain nearly 200 regulations, and are very interesting to those who study weather predictions.

THE *Meteorologische Zeitschrift* (Vienna) for June contains the first part of an epitome of Dr. von Bezold's papers on the thermodynamics of the atmosphere, which have already been summarized in our notices of Societies.—Dr. J. Hann contributes a valuable article on the results of the meteorological observations of the late Prof. A. Ackermann at Port-au-Prince, Hayti, 1854-68, being a part of the world where they are of special value. The

observations were rescued from entire loss by the exertions of Dr. Hann and Prof. J. Scherer, the originals having been wilfully destroyed. The distribution of rainfall is much affected by the mountain features of the island; in the north the rainy season is from December to April, while in the south it is from May to July. The average yearly rainfall at Port-au-Prince, from the above observations, was 61 inches, on 153 days. The greatest daily fall was 5.6 inches in May 1865, the rain lasting four hours. The climate is very equable; the mean of the absolute maximum temperature was 98° 2, and of the minimum 56° 8.—Dr. von Lepel describes his experiments in passing electric sparks through glass tubes lined with a thin coating of paraffin, and containing a small amount of moisture, and points out that during thunderstorms many similar discharges may be observed, and may find their explanation in these experiments. The sparks differ in character and in colour, and the author argues that the humidity in the tube may be compared to the particles of vapour in the thunder-clouds, and that the coating of paraffin may have the same optical effect as the translucent clouds themselves. He gives the results of his thunderstorm observations on these lines during the summer of the year 1888.

SOCIETIES AND ACADEMIES.

LONDON.

Physical Society, June 22.—Prof. Reinold, F.R.S., President, in the chair.—The following communications were made:—Note on some photographs of lightning, and of "black" electric sparks, by Mr. A. W. Clayden. The lightning photographs, three in number, were obtained during the storm on June 6. Two flashes, seen on one plate, show complicated and beautiful structure: one of them is a multiple flash, and flame-like appendages point upwards from every angle; the other is a broad ribbon, and, although the plate shows signs of movement, the displacement is not in a direction such as would produce a ribbon-like effect from a linear flash. The second plate shows four flashes, none of which are ribbon-like, though the camera had moved considerably. The third plate was exposed to six flashes, one of which was believed to pass down the middle of the plate; but, on development, only a triple flash in one corner of the plate was seen. Careful search, however, revealed the central flash as a dark one with a white core, and other dark flashes were subsequently found. The plate was very much over-exposed, and this suggested that black flashes might be due to a sort of cumulative action caused by the superposition of the glare from a white cloud upon the normal image of the flash. To test this, sparks from a Wimshurst machine were photographed, and, before development, the plates were exposed to diffused gas-light for a short time. The bright sparks yielded normal images with reversed margins, and the faint ones were completely reversed. Other experiments showed the reversal to spread inwards as the time of exposure to gas-light increased. Finally, reversal was effected by placing a white screen behind the spark, to represent a white cloud, the only illumination being that of the spark itself. In the discussion which followed, Mr. W. N. Shaw exhibited a photograph taken during the same storm, which is particularly rich in dark flashes branching outwards from an intensely bright one. In some places the bright line has dark edges, and in one part a thin bright line runs along the middle of an otherwise dark portion of the flash. In answer to Mr. Inwards, Mr. Shaw said the plate was exposed about half a minute, and the former thought that, under those conditions, the appearance of the plate did not contradict Mr. Clayden's hypothesis. Speaking of the same photograph, Prof. Perry considered that Mr. Clayden's observations would explain the result, for a bright flash required more exposure to diffused light to reverse it than a faint one did. Prof. Ramsay reminded the meeting that Prof. Stokes's "oxides of nitrogen" explanation was still a possible one; and Mr. C. V. Burton asked whether they may be due to faint sparks cutting off light from brightly illuminated clouds, just as a gas-flame absorbs light from a brighter source. In reply, Mr. Clayden thought the "oxides of nitrogen" hypothesis improbable, and said his experiments did not enable him to answer Mr. Burton's question. As regards Mr. Shaw's plate, he believed the diffused light from the clouds would be sufficient to reverse the fainter tributary flashes, although it was insufficient to reverse the primary one. From data obtained when the ribbon-flash was taken, he had made some calculations which gave the height of the clouds about 1000 yards, and the ribbon-flash 1300 yards