occurs in Nevis, whence the Zoological Society received living specimens (presented by Mr. Graham Briggs) in 1870.

The only West Indian island in which *Quadrupana* of the *Americana* type occurs is Trinidad, which was, doubtless, formerly part of the mainland of South America.

3 Hanover Square, W., February 17. P. L. SCLATER.

Galls.

I HAD not intended to take any further part in this correspondence; but the interesting suggestion which has now been made upon the subject by Mr. T. D. A. Cockerell (*NATURE*, Feb. 13, p. 344) induces me to withdraw the sentences that he quotes from my previous letters, to the effect that it seems impossible to imagine any way in which galls can be attributed to natural selection acting on the plants *directly*. In my own consideration of the matter this seemed "obvious," and therefore my motive in taking up the difficulty as presented by Mr. Mivart was that of "asking whether anybody else had a better explanation to offer" than the one which my letter suggested—viz. "that natural selection may operate on the plants *indirectly through the insects*," by always selecting those insects the character of whose secretions is such as will best cause the plants to grow the particular kind of morphological abnormality which the larvæ require. Mr. Cockerell, however, has now furnished what seems to me an extremely plausible hypothesis, showing that there is a way in which it is quite conceivable that the growth of galls may be an actual benefit to the plants, and therefore that natural selection may act directly on the plants themselves in evolving these sometimes highly specialized structures for the use of their parasites. Mr. Cockerell informs me in a private communication that he has been verifying this hypothesis by observations in detail; but whether or not he will be able to establish it, I think at any rate he has done good service in thus suggesting another possibility.

On the other hand, I cannot see that Mr. Ainslie Hollis has helped us at all (*NATURE*, January 23, p. 272). For he merely enunciates the truism that trees which were not endowed with sufficient "developmental vigour" adequately to resist the attacks of gall-making insects "would doubtless have long ago succumbed in a struggle for existence." And this truism he appears to suppose furnishes an explanation of how "natural selection, operating in the ordinary manner," has produced galls for the exclusive benefit of the insects. But it is obvious that the more detrimental the growth of galls has proved to trees, the less reason there must have been for natural selection, "operating in the ordinary manner," to have developed these often highly specialized structures for the benefit of parasites.

London, February 13. GEORGE J. ROMANES.

The Supposed Earthquakes at Chelmsford on January 7.

NATURE for January 16 (p. 256) reprints from the *Essex County Chronicle* a short account of two supposed earthquake-shocks felt at and near Chelmsford on January 7, at 12.30 and 1.25 p.m. Being engaged in the study of British earthquakes, I made inquiries in the district referred to, and the result of these is to show that the shocks were almost certainly due to the firing of unusually heavy guns at Woolwich. It may be worth while to state the evidence for this conclusion somewhat fully, as it will be difficult to obtain it in after years.

(1) I applied to the authorities at Woolwich and Shoeburyness as to the nature of the firing on January 7. At the latter place, the only practice was from 9-inch and 10-inch guns, the maximum charge used was 70 pounds of powder, and therefore not capable of producing the shocks felt at Chelmsford. At Woolwich, however, the 110-ton gun, "the heaviest in *H.M. service*," was fired at the times mentioned.

(2) *Form of the Disturbed Area*.—The only accounts I have as yet received are from the following places: Great Warley (near Romford), Brentwood, Epping, Ingatestone, on the road between Ongar and Fyfield, Roxwell, Chelmsford, Chignall St. James, and Chipping Hill (Witham); which are respectively at about 6, 12½, 14, 16, 16, 21, 24, 24, and 32 miles distance from Woolwich. Referring to a map of Essex, it will be seen that these places all lie close to a line drawn from Woolwich in a north-easterly direction; with the exception of Epping, the direction of which is about north by east from Woolwich. According to the *Times* weather report of January 8, southerly and

south-westerly breezes prevailed very generally throughout the kingdom on the previous day.

(3) *Nature of the Shock*.—In four cases, the shock was in the first instance attributed to the firing of heavy guns. If there was any vibration of the earth, it must have been very slight, and the following descriptions seem to leave little doubt that the rattling of windows noticed was due to an air-wave.

Great Warley—The shock "broke a pane of glass 4 feet x 2 feet on my job."

Brentwood—"The shocks commenced as a low rumble, increasing till the doors shook and rattled, as though the rumbling was followed by a bang or explosion."

Between Ongar and Fyfield (the observer driving)—"The ground felt as if it were sinking," and there was "a rumbling noise something like guns in the distance."

Roxwell—The sound "exactly resembled the report of the big guns at Shoebury, but was far louder than we usually hear them."

Chelmsford (the observer walking)—There was "a noise as of a very heavy weight being rolled across the floor of the room of the house to the south of him, which he was passing."

Chignall St. James—"The shock was extremely slight, but there was a most pronounced concussion in the air which made a sound on the windows as if a person had thumped the centre of the window frame with the soft part of his hand. There was no tremulous motion felt."

Witham—The observer "heard a strange rumbling sound which seemed to slightly deafen him, but he felt no vibration of the earth."

That the disturbances recorded had only one origin is, I think, evident, (1) from the decrease in intensity (roughly speaking) as the distance from Woolwich increases, and (2) from there being no considerable gap between the places of observation. Records from the immediate neighbourhood of Woolwich could hardly be expected, as there they would naturally be attributed to their proper source.

I am indebted to the editor of the *Essex County Chronicle* for inserting a letter asking for observations on the shocks, and to several gentlemen for the courtesy and kindness with which they replied to this letter and to other inquiries that I made in the surrounding district.

CHARLES DAVISON.

38 Charlotte Road, Birmingham, February 13.

Shining Night-Clouds.

IN July last, on a fine night, about 8 p.m. (two hours after sunset), I noticed a fleecy cloud lit up by a yellowish light, directly over the back of a range of hills due west from this place. As it did not move, it struck my attention, and I observed that what little wind there was carried the few floating clouds north-east to south-west. I continued to watch the cloud, which covered 4° or 5°, until 11 p.m., and concluded that as in that direction lay the Puracé volcano, about 40 miles away, the light and cloud probably came from it. But I made inquiries by telegraph, and found that no eruption had taken place in the Puracé, which has been quiet now for many years. I regret, seeing now that the subject is interesting, that I did not observe more carefully. I may add that in the direction of the cloud no prairie or forest fire could have occurred to account for it.

ROBERT B. WHITE.

Agrado (lat. 2° 20' N.), Department of Tolima,
U.S. of Colombia, S.A., December 22, 1889.

A Greenish Meteor.

TO-NIGHT (Jan. 30), at 8.15 p.m., I saw a meteor which, notwithstanding a bright moon, shone out exceedingly brightly, exceeding any star. It appeared to travel south, for about 10°, vanishing about 15° above the horizon. Its colour differed from that of any meteor I have seen before, being pale green or greenish.

T. D. A. COCKERELL.

West Cliff, Custer Co., Colorado, January 30.

THE MOLECULAR STABILITY OF METALS, PARTICULARLY OF IRON AND STEEL.

(1) ALLOW me to add some words relative to the very timely lecture on the hardening and tempering of steel, recently published by Prof. Roberts-Austen

(NATURE, xli. pp. 11, 42). I desire, in the first place, to point out the bearing of the singular minimum of the viscosity of hot iron (*loc. cit.*, p. 34) on the interpretation given of Maxwell's theory of viscosity (*Phil. Mag.* (5), xxvi. pp. 183, 397, 1888; xxvii. p. 155, 1889). When iron passes through Barrett's temperature of recalcence, its molecular condition is for an instant almost chaotic. This has now been abundantly proved (cf. John Hopkinson, *Phil. Trans.*, London, clxxx. p. 443, 1889, where the literature may be found; cf. Osmond, below). The number of unstable configurations, or, more clearly, the number of configurations made unstable because they are built up of disintegrating molecules, is therefore at a maximum. It follows that the viscosity of the metal must pass through a minimum. Physically considered, the case is entirely analogous to that of a glass-hard steel rod suddenly exposed to 300°. If all the molecules passed from Osmond's β state to his α state together, the iron or steel would necessarily be liquid. This extreme possibility is, however, at variance with the well-known principles of chemical kinetics. The ratio of stable to unstable configurations cannot at any instant be zero. Hence the minimum viscosity in question, however relatively low, may yet be large in value as compared with the liquid state.

(2) My second point has reference to the function of carbon in steel. It is not to be understood that we ignore the importance of the changes of carburization produced by tempering steel. To explain the varied physical phenomena which accompany temper, it is sufficient to recognize some *special* instability in the tempered metal. This is given by the carbide configuration, and the physical explanations in question may be made without specifying its nature further. Hence the permissibility of the purely physical considerations.

On the other hand, it is indeed surprising that, on the part of engineers and chemists, the important subject of temper has been but inadequately dealt with, as Prof. Austen justly remarks. Sir Frederick Bramwell (cf. NATURE, xxxviii. p. 440), in his inaugural address at Bath, in 1888, dwelt at some length on the subject of temper. The question is again touched upon by Mr. Anderson at the Newcastle meeting of the British Association. Neither of these gentlemen, however, really shows forth the gist of the matter. Indeed, even in Ostwald's massive "*Lehrbuch der Allgemeinen Chemie*" (Leipzig, W. Engelmann, 1887), full of examples as it is, bearing on all points of chemical physics, the frequent and exceptionally important case of tempered steel is altogether absent. And yet the chemical interpretation to be given to the phenomena of temper seems to be closely at hand. Dr. Strouhal and I (*Wied. Ann.*, xi. p. 390, 1880; Bulletin U.S. Geol. Survey, No. 14, chap. ii., 1885) showed that, by the process of hardening, the electrical resistance of steel may be increased by more than three times its value for the soft metal. If the hard rod is now softened, the resistance again decreases by an amount depending on the temperature to which the hard metal is exposed and on the time of such exposure, in a way which, throughout the whole research, is beautifully sharp and characteristic. Eventually, the relatively low resistance of soft steel is again reached. Now suppose the carbon molecule of steel to be dissolved in the metal, forming an alloy of Matthiessen's Class II. Seeing that the quantity of carbon contained is not large, the electrical resistance of hard steel is at once an expression of its chemical composition, structurally unknown though it be. Hence in the electrical diagram of the phenomena of temper constructed by Dr. Strouhal and myself, the time variations of resistance of hard steel at any given temperature may be interpreted as a case of Wilhelm's (*Pogg. Ann.*, lxxxi., pp. 413, 499, 1850) rate of chemical reaction (*Reaktionsgeschwindigkeit*), and expressed in accordance with his well-known exponential law. This indeed is the character

of the observed time curves. Hence also the full diagram of the phenomena of temper, considered both in their variation with time and with temperature, is available for the elucidation of most points relative to the effect of temperature on rate of chemical reaction.¹

(3) A further remark may be made relative to Osmond's (*Annales des Mines*, July-August, 1888, pp. 6-7; *Mém. de l'Artillerie de la Marine*, Paris, 1888, p. 4) iron of the α and the β type. The assertion that mere strain partly changes α into β iron is in conformity with the viscous behaviour of the metal. For it appears that the effect of any mechanical strain as well as of temper, is marked decrease of the viscosity of the metal. Osmond's theory, however, appears to explain too much. Since most metals can be similarly hardened by straining, it would follow that there should be α and β varieties in all these cases, even though a molecular change corresponding to Gore's phenomenon in iron has only in a few instances been observed (iron, nickel, platinum-iridium alloy). I believe, however, that there is reason to be urged even in favour of this extreme view.² The ion theory of metallic conductivity is fast gaining ground.

J. J. Thomson states it in his well-known book ("Applications of Dynamics," p. 296). Giese (*Wied. Ann.*, xxxvii. p. 576, 1889) has outlined an ion theory of electric conduction, uniformly applicable to metals, electrolytes, and gases. It seems to me, if a preliminary hypothesis be made relative to the evolution of a magnetic field out of an electric field; if advantage be taken of the spiral distribution of points which frequently results from the symmetrical interpenetration of two congruent Bravais systems;³ if, finally, in metals, the function performed by a bodily transfer of ions can also be performed by an exchange of the charges of charged atoms (Giese, indirectly Helmholtz), that the possibility of an ion theory of magnetism may be suspected. Quite apart from the influence of a field, the conditions of exceptionally close approach favourable to the transfer of charges from atom to atom, are given by the distribution of the heat agitation in the metal.

(4) I will close this note by some remarks on the change of the character of diffusion when occurring in solids. Studying the coloured oxide coats on iron, Dr. Strouhal and I (*Bull. U.S.G.S.*, No. 27, p. 51, 1886) pointed out that the outer surface of the film is oxidized as highly as possible in air; and that the inner surface of the film, continually in contact with iron, is reduced as far as possible. This distribution of the degree of oxidation along the normal to the layer, is equivalent to a force in virtue of which oxide is moved through the layer, from its external surface to its internal surface. The formation of an oxide coat is thus a case of diffusion. Conformably with this view, the film, during its formation, behaves like an electrolyte, as was pointed out by Franz, Gauguin, and Jenkin, and more recently by Bidwell and by S. P. Thompson.

We then adverted to the crucial difference between diffusion in solids and diffusion in liquids, inasmuch as in the former case (solids) diffusion demonstrably *ceases* after a certain small thickness is permeated. The limit thickness of the film is reached asymptotically, through infinite time. It has a definite value for each temperature, increasing as temperature increases. In the light of other evidence since gained, this explanation is substantiated. The formation of the

¹ An ulterior consideration presents itself here relative to an extension of the theory of Arrhenius (*Wied. Ann.*, iv. p. 391, 1878) to metallic conductivity. Arrhenius and Ostwald find in the maximum of electrolytic conductivity a measure of rate of reaction. I must pass over this question here, since it is without immediate bearing on the text.

² I have spent much time in endeavouring to throw light on this question, and will indicate the results later. My methods were (1) to find the effect of mechanical strain on the carburization of steel; (2) to find the effect of strain on the rate of solution; (3) to find the hydro-electric effect of stretching.

³ A good account of the relations of the Bravais and the Sohncke system is given by H. A. Miers, in NATURE, xxxix. p. 277.

oxide coat is a case of solid diffusion, and as such it bears the same relation to the diffusion of liquids, that the viscosity of solids bears to the viscosity of liquids. The two phases (solid, liquid) of each phenomenon are to be correlated in ways essentially alike. The available stress, as compared with the available instability at a given temperature, determines the time character of the result.

CARL BARUS.

Physical Laboratory, U.S. Geological Survey,
Washington, D.C.

CHRISTOFORUS HENRICUS DIEDERICUS
BUYS BALLOT.

BUYS BALLOT was born on October 10, 1817, at Kloetinge in Zeeland; was a student in arts and the natural sciences at the University of Utrecht, where he first became Lector of Physics and Chemistry in 1844, and then successively Professor of Mathematics in 1847, and of Experimental Physics in 1870, which latter chair he ceased to hold in November 1887 on completing his fortieth year as Professor. He was appointed Director of the Royal Meteorological Institute of the Netherlands in 1854, and held this position with great ability and distinction till his death on Monday, the 3rd of the present month.

His first contribution to science was a paper on a chemical subject in 1842, this being a science of which he was Lector at the time; but soon thereafter he turned his attention to meteorology, which he emphatically made the business of his life. The following are among the earlier of his papers on the subject, and they are, it will be seen, very significant of his future work:—"On the Influence of the Rotation of the Sun on the Temperature of our Atmosphere," in 1846; "On the Importance in Meteorology of Deviations from the Mean States of the Atmosphere," in 1850; "Results of the Observations of 1849 and 1850 in different places in Holland," in 1851; and "On Synchronous Representations of Weather Phenomena," in 1854.

In these early times of meteorology, when instruments and modes of observing still greatly needed the guiding hand of science towards the founding of international meteorology, Dr. Buys Ballot was wisely led to attempt the construction of no general isobaric and isothermal maps in investigating storms and other weather phenomena, but contented himself in investigating weather disturbances by representing them over the surface of Europe by means of deviations from the means, or averages, of the places represented. In this mode of working he made several of his more important contributions to meteorology, and out of it developed the system of storm warnings he issued for Holland. In this connection his barometric and thermometric means for a very large number of places over Europe will long be a standard work. Of these contributions, unquestionably the most important is that known as BUYS BALLOT'S LAW OF THE WINDS, which states the relation between the direction of the wind and the distribution of atmospheric pressure at the time the wind is blowing. This relation was further developed by Dr. Buchan in 1869, in his paper on the mean pressure of the atmosphere and prevailing winds of the globe, in which it was shown that the prevailing winds of all climates are simply the result of the distribution of pressure.

One of the most exhaustive discussions of the influence of the moon on weather was made by Dr. Ballot. The discussion covered a period of about a century, and he showed that the longer the period the closer do the cases for or against any such influence approach equality. Subsequent to Maury, Dr. Ballot was one of the earlier and most energetic and successful workers in maritime meteorology, and his meteorological charts of the routes of

Dutch ships over the great oceans is a standard work. Dr. Ballot also took an active and efficient part in the Meteorological Conferences and Congresses held at intervals from 1872 to 1888, which have brought about a greater uniformity in meteorological observations and discussions than previously existed. He was chosen, by the first Congress, President of the Permanent Committee. Among his last works was the proposal of a method of developing and representing the variability of the weather and climates by the values of the deviations of the daily observations from the averages, irrespective of sign.

The great merits of his indefatigable services to science, but more particularly to meteorology, were recognized by his being made LL.D. of Edinburgh University, Knight of the Order of the Netherland Lion, Commander of the Order of Franz Joseph of Austria, and of St. James of the Sword of Portugal, and Knight of second class of the Prussian Order of the Crown. But above all, his ever readiness in every degree to oblige, the genial sunshine of his face, and his loveliness, make his death to be felt by many of us as a sharp personal bereavement.

NOTES.

ON Tuesday evening the Cambridge University Natural Science Club and the Master of Downing (Dr. Alex. Hill) gave a *conversazione* at Downing Lodge, at which 260 guests, including many distinguished residents and non-residents, were present. The several scientific professors were very liberal in lending the treasures from their museums, and as this is the first entertainment of the kind which has been given in Cambridge, many objects of great historic interest, such as Babbage's calculating machine, Cavendish's apparatus, &c., were exhibited. Artificial silk was spun, quartz filaments drawn, smokeless gunpowder and other scientific novelties shown. One of the most interesting exhibits was a series of Egyptian heads unwrapped from their mummy cloths, and artfully "restored" by Prof. Macalister. A very attractive feature of the entertainment was an address by Dr. Lauder Brunton, who had much that was interesting to say about his recent experiences in India. Mr. Gardiner illustrated the dispersion of seeds by the aid of the limelight and boxes of seeds of various kinds suspended from the ceiling.

THE annual general meeting of the Geological Society of London will be held to-morrow (Friday) at 3 o'clock, and the Fellows and their friends will dine together at the Criterion Restaurant at 7.30 p.m.

BEFORE the next ordinary meeting of the Royal Microscopical Society, it will have moved its quarters from the rooms hitherto occupied by it in King's College, which are now required for the purposes of the College, to 20 Hanover Square. The ordinary meetings will in future be held on the third instead of the second Wednesday in the month, and the annual meeting in January instead of February. The Quekett Microscopical Club has also transferred its place of meeting to 20 Hanover Square since the commencement of the year.

WE regret to have to record the death of Sir Robert Kane, F.R.S. He died after a short illness on Sunday, the 16th inst., at his residence in Dublin.

THE fine buildings of the University of Toronto were almost wholly destroyed by fire last Friday. The flames were unfortunately fanned by a strong wind, and the fire spread so rapidly that hardly anything could be saved. A small number of specimens in the museum, and some of the scientific apparatus, were brought out by students, but they were mostly broken while