with an aluminium frame, has been made for the Observatory at Kensington, and gives every satisfaction. The comparison spectrum used at Potsdam has been furnished by a Geissler tube placed directly in the cone of rays of the refractor, at a distance of 40 cm. from the slit, the tube being at right angles to the optical axis of the refractor and the slit. The slit is set parallel to the line of the diurnal motion, and width is given to the spectrum by making the driving-clock move slightly slower or faster than its proper rate. A uniform exposure of one hour has been employed, the proper intensity being obtained by changing the rate of the driving-clock, so that the error increases with increase of brightness. The photographs are measured with the aid of a microscope having a sliding apparatus on its table, movable by a fine micrometer screw. One revolution of the screw corresponds to a difference of wave-length of $0.324 \mu\mu$, which, expressed in miles per second, is 139.13 . After describing the methods of measuring the displacement of lines in stars of different types of spectra, Prof. Vogel brings together the results which have formed the subject of several previous communications. It is said that the probable error in the determination of the radial velocity of a star of Class II. is ± 1.34 miles per second, and for stars of Class I., ± 2.31 miles. Measurements have been made independently by Prof. Vogel and Dr. Scheiner, and each star has been observed on the average 3.3 times, wherefore it is concluded "that the probable error of the definitive values for both spectral classes will amount to less than one mile." A list of the observed velocities of forty-seven stars will soon be published. The mean motion in the line of sight is 10.6 English miles per second; six stars have a velocity less than 2 miles per second, and five greater than 20 miles. α Tauri heads the list with a velocity of about + 36 miles per second. Fifteen of the stars have a positive, and thirty-two a negative motion.

ORTHOCHROMATIC PLATES FOR ASTRONOMICAL PHOTOGRAPHY.—MM. Fabre and Andoyer photographed the eclipsed moon at Toulouse Observatory on November 13, 1891; and some of the pictures obtained were exhibited by them at the meeting of the Paris Academy of January 11, with a note on the method of production. Collodion-bromide and collodion-chloride plates were employed, both kinds being treated with eosin and cyanin to render them orthochromatic. The former kind of plate was found to be relatively more sensitive to red and yellow rays than the latter, although both were stained with the same dyes. It is proposed, therefore, to use collodion-bromide orthochromatic plates to obtain photographs of Mars, Jupiter and the red spot, and coloured stars.

DREDGING OPERATIONS IN THE EASTERN PACIFIC.

THE *Bulletin* of the Museum of Comparative Zoology at Harvard College, published in June, contains three letters from Prof. Alexander Agassiz to the Hon. Marshall McDonald, United States Commissioner of Fish and Fisheries, on the dredging operations off the west coast of Central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California. The operations, which were in charge of Prof. Agassiz, were carried on by the U.S. Fish Commission steamer *Albatross*, Lieutenant Commander Z. L. Tanner, U.S.N., commanding.

I.

Steamer "*Albatross*," Panama, U.S. of Colombia,
March 14, 1891.

MY DEAR COLONEL McDONALD,—We returned yesterday from our first trip. The route extended from Panama to Point Mala, and next to Cocos Island; from there we ran in a southerly direction, then north-westerly to Malpelo Island, and back to the hundred-fathom line off the Bay of Panama. We spent several days trawling off the continental plateau of the Bay. This trip being rather in the nature of a feeler, I cannot tell you just what I think it means. But I believe I can to some extent conjecture probabilities from what has been accomplished.

I have found, in the first place, a great many of my old West Indian friends. In nearly all the groups of marine forms among the Fishes, Crustacea, Worms, Mollusks, Echinoderms, and Polyps, we have found familiar West Indian types or east coast forms, and have also found quite a number of forms whose wide

geographical distribution was already known, and is now extended to the Eastern Pacific. This was naturally to be expected from the fact that the district we are exploring is practically a new field, nothing having been done except what the *Albatross* herself has accomplished along the west coast of North and South America. The *Challenger*, as you will remember, came from Japan to the Sandwich Islands, and from there south across to Juan Fernandez, leaving, as it were, a huge field, of which we are attacking the middle wedge. As far as we have gone, it seems very evident that, even in deep water, there is on this west coast of Central America a considerable fauna which finds its parallel in the West Indies, and recalls the pre-Cretaceous times when the Caribbean Sea was practically a bay of the Pacific. There are, indeed, a number of genera in the deep water, and to some extent also in the shallower depths, which show far greater affinity with the Pacific than with the Atlantic fauna. Of course, further exploration may show that some of these genera are simply genera of a wider geographical distribution; but I think a sufficiently large portion of the deep-sea fauna will still attest the former connection of the Pacific and the Atlantic.

I am thus far somewhat disappointed in the richness of the deep sea fauna in the Panamic district. It certainly does not compare with that of the West Indian or Eastern United States side. I have little doubt that this comparative poverty is due to the absence of a great oceanic current like the Gulf Stream, bringing with it on its surface a large amount of food which serves to supply the deep-sea fauna along its course. In the regions we have explored up to this time, currents from the north and from the south meet, and then are diverted to a westerly direction, forming a sort of current doldrums, turning west or east or south or north according to the direction of the prevailing wind. The amount of food which these currents carry is small compared with that drifting along the course of the Gulf Stream. I was also greatly surprised at the poverty of the surface fauna. Except on one occasion, when, during a calm, we passed through a large field of floating surface material, we usually encountered very little. It is composed mainly of *Salpæ*, *Doliolum*, *Sagittas*, and a few *Siphonophores*—a striking contrast to the wealth of the surface fauna to be met with in a calm day in the Gulf of Mexico, near the Tortugas, or in the main current of the Gulf Stream as it sweeps by the Florida Reef or the Cuban coast near Havana. We also found great difficulty in trawling, owing to the considerable irregularities of the bottom. When trawling from north to south, we seemed to cut across submarine ridges, and it was only while trawling from east to west that we generally maintained a fairly uniform depth. During the first cruise we made nearly fifty hauls of the trawl, and, in addition, several stations were occupied in trawling at intermediate depths. In my dredgings in the Gulf of Mexico, off the West Indies, and in the Caribbean, my attention had already been called to the immense amount of vegetable matter dredged up from a depth of over 1500 fathoms, on the lee side of the West Indian Islands. But in none of the dredgings we made on the Atlantic side of the Isthmus did we come upon such masses of decomposed vegetable matter as we found on this expedition. There was hardly a haul taken which did not supply a large quantity of water-logged wood, and more or less fresh twigs, leaves, seeds, and fruits, in all possible stages of decomposition. This was especially noteworthy in the line from the mainland to Cocos Island, and certainly offers a very practical object-lesson regarding the manner in which that island must have received its vegetable products. It is only about 275 miles from the mainland, and its flora, so similar to that of the adjacent coast, tells its own story. Malpelo, on the contrary, which is an inaccessible rock with vertical sides, and destitute of any soil formed from the disintegration of the rocks, has remained comparatively barren, in spite of its closer proximity to the mainland.

The most interesting things we have found up to this time are representatives of the *Ceratis* group of Fishes, which the naturalists of the *Albatross* tell me they have not met before on the west coast of North America. The Crustacea have supplied us with a most remarkable type of the Willemoesia group. The paucity of Mollusks, and also of Echini, is most striking, although we brought up in one of the hauls numerous fragments of what must have been a gigantic species of *Cystechinus*, which I hope I may reconstruct. We were also fortunate enough to find a single specimen of *Calamocrinus* off Morro Puercos, in 700 fathoms, a part of the stem with the base,

showing its mode of attachment to be similar to that of the fossil Apicrinidae. The number of Ophiurans was remarkably small as compared with the fauna of deep waters on the Atlantic side, where it often seems as if Ophiurans had been the first and only objects created. The absence of deep-sea corals is also quite striking. They play so important a part in the fauna of the deeper waters of the West Indies, that the contrast is most marked. Gorgoniæ and other Halcyonoids are likewise uncommon. We have found but few Siliceous Sponges, and all of well-known types. Star-fishes are abundant, and are as well represented in the variety of genera and species as on the Atlantic side of the Isthmus. I may also mention the large number of deep-sea Holothurians (Elasipoda) which we obtained, as well as a most remarkable deep-sea Actinian, closely allied to *Cerianthus*, but evidently belonging to a new family of that group. We found the usual types of deep-sea West Indian Annelids, occasionally sweeping over large tracts of mud tubes in the region of green mud. Although we dredged frequently in most characteristic Globigerina ooze, I was much struck with the absence of living Globigerinæ on the surface. Only on two occasions during a calm did we come across any number of surface Globigerinæ and Orbulinæ. On one occasion the trawl came up literally filled with masses of a species of *Rhabdamina* closely allied to *R. lineata*. Thus far, no pelagic Algae have been met with.

It is interesting to note that at two localities we came across patches of modern greensand similar in formation to the patches discovered off the east coast of the United States by the earlier dredgings of the Coast Survey, of Pourtales, and of the *Blake*. Having always been more or less interested in pelagic fauna, and having paid considerable attention to its vertical distribution during my earlier cruises in the *Blake*, I was naturally anxious to reconcile the conflicting statements and experiences of the naturalists of the *Challenger* and *Gazelle* on one side, and my own observations on the other. Both Murray and Studer contended that, in addition to the deep-sea and pelagic fauna, there was what might be called an intermediate fauna, with characteristic species, having nothing in common with the other two; while I maintained, on the other hand, from my experiments in the *Blake*, that there was no such intermediate fauna, but that the pelagic fauna might descend to a considerable depth during the daytime to escape the effects of light, heat, and the disturbing influence of surface winds, and that this surface fauna on the Atlantic side—off shore in deep water—did not descend much deeper than 150 to 200 fathoms. In order to test this point, Dr. Chun, under the auspices of the Naples Station, made an expedition to the Ponza Islands. Dr. Chun applied to a tow-net an apparatus for closing it, similar to the propeller in use on our thermometer and water-cups. He towed to a depth of 1400 metres, if I am not mistaken, but never at any great distance from the mainland or from the islands of the Gulf of Naples, and came to the conclusion that the pelagic fauna existed all the way to the bottom. At the time, I considered his experiments inconclusive, and was, of course, anxious to repeat them in a strictly oceanic district, in great depths, and at a considerable distance from shore. I had an apparatus constructed by Ballauf, of Washington, similar to that used by Dr. Chun. Unfortunately, in testing it we found the pressure of the tow-net against the propeller shaft so great as to make the machine useless, or, at any rate, most unreliable. Thanks to the ingenuity of Captain Tanner, we overcame these obstacles. He devised a net which could be closed at any depth by a messenger, and which worked to perfection at 200, 400, 300, and 1000 fathoms, and had the great advantage of bringing up anything it might find on its way up above the level at which it was towed. The lower part of the bag alone was closed by a double set of slings pulled by two weights liberated from a bell crank by a messenger. We found that, in towing the net at 200 fathoms for twenty minutes, we got everything in any way characteristic of the surface fauna which we had fished up with the tow-net at the surface. In addition to this, we brought up five species of so-called deep-sea Fishes, *Scopelus*, *Gonostoma*, *Beryx*, and two others, which had thus far been brought up in the trawl, and considered characteristic of deep water. Also a peculiar Amphipod, and the young of the new species of *Willemoesia* mentioned above. We then tried the same net at 300 and 400 fathoms, and in neither case did we bring up anything in the closed part of the bag, while the upper open part brought up just what we had found previously at a depth of 200 fathoms, plainly showing that in this district the surface fauna goes down

to a depth of 200 fathoms, and no farther. Next came our single attempt to bring up what might be found, say within 100 fathoms of the bottom, and Captain Tanner's net was towed at a depth of 1000 fathoms where the soundings recorded 1100. Unfortunately, we deepened our water while towing only twenty minutes to over 1400 fathoms, so that we failed in our exact object. But we brought up in the closed part of the bag two species of Crustacea, a Macruran and an Amphipod, both entirely unlike anything we had obtained before. I hope in the next cruise to follow this up, and determine also the upper limits of the free-swimming deep-sea fauna. In the upper part of the bag (the open part) we brought up a couple of so-called deep-sea Medusæ, which must have been collected at a comparatively moderate depth, judging from their perfect state of preservation.

I can hardly express my satisfaction at having the opportunity to carry on this deep-sea work on the *Albatross*. While of course I knew in a general way the great facilities the ship afforded, I did not fully realize the capacity of the equipment until I came to make use of it myself. I could not but contrast the luxurious and thoroughly convenient appointments of the *Albatross* with my previous experiences. The laboratory, with its ingenious arrangements and its excellent accommodations for work by day and by night, was to me a revelation. The assistance of Messrs. Townsend and Miller in the care of the specimens was most welcome, giving me ample time to examine the specimens during the process of assorting them, and to make such notes as I could between successive hauls, while paying some attention also to the work of the artist, Mr. Westergren. He has found his time fully occupied, and we have in this trip brought together a considerable number of coloured drawings, giving an excellent general idea of the appearance of the inhabitants of the deep waters as they first come up. These drawings can be used to great advantage with the specimens in making the final illustrations to accompany the reports of the specialists who may have charge of working up the different departments. . . .

We left Panama on February 22, and returned to Panama after an absence of twenty days.

II.

"*Albatross*," *Acapulco*, April 14, 1891.

We have reached the end of our second line of explorations. After coaling we left Panama, and reached Galera Point, where we began our line across the Humboldt Current, which was to give us a fair idea of the fauna of that part of the coast as far as the southern face of the Galapagos. With the exception of three good casts, the trawling on that part of the sea bottom proved comparatively poor, nor did the sea face of the southern slope of the Galapagos give us anything like the rich fauna I had expected. Theoretically, it seemed certain that a sea face like that of the Galapagos, bathed as it is by a great current coming from the south and impinging upon its slope, and carrying upon its surface a mass of animal food, could not fail to constitute a most favourable set of conditions for the subsistence and development of a rich deep-sea fauna.

In the deeper parts of the channel between Galera Point and the southern face of Chatham Island, we found a great number of *Elasipoda*, among them several genera like *Peniagone*, *Bathodytes*, and *Euphrosyne*, represented by numerous species. The Star-fishes of this, our second cruise, did not differ materially from those collected during our first trip, but we added some fine species of *Freyella*, *Hymenaster*, *Astrogonium*, *Asterina*, and *Archasteridae* to our collections. Among the Sea-urchins on two occasions we brought up fine hauls of a species of *Cystechinus* with a hard test, many specimens of which were in admirable state of preservation. Among the Ophiurans nothing of importance was added, unless I may except a lot of Ophiocreas attached to a *Primnoa*, and a pretty species of *Sigsbea* attached to a species of *Allopora*, from the south side of Chatham Island.

The Gorgonians were remarkably few in number, which is undoubtedly due to the unfavourable nature of the bottom we worked upon. Nearly everywhere except on the face of the Galapagos slope we trawled upon a bottom either muddy or composed of Globigerina ooze, more or less contaminated with terrestrial deposits, and frequently covered with a great amount of decayed vegetable matter. We scarcely made a single haul of the trawl which did not bring up a considerable amount of decayed vegetable matter, and frequently logs, branches, twigs, seeds, leaves, fruits, much as during our first cruise.

Our Crustaceans, from the nature of the bottom, naturally consisted of the same groups of deep-sea types which we obtained before. I may, however, mention a haul containing a goodly number of *Nephrops*, a genus we had not previously obtained.

Among the Worms the Maldanæ and Limicolous types were unusually abundant at some localities, the empty mud tubes often filling the bottom of the trawl. Some very large specimens of *Trophonia* were collected, and remarkably brilliantly coloured (orange and carmine) Nemerteans and Planarians.

The Mollusks were very scanty, and the absence of Comatulæ or other Crinoids was equally disappointing, even when trawling on the extension of the line started three years ago by the *Albatross*, on the eastern face of the Galapagos slope, when on her way from Chatham Island to San Francisco. We took up this line off Indefatigable Island, hoping to obtain from that quarter our best results, but our hauls were very disappointing. The ground proved not only most difficult to dredge upon, but also comparatively barren, and it was not till we got into the oceanic basin again, between the Galapagos and Acapulco, that our catches improved. But even then they were not to be compared with the hauls at similar depths in the Atlantic off the West Indies, or along the course of the Gulf Stream.

Among the Fishes, our most important catches were fine specimens of *Bathyonus*, of *Bathybrissa*, of *Bathypteroides*, and a few specimens of *Ipnops* in excellent condition.

From the nature of the bottom we naturally expected rich hauls of Siliceous Sponges, but we did not find many, and I do not think there were many novelties among those we have collected. On two occasions, a number of specimens of Ascidians were brought up; among them was a fine white translucent *Corinascidia*.

Among the Bryozoans, the most noteworthy haul was a number of beautiful specimens of the delicate *Naresia*, in excellent condition. On the line from the Galapagos to Acapulco we brought up a good many Foraminifera from the mud bottoms. On several occasions the bottom must have been covered with huge masses of a new type of an arenaceous Foraminifer, forming immense curling sheets attached by one edge to stones or sunk into the mud. This Foraminifer seems to increase in size by forming irregular more or less concentric crescent-shaped rings. When it comes to the surface, it is of a dark olive-green colour.

During this second cruise we continued our experiments with the Tanner closing tow-net, in order to determine the lower limits of the surface pelagic fauna, and to determine also if there is any so-called intermediate pelagic fauna at other depths, or within a short distance from the bottom.

On March 25, at a point not quite half-way between Cape San Francisco and the Galapagos, in 1832 fathoms of water, the Tanner net was sent down to tow at a depth which varied from 1739 to 1773 fathoms. The net was towed within these limits for a period of something over twenty minutes. The messenger was then sent down to close the net; time occupied seven minutes. The net was then drawn up to the surface. The lower part of it was found to have closed perfectly, and contained nothing beyond a few fragments of leaves. The lower bag was carefully washed in water which had been strained, and the water examined with all possible care, and sifted again. It contained nothing. The upper part of the net, however, which had remained open on its way up, was found to contain the identical surface things which on former occasions we had found in the Tanner net down to a depth of 200 fathoms. They were a small species of *Sagitta*, and species of *Doliolum*, *Appendicularia*, a huge *Sagitta*, a large number of *Leucifer* and *Sergestes*, and several species of *Schizopods* and *Copepods*; two species of *Hyperia*, probably parasitic on a *Salpa*, which was also quite abundant; several finely coloured *Calanus*, some *Isopods*, and a number of fragments of what must have been a very large *Beroë*, measuring from five to six inches in diameter; *Leptocephalus*, several specimens of *Stomias*, of *Scopelus*, of *Melamphæx*, and other species, many of which, like some of the *Schizopods*, had been considered as typical deep-sea forms. Among the so-called deep-sea *Medusæ*, several specimens of *Atolla* and *Periphylla* were also found in the open part of the net. I may mention also as of special interest a huge *Ostracod*, allied to *Crossophorus*, with a thin semi-transparent carapace, and measuring somewhat more than one inch in length. The largest *Ostracod* previously known is not more than one-third of an inch long. On two other occasions this same *Ostracod* was brought up in the tow-net from a depth of less than 200 fathoms.

The surface at this point was also examined with the tow-net, and the pelagic animals found to be the same as those brought up in the open part of the tow-net on its way from the bottom. The number both of species and specimens was, however, much less than in the Tanner net. On the following day the Tanner tow-net was sent to be towed at a depth of 214 fathoms. In twenty minutes the messenger was sent down and the net hauled up. The bottom part of the net came up tightly closed. Its contents were examined in the same manner as before in well sifted water, and the water was found to be absolutely barren, while the upper part of the net, which came up open, and was not more than eight or nine minutes on the way, was well filled with surface life. The net contained this time a number of *Hyaleas* and *Criseis*, in addition to the things collected the day before. An examination of the surface fauna at this same point with the tow-net showed the presence only in smaller numbers of the same species which the open part of the same net contained, except that there were a larger number of bells and fragments of *Diphyes* and of *Cristalloides* than in the Tanner net. The point at which this experiment was made was about 250 miles from the Galapagos, and about the same distance from Cape San Francisco. There were myriads of *Nautilograpsus* swarming on the surface of the water; they literally filled the surface tow-net. On two other occasions, once at a distance of 350 miles in a south-easterly direction from Acapulco (depth 2232 fathoms), we tried the same experiment with the Tanner net, and invariably with the same result. The net was towed at a depth of 100, of 200, and of 300 fathoms, each time for twenty minutes, the messenger sent down, and the bottom part closed. At the depth of 100 fathoms, the closed part of the net contained practically the same things as the open part of the net; at 200 fathoms, the lower part of the net contained but few specimens of the surface life; and at 300 fathoms, the closed bottom net came up empty.

On the following day the surface was carefully examined, and the tow-net sent to 175 fathoms, where it was towed for twenty minutes, and the messenger sent down to close it. The lower net came up well filled with the surface pelagic species, which on this day were unusually varied, it having been smooth and calm the previous night, and the morning before the towing was made. This haul was made in the evening, at 8 p.m. The previous hauls had been made at about 10 a.m., in a brilliant sunlight. Again, on April 11, about thirty miles south-east of Acapulco, in a depth of over 1800 fathoms, the Tanner net was sent to a depth of 300 fathoms, and the messenger sent down to close it. There was nothing in the lower part of the net which had been closed, while the open part contained an unusually rich assortment of surface species, and among them a large number of *Scopelus*, of *Schizopods*, and of *Rhizopods*, mainly *Collozoun* and *Acanthometra*.

These experiments seem to prove conclusively that in the open sea, even when close to the land, the surface pelagic fauna does not descend beyond a depth of 200 fathoms, and that there is no intermediate pelagic fauna living between that depth and the bottom, and that even the free-swimming bottom species do not rise to any great distance, as we found no trace of anything within 60 fathoms from the bottom, where it had been fairly populated.

The experiments of Chun regarding the distribution of the pelagic fauna have all been made in the Mediterranean, within a comparatively short distance from the shore, and in a closed basin showing, as is well known, special physical conditions, its temperature to its greatest depths being considerably higher than the temperature of oceanic basins at the limit of 200 fathoms, or thereabout, which we assume now to be the limit of the bathymetrical range of the true oceanic pelagic fauna. At 200 fathoms our temperature was from 49° to 53°, while, as is well known, the temperature of the Mediterranean soon falls at 100 fathoms even to about 56°, a temperature which is continued to the bottom in this closed basin. Of course, if temperature is one of the factors affecting bathymetrical distribution, there is no reason except the absence of light which would prevent the surface pelagic fauna from finding conditions of temperature at the greatest depth similar to those which the surface fauna finds within the limit of 200 fathoms in an open oceanic basin.

Arriving as we did at the Galapagos at the beginning of a remarkably early rainy season, I could not help contrasting the green appearance of the slopes of the islands, covered as they were by a comparatively thick growth of bushes, shrubs, and

trees, to the description given of them by Darwin, who represents them in the height of the dry season as the supreme expression of desolation and barrenness. Of course, here and there were extensive tracts on the sea-shore where there was nothing to be seen but blocks of volcanic ashes, with an occasional cactus standing in bold relief, or a series of mud volcanoes, or a huge black field of volcanic rocks, an ancient flow from some crater to the sea; but as a rule the larger islands presented wide areas of rich fertile soil, suitable for cultivation. The experiments at Charles Island, where there is a deserted plantation, and at Chatham Island, where Mr. Cobos has under successful cultivation a large plantation producing sugar, coffee, and all the tropical fruits, as well as extensive tracts on which his herds of cattle, sheep, and donkeys roam towards the higher central parts of the island, show the fertility of these islands. They are indeed as favourably situated for cultivation as the Sandwich Islands or Mauritius, and there is no reason why, if properly managed, they should not in the near future yield to their owners as large returns as do those islands.

I obtained from Mr. Cobos a piece of the so-called sandstone, said to occur on Indefatigable Island, and which of course I was most anxious to see, as the occurrence of true sandstone would have put quite a different face on the geological history of the Galapagos from the one usually received. This I found to be nothing but coral rock limestone, either a breccia, or slightly oolitic, identical with the formation found back of the beach at Wreck Bay on Chatham Island. I found there an old coral rock beach, extending on the flat behind the present beach, composed entirely of fragments of corals, of mollusks, and other invertebrates, cemented together into a moderately compact oolitic limestone, which when discoloured, as it often is, and turned gray, would readily be mistaken for sandstone. This coral rock is covered by just such a thin, ringing coating of limestone as characterizes the modern reef rock of other localities. On nearly all the islands there are a number of sandy beaches made up of decomposed fragments of corals and other invertebrates, and cemented together at or beyond high-water mark into the modern reef rock I have described. The coral is mainly made up of fragments of *Pocillopora*, which is found covering more or less extensive patches off these coral sand beaches, but which, as is well known, never forms true coral reef in the Panamic district. The only true coral reef belonging to this district is that of Clipperton Island (if we can trust the Admiralty charts), situated about 700 miles to the south-west of Acapulco. But neither at Cocos Island, nor at the Galapagos, nor anywhere in the Panamic district, do we find true coral reefs—nothing but isolated patches of reef-building coral. The absence of coral reefs in this district has of course already been noted by other naturalists, who have been struck by this feature in an equatorial region. Dana has ascribed it to the lower temperature of the water due to the action of the Humboldt current coming from the south, pouring into the Bay of Panama, and then flowing westward with the colder northerly current coming down the west coast of Mexico and Central America. From the investigations made this year by the *Albatross*, I am more inclined to assume that the true cause of the absence of coral reefs on the west coast of Central America is due to the immense amount of silt which is brought down the hill and mountain sides every rainy season, and which simply covers the floor of the ocean to a very considerable distance from the land, the land deposits being found by us even on the line from the Galapagos to Acapulco at the most distant point from the shore to the side or extremities. The mud in Panama Bay to the hundred-fathom line is something extraordinary, and its influence on the growth of coral reefs is undoubtedly greatly increased from the large amount of decomposed vegetable matter which is mixed with the terrigenous deposits.

The course of the currents along the Mexican and the Central and South American coasts clearly indicates to us the sources from which the fauna and flora of the volcanic group of the Galapagos has derived its origin. The distance from the coast of Ecuador (Galera Point and Cape San Francisco) is in a direct line not much over 500 miles, and that from the Costa Rica coast but a little over 600 miles, and the bottom must be for its whole distance strewn thickly with vegetable matter. The force of the currents is very great, sometimes as much as 75 miles a day, so that seeds, fruits, masses of vegetation harbouring small reptiles, or even large ones, as well as other terrestrial animals, need not be afloat long before they might safely be

landed on the shores of the Galapagos. Its flora, as is well known, is eminently American, while its fauna at every point discloses its affinity to the Mexican, Central or South American, and even West Indian types, from which it has probably originated; the last indicating, as well as so many of the marine types collected during this expedition, the close connection that once existed between the Panamic region and the Caribbean and Gulf of Mexico.

I have already referred to the physiognomy of the deep-sea fauna, showing relationship on the one side to Atlantic and West Indian types, and on the other to the extension of the Pacific types, which mix with the strictly deep-sea Panamic ones. The western and eastern Pacific fauna, while as a whole presenting very marked features in common, yet also present striking differences. The vast extent of territory over which some of the marine types extend, through all the tropical part of the Pacific, may readily be explained from the course of the great western equatorial current and the eastern counter current, which cannot fail to act as general distributors in space for the extension of a vast number of marine Vertebrates and Invertebrates.

Mr. Townsend made quite a large collection of birds from Chatham and Charles Islands, considering the short time we were there.

As soon as we have reached Guaymas, I shall be able to give you a better *résumé* of the character of the deep-sea fauna of the Panamic region, and of its relationship on the one side to the Pacific fauna and on the other to the West Indian region.

III.

Guaymas, April 25, 1891.

We left Acapulco on April 15 for our third cruise into the Gulf of California, and steamed as far as Cape Corrientes without attempting to do any trawling. The character of the bottom, as indicated on the charts, promised nothing different from what we had dredged off Acapulco, and on the line from there to the Galapagos Islands. We made one haul off Cape Corrientes, bringing up nothing but mud and decomposed vegetable matter. This induced us to keep up the Gulf of California till we were off the Tres Marias. We there made several hauls, and obtained some *Umbellulæ*, *Pennatulæ*, *Trochophilum*, *Anthopitulum*, and a fine *Antipathes*, a few *Comatulæ*, a large *Astropecten*, some fine specimens of *Urechinus* and of *Schizaster*, a few *Holothurians*, *Lophothuria*, *Trochostoma*, and two species of *Elasipoda*, besides a few fragments of *Gasteropods*, with an empty shell of *Argonauta*.

Among the Crustacea there came up the usual types found living upon muddy bottom, such as *Glyphocrangon*, *Heterocarpus*, *Notostoma*, *Pentacheles*, *Nematocarcinus*, *Nephrops*, together with species of *Lithodes* and of *Munida*. The usual types of *Limicolous Annelid* also were found here, *Halinæcia*, *Terebella*, *Maldania*, and the like, a few *Ophiurans*, *Ophiopholis*, and *Ophiocantha*, a few fragments of *Farrea*, and a huge *Hyalonema* of the type of *H. toxeres*. Among the Fishes there were a few *Macrurans*, *Bathypetrolides*, *Lycodes*, and *Malthe*. The trawl was usually well filled with mud, and with the mud came up the usual supply of logs, branches, twigs, and decayed vegetable matter.

On going farther north into the Gulf of California, the nature of the bottom did not change materially, and we found the trawling most difficult from the weight of the mud brought up in the trawl. But occasionally a haul was made which more than repaid us for the time spent on the less productive ones. Two of the hauls are specially worthy of mention, as being characteristic of the deep-water fauna of the Gulf of California, one made in 995 fathoms, and the other in 1588 fathoms. We obtained in these hauls a number of *Ophiomusium* and *Ophiocreas*, some fine specimens of *Schizaster*, a new genus allied to *Palæopneustes*, and also the same species of *Cystechinus*, with a hard test, and of *Phormosoma*, which we had obtained before on the line from the Galapagos to Acapulco. Beside these there came up a number of specimens of an interesting species of *Pourtalesia*, most closely allied to *Pourtalesia miranda*, the first type of the group dredged in the Florida Channel by Count Pourtales.

The deeper haul was specially rich in *Holothurians*, among them a fine large white *Cucumaria*, some specimens of *Trochostoma*, several species of *Bathodytes*, some of them remarkable for their white colour, their huge size, and comparatively small

number of ventral tentacles. With these were numerous specimens of an interesting species of Euphronides. In this haul I was specially struck with the Elapipoda, and the great variety in the consistency of the skin in individuals of one and the same species; it varied in different individuals from extreme tenuity to a comparatively tough gelatine-like consistency. On carefully sifting the mud, we found a number of interesting Foraminifera, and of delicate and minute Gasteropods and Lamellibranchs, fragments of the shell of an Argonauta, and two species of a huge ribbed Dentalium. Among the Star-fishes were specially noticeable a large Brisinga, a long-armed Cribrella, and several species of Astropecten. The usual types of Worms were found in the mud at these greater depths. In addition to a number of Macrurids, we obtained a pink Amphionus, a large black Beryx-like fish, a fine Nettastoma, and a couple of species of Lycodes. The usual surface species of Stomias and of Scopelus also came up in the trawl. Among the Crustaceans were a fine lot of Arcturus, of Colossendeis, of Glyphocrangon, and of a Caridid with a deep blue patch on the base of the carapace, making the strongest possible contrast to the dark crimson colouring of the rest of the body. Blue is a very unusual colour in the deep-sea types, although the large eggs of some of the deep-sea Macrurans are often of a light blue tint.

We brought up in the trawl at various times, and subsequently also in the Tanner net, from depths of less than 200 fathoms, the same gigantic Ostracod which I mentioned in one of my previous letters, several specimens of Atolla, and fragments of a huge Periphylla, which must have been at least 15 inches in diameter. Also a most interesting new type of Bougainvillia, remarkable for having eight clusters of marginal tentacles, but only four chymiferous tubes.

We continued our experiments with the Tanner tow-net. On April 16, about 120 miles from Acapulco, we sent the net to tow at a depth of 175 fathoms, and after towing for about twenty minutes, sent the messenger to close it. On examining the bottom part of the net, which came up tightly closed, we found it to contain practically the same things as we obtained in the surface net at the same spot.

On two occasions we sent the net to be towed at depths of 800 fathoms and of 700 fathoms, the depths at these points being in one case 905 fathoms and in the other 773 fathoms. At the greater depth, the water shoaled somewhat while towing, as the closed part of the net came up partly filled with fine silt; while during the second haul, the twisting of the swivel wound the straps of the weights round the rope, and the net came up open, but must have dragged very close to the bottom, as it contained a fine specimen of Nettastoma, and some Penæids, which we supposed to be deep-sea types. Otherwise the net contained only the customary surface species of Sagitta, Pteropods, Copepods, Schizopods, Tunicates, and Fishes. These two hauls were made about the middle of the Gulf of California, at a distance of some fifty miles in a south-westerly direction from Guaymas.

On April 23, a few hours before reaching Guaymas, we made one more attempt with the Tanner tow-net, at a depth of 620 fathoms, sending the net to be towed at a depth of from 500 to 570 fathoms. We found in this case in the bottom part of the net, which came up tightly closed, a Scopelus, a Penæid, and a Hyalea, while the upper open part of the net contained the same surface species we had obtained before.

My experience in the Gulf of California with the Tanner self-closing net would seem to indicate that in a comparatively closed sea, at a small distance from the land, there may be a mixture of the surface species with the deep-sea bottom species, a condition of things which certainly does not exist at sea in an oceanic basin at a great distance from shore, where the surface pelagic fauna only descends to a comparatively small depth—about 200 fathoms—the limits of the depth at which light and heat produce any considerable variation in the physical condition of the water. The marked diminution in the number of species below 200 fathoms agrees fairly with the results of the *National Expedition*.

The more I see of the *Albatross* the more I become convinced that her true field is that of exploration. She is a remarkably fine sea boat, and has ample accommodation for a staff of working specialists such as would be needed on a distant expedition. The time will soon come when the Fish Commission will hardly care to continue to run her, and I can conceive of no better use for so fine a vessel than to explore a belt of 20° latitude north

and south of the equator in the Pacific, from the west coast of Central America to the East Indian Archipelago.

The success of the *Albatross* thus far has depended entirely upon the zeal, energy, intelligence, forethought, and devotion of Captain Tanner, if I may judge of the past by the present. He never spares himself, and he is always ready to make the most of the time at his disposal for the benefit of the special object he has in charge. He looks after every haul of the trawl himself, and will not allow anyone else to jeopard in any way the material of the vessel, or the time it requires to make a haul. That responsibility he assumes himself, and it constitutes his daily work. In looking over the records of the *Albatross* during her voyage from New York to San Francisco, I am struck with the amount of work which has been accomplished. It would be but a just return to Captain Tanner if Congress would make the necessary appropriations to work up and publish all that he has brought together, not only on that cruise, but also what has been left untouched thus far of the immense collections made by him in the Caribbean, and off the east coast of the United States, to say nothing of his explorations in the Gulf of California, on the coast of California, on the coast of Alaska, and in the Behring Sea, from which he has accumulated endless and most interesting material, which no other ship could get together unless she had another Tanner in command.

We reached Guaymas on April 23, in the afternoon, and I parted from the ship with great regret, but more than satisfied with the results of this expedition.

Allow me, in concluding, to thank you most cordially for having given me the opportunity to join the *Albatross* on this extended cruise, and for your kindness in urging the President to allow the vessel to be detailed for this work.

As soon as it may become practicable, I shall send you a full *résumé* of our work, accompanied with sketches of the Tanner tow-net and a detailed chart of the route we followed.

Very respectfully yours,

ALEXANDER AGASSIZ.

THE ORIGIN OF THE ASS, THE CAT, AND THE SHEEP IN CHINA.

AT a recent meeting of the China Branch of the Royal Asiatic Society in Shanghai, Dr. Macgowan, a well-known Chinese scholar, read a paper on the probable foreign origin of the ass, the cat, and the sheep in China. He said that the Chinese, in their numerical co-ordination of concrete and abstract nature, give the "six domestic animals" as the horse, ox, goat, pig, dog, and fowl; which seems to indicate that when that formula was framed, neither cat, sheep, nor ass had been domesticated there. When familiar beasts were selected to denote years of the duodenary cycle, to the "six domestic animals" were added the rat, tiger, hare, dragon, serpent, and monkey, to complete the dozen, as if the ass, sheep, and cat were too little known to meet the object in view, which was the employment of the most familiar representations of animated nature for the duodenary nomenclature. Still more striking is the absence of the ass, sheep, and cat from the twenty-eight zodiacal constellations, which are represented by the best-known animals.

With regard to the ass, there is ample reason to regard it as being excluded from the list of domestic animals because it was not archaic. The hybrid mule is of comparatively modern origin in China, dating back only about a score of centuries. A miscellany of the Sung era states that "the mule was not seen during the Hsai, Shang, and Chou dynasties; that it was a cross between the ass and horse from Mongolia. It is regularly bred in the north, and is worth in the market twice as much as the horse; it is popularly reported that its bones are marrowless, which is the reason of its inability to produce its kind." Again, it is recorded in a Ming cyclopædia: "The mule is stronger than the horse, and is not a natural product of China; in the Han era it was regarded as a remarkable domestic animal." Is it likely that, if the ass existed during the three ancient dynasties, there was no crossing with the horse?

With regard to the cat, Dr. Macgowan proceeded to state that there was a quotation from a standard work which dis-