under consideration the size of the image thrown on the screen can be varied at will by simply altering the distance between the elements, but the further the lens is from the focussing screen, the more will be the time of exposure.

With such a lens as this Mr. Dallmeyer has taken many excellent pictures, but perhaps the best idea of its properties will be gathered from the facts obtained by photographing—by means of two cameras, one supplied with a “long focus landscape lens,” and the other with the “new telephotographic lens”—the flame of an oil lamp placed at a distance of 20 feet. With equal extensions of the camera the image of the flame given by the new lens was five times greater than that by the other.

In the compound lens the anterior element before referred to is here replaced by a complete portrait lens, while a negative symmetrical combination takes the place of the posterior element. This lens may be said to be more perfect than the simple lens, Mr. Dallmeyer having been able to introduce considerable improvement in the construction.

Some excellent work done with this lens has been exhibited by Messrs. F. Mackenzie and Annan at the Camera Club. The pictures represented a building at a distance of 500 yards. The first, taken with an ordinary rapid rectilinear lens with an extension of 14 inches, gave the house as $\frac{3}{4}$ of an inch long. The second—with the compound tele-photo lens, extension 9 inches from the back lens—gave $2\frac{1}{2}$ inches as the size of the house, while the third, with 30 inches' extension, gave the house as $6\frac{1}{2}$ inches. Although these numbers can give one a very good idea of what this new lens can accomplish, yet the direct copies from photographs inserted in the pamphlet under consideration convey a more vivid impression.

There is no doubt that this lens will find some very valuable applications, that of astronomical photography not being the least of them, for every one knows the great advantage a short telescope has over a long one if the degree of magnification in both are equal. W.

ARBORESCENT FROST PATTERNS.

WE have received the following letters with regard to arborescent frost patterns, to which Prof. Meldola called attention in last week's NATURE:—

I AM very glad that Prof. Meldola has called attention to the curved figures of frozen mud (of which the specimens on December 4 were unusually fine), because I hope that some one will explain why the sexangular crystallization which is universal in snow, and general in water, is exchanged both on windows and on muddy pavements for curves. Probably I ought to know all about it, but I cannot remember seeing an explanation, and shall be obliged by reference to one, which will probably be of interest to others besides G. J. SYMONS.

62, Camden-square, London, N.W.

THE interesting “fronds” of muddy ice observed by Prof. Meldola (p. 126) are not very uncommon on the pavements in these “Northern Heights.” I saw them on the date which he named, and have more than once studied them. I then noticed that the “interstitial” pavement seemed partly cleared of mud, as if the water had drawn this towards the groups of crystals. The mode of formation recalled to my mind certain phenomena in crystal building within rocks, and I suspect the mud has its influence. Indeed, it seems to me very probable that all these “dendritic” growths of crystals are the results of “impeded” or “constrained” crystallization, to some of which I have called attention in noticing a structure in the Charnwood syenite (*Quart. Jour. Geol. Soc.*, 1891, p. 101). On this point Prof. Sollas makes some important remarks in his well-known paper on the Wicklow granites. T. G. BONNEY.

THE beautiful curved forms assumed by the ice on the paving flags last Sunday were very noticeable in this neighbourhood and Hampstead as well as in other parts of London. What I observed were not quite like those described and figured by Prof. Meldola, but resembled rather the scrolls and volutes which are frequently used in decorative art. The finest piece that I saw was in this square, where several of these scrolls radiated from a central point, and spread over several feet of the pavement. A friend, Mr. E. Swain, observed that where one of these scrolls came upon a puddle of clear water the crystals were continued in a straight line. Such forms are not at all unusual in the freezing of muddy water, and at the present moment the puddles in the road opposite my house are

filled with rectilinear crystals of ice, which assume a curved form in the mud at their margins. The peculiarity on Sunday was their large size and beauty. Something analogous takes place when gold or silver is reduced from solutions of its salts by more electro-positive metals. Under certain circumstances

the metal will present itself in the form of curved crystals, if the term be allowable. A pretty spray of gold of this character is figured in the report of my lecture “On the crystallization of silver, gold and other metals,” in the proceedings of the Royal Institution, vi., 428. If a piece of caprous oxide be immersed in a solution of nitrate of silver, there shoot from its surface thin threads of silver, which, after proceeding straight forward for a while, suddenly turn at an angle of 120° or 60° , and make perhaps many other deviations: but sometimes these threads, instead of being straight, are curved; and in that case the threads that branch from them are curved likewise. A magnified drawing of such a formation is given herewith. These strange departures from the usual rectilinear course of crystal formation are very curious, and deserve more study than has hitherto been given them. J. H. GLADSTONE.

17, Pembroke-square, December 10.

PROF. MELDOLA'S letter (p. 125) has been interesting to me, as I noted a striking and similar phenomenon here on Thursday, December 8, in the forenoon. The trottoirs of several streets (east, west, north and south) were covered all over with beautiful patterns, somewhat different from Prof. Meldola's illustration, there being innumerable dark, broad, sharply-contoured leaf-like patches, distant several inches from each other, and connected by finely curved and branched tendril-like stalks. Foggy, with a faint north breeze. I should presume the “leaves” were due to sparse drops of sleet fallen during the night. Freiburg, Badenia, December 10. D. WETTERHAN.

THE graceful arborescent frost patterns described by Prof. Meldola in last week's NATURE were very conspicuous on the foot-bridge by the side of Charing Cross railway bridge, on the same morning, this being a situation still more exposed to the wind which he mentions as the probable cause.

December 12.

J. T. RICHARDS.

I OBSERVED the same phenomenon as Prof. Meldola describes in NATURE of December 8, on the same date, December 4, on pavements in Cheltenham, about 10.45 a.m.; after mid-day they had gone. I saw the patterns on pavements running north and south, as well as east and west. They were most exquisite; some like the illustration, others much more minute; but always in a connected design over the whole flag. They had all the appearance of fossil vegetation. I never saw anything of the kind before. J. J. ARMITAGE.

December 13.

MR. A. W. BENNETT and Mr. E. L. Garbett have also sent communications corroborating the phenomenon observed by Prof. Meldola. The former attributes it to “defoliation of the stones as the result of weathering or wear.”