stereo isomerism; the essential part of it is retained in the assumption that the carbon atom is stereo-chemically to be regarded as tetrahedral in shape. If any student carries away the notion that this is believed to be the actual shape of the atom, there is no more mischief done than in that student's case who gathers the impression that the two carbon atoms united by an ethylene linkage are held together by two pairs of forces which do not act along the line joining the two atoms, but meet at an angle in empty space. G. S. TURPIN.

Huddersfield, September 24.

Careless Writing.

PROF. HILDEN, in his review, published in NATURE, September 20, takes exception to the loose phraseology adopted by writers on chemical subjects. This is, alas! only too common.

For example, in one of the best works on inorganic chemistry, written by a Professor of Chemistry whose writings are characterised by their logical clearness and philosophic reasoning, one may read:—

"When a molecule of hydrogen acts upon a molecule of chlorine to form two molecules of hydrochloric acid gas, 44,000 c. of heat are evolved." Of course, nobody can fail to understand what is meant. But as the words stand, it is certainly one of the most remarkable feats of science, and makes us feel that some happy mortal has succeeded in refining his faculties down to the degree of fineness, popularly ascribed only to a certain species of "demon." F. G. DONNAN.

Holywood, Belfast.

ON THE DOCTRINE OF DISCONTINUITY OF FLUID MOTION, IN CONNECTION WITH THE RESISTANCE AGAINST A SOLID MOVING THROUGH A FLUID.¹

II.

§ 6. IN every case in which vacuum is formed at an edge of a solid moving in an inviscid incompressible fluid, under pressure constant at all infinitely great distances from the solid, a succession of finite individual vortices is sent from the edge into the liquid, and the motion is essentially unsteady. Each individual vortex has a finite endless vacuum for its core instead of the rotationally moving ring of fluid of the Helmholtz vortex ring. But it should be noticed that it would not be rings of vacuum, but bubbles, that would in many cases be first detached from the solid; that by the tumultuous collapse of bubbles they become rings; and that the case in which the collapse of a bubble, in our ideal fluid, could be completed to an annulment of volume, is of necessity infinitely rare; and that the case in which, when a bubble becomes a ring by the meeting of two points of collapsing boundary, there is exactly no circulation through the aperture, is infinitely rare.²

§ 7. In the case of our circular disc, it would be circular vortex rings that, if the water were inviscid, would be shed off from its edge when the depth is less than 63 feet. If the depth is very little less than 63 feet these rings would be exceedingly fine, and would follow one another at exceedingly short intervals of time. Thus quite close to the edge there would be something somewhat like Stokes' "rift," but with a rapid suc-

cession of vacuum rollers, as it were; and no slipping between the portions of the fluid on its two sides.

§ 8. At greater depths than 63 feet, if the water had absolutely no viscosity,¹ the motion would be continuous and irrotational, as described in § 4, text and foot-note: but any degree of viscosity, however slight, would, if the edge were infinitely sharp (instead of having a radius of curvature of $1/2000$ of an inch, as has our supposed disc), give rise to a state of motion in its neighbourhood somewhat like to Stokes' rift,² "a surface of discontinuity extending some way into the fluid," but with the difference that there is no slip of fluid on fluid. A trail of rotationally moving liquid, a Helmholtz "vortex sheet" of exceedingly small thickness, is thus left in the wake of the circular edge; which, while becoming thicker as it gets farther from the edge, becomes rolled up in a wildly tumultuous manner, giving the appearance of an irregular crowd of detached circular ring-vortices. This crowd follows the disc at an ever diminishing speed and widens outward farther and farther, and inwards encroaches more on the comparatively undisturbed middle of the wake, as it is left farther and farther behind the disc.

§ 9. Whether as in § 7 for an ideal inviscid incompressible fluid, or as in § 8 for a natural liquid such as water, the "wake," that is to say, the fluid on the rear side of the plane of the disc, as far as it is sensibly affected by the motion of the disc, must be as described in the last sentence of § 8. The rear of the wake is always moving forwards, that is to say, following the disc; but at a continually diminishing speed. Hence, if the disc has been set in motion from rest some finite time, t , ago, the whole wake must be included between the plane from which the disc started, and the plane in which the disc is now, at the time when we are thinking of it. These two planes are at the finite distance, Vt , asunder. In other words the wake extends to some distance less than Vt , rearwards from the disc.

§ 10. The shedding off of vortex rings from the edge of the disc, to follow in its wake at less speed than its own, essentially gives a contribution to negative pressure on the rear side of the disc equal to $\frac{d}{dt} \Sigma \kappa$; where $\Sigma \kappa$ denotes the sum of the circulations of all the coreless ring vortices, or of all the rotationally moving liquid, which have or has left the edge since the beginning of the motion. This, with the commonly assumed velocities of the fluid on the two sides of the rigid plane, seems insufficient to account for the excess of observed pressure above that calculated for a long blade by Lord Rayleigh's formula³ referred to in my letter to NATURE, "Towards the Efficiency of Sails, &c.," and leaves some correction to be made on those assumed velocities. But the working out of this interesting piece of mathematical hydrokinetics must be deferred for a continuation of the present article in which supposed discontinuity of fluid motion, extending far and wide, as taught by many writers in many scientific papers and text books since Stokes' infinitesimal rift started it in 1847, will be considered. KELVIN.

(To be continued.)

MR. SCOTT ELLIOT'S RUWENZORI EXPEDITION.

ABOUT three years ago, in NATURE (November 5, 1891), I gave an account, rescued from an American periodical, of the botanical results, slender enough it is true, but not without interest, of the Emin Relief Ex-

¹ Viscosity is resistance to change of shape in proportion to the speed of the change.

² Stokes, "Mathematical and Physical Papers," vol. i. p. 310.

³ In lines 9 and 10 of the printed letter (NATURE, Aug. 20, 1894, p. 425), for "something like five or ten," substitute 4.8. I unfortunately had not Lord Rayleigh's formula by me at the time the letter was written.

¹ Continued from p. 525.

² The whole subject of the motion of an incompressible inviscid fluid with vacuum on the other side of the whole or any part of its boundary is of surpassing interest. Consider for example an open fixed basin, with water poured into it and left not quite at rest under the influence of gravity; swinging slightly from side to side; let us suppose for example; the water perfectly inviscid, and vapour-less, but may be either cohesionless or cohesionless; and there being perfect vacuum over all its free surface. Very soon it will certainly throw up a drop somewhere; and before very long it will become covered with spin-drift and will thus illustrate Maxwell's important allegation (which I believe to be true though it has been much doubted), that any conservative system must "sooner or later" pass through every possible configuration.