characters: it will be capable of forming an oxide RO_3 ; its hydride, if it exists, will be even more unstable than tellurium hydride; its compounds will be easily reduced; it will form definite alloys with metals; and have the specific gravity of about 9.3. Since then Prof. W. Preyer, in his "Genetisches System der chemischen Elemente," Berlin, 1893, p. 100, among unknown elements, gives one in the sulphur group, thus predicted by Mendeléef. He goes into further details, representing the atomic weight as 213; specific gravity, 8.6; atomic volume, 24.7; specific heat, 0.03; electro-negative, divalent, diamagnetic.

A. Grünwald (Bl. 1891, 5, p. 21) appears to have seen the spectrum of this element, whilst observing spectra of tellurium, copper and antimony.

Norwegium is considered by some chemists to be this missing element; Dr. Preyer is not of that opinion, regarding the claims of norwegium to be considered an element to be insufficiently supported.

In consideration of Dr. Brauner's work on the atomic weight of tellurium, and the priority of his suggestion of the existence of an accompanying element of higher atomic weight, coupled with his excellent work generally in furthering the claims of the Periodic Law (notably in the cases of beryllium and of the cerium, lanthanum and didymium series), I would suggest that this new element, when isolated from its close union with tellurium, be called Bohemium, in his honour.

C. T. BLANSHARD.

Bright Meteors.

A PRETTY bright meteor crossed the eastern sky here on Saturday evening last, at about 7.54 p.m. I saw only the flash of light which it cast on the ground and in the sky towards the east, like a momentary weak red flash of lightning. "A shooting-star," said a bystander close to me, who saw it fairly well, and who gave me, roughly, this description, by the stars, of its apparent course: From about R.A. 340° , Decl. $+25^\circ$, to about 336° , $+12^\circ$. It described this course of $12''$ in about a second, and was red in colour, and broke up at last with a red flash, leaving no train of light or of sparks along the track which it had traversed, so long as for a second or two in which I had time to look towards the direction where he pointed. Tree-tops intercepted his view beyond the point of this disruption, but the light's sudden extinction there made a much further extension of the track unlikely.

About an hour afterwards, at about 8.49 p.m., I chanced to see another rather fine shooting-star, of about the brightness and colour of the planet Mars at present, descending some 16° or 18° across the northern sky in one and a-half or two seconds, from R.A. 47° , Decl. $+70^\circ$, to 100° $+69^\circ$, beginning and disappearing rather suddenly, and of nearly uniformly bright appearance all along its course. It projected no sparks, and, like the earlier meteor, left no train of light or enduring sparks along its track to mark its course.

Prolonged backwards the apparent paths of these two meteors diverge from near α Cassiopeiæ, and it may perhaps be that a shower of bright, ruddy meteors from the direction of Cassiopeia was in progress on that date, of which the two meteors here described may have been bright enough members to have been generally noted. The above observations, although those of the brighter meteor are only of rather slender accuracy, will perhaps be useful, in that case, to serve for comparison with other records which, at least of the larger meteor of the two, may not be impossible to have been noted and preserved elsewhere.

A. S. HERSCHEL.

Observatory House, Slough, October 1.

Tan-Spots over Dogs' Eyes.

CAN any of your readers explain the meaning of the tan-spots seen so commonly over the eyes in black-and-tan dogs of most breeds?

When in Melbourne last year, I went carefully over all the dogs in a show, with one of the stewards, and we found the spots in all the black-and-tan terriers, foxhounds, deerhounds, collies, lurchers, &c.; but I could get no information regarding them from the experts.

In some of the highly-bred toy dogs, as the small black-and-tan terriers, I found on inquiry that these spots, formerly so

very conspicuous, were being bred out, and had nearly disappeared. Their persistence through so many strongly-marked varieties, except those of late date, is singular, for there is fairly good proof that when first domesticated the dog was red or bright brown, like the pariah, dingo, &c.

As far as I can see, we do not find the spots white on a black or dark ground; nor yet black or dark on a white or light ground. My explanation is that they have arisen as a permanent marking after the dogs "sporting" to black under domestication, and have been preserved and developed through natural selection. Possibly they are protective, and simulate eyes.

One morning, just at dawn, I had occasion to go out into the garden, and while stooping to examine some flowers, near a fence partly covered with creepers, I suddenly saw an animal's head looking through, and what seemed to be two black, and seemingly large, eyes glared at me. Suspecting that a black leopard was about to spring over, I started back, clapped my hands, and shouted. To my relief, however, I saw a tail wag, and found that the spectator was a coolie's dog I knew very well, and which recognised me. The use of the tan-spots—in this case at least—then occurred to me.

May it not be that the spots thus serve a protective purpose, and have often saved the lives of dogs (black dogs) from their enemies, the smaller felines, such as the clouded leopard, &c.? Perhaps the matter is not new; but if it is, it seems worth looking into.

I have several dogs about here now with black bodies and heads; the tan-spots, rather pale, are of the size of a shilling. I have shot one, keeping the skin of the head as a curiosity.

Sibsagar, Asam, September 7.

S. E. PEAL.

Flight of the Albatross.

AT the request of several friends, I enclose, for your inspection, a snap-shot of a northern albatross, which I took *en route* from Victoria, British Columbia, to San Francisco.

The photograph gives the bird in a position in which the human eye is incapable of seeing it. Strangely enough, the flight of the bird which I photographed had been during the whole course of the morning the subject of much discussion: none of us could imagine how the force which enabled it to fly at such

great speed was generated. I remembered having seen some discussion in the papers on the subject, and stated that the matter had been settled to the satisfaction of scientific people, and did my best to explain the theory.

I chose the moment for my photograph when the bird was about fifteen feet from the camera, and sailing alongside of the steamer. The stretch of the wings was estimated by the