A striking contrast is afforded by placing a piece of plain white paper over a portion of the screen, and observing the whole by the light either of a Geissler tube or of the Tesla brush.

Under these conditions the white paper will appear very dark, while the fluorescent surface is brilliantly luminous.

A. A. C. SWINTON.

66 Victoria Street, London, S.W., November 3.

IN continuation of our letter of October 29, we find that a reflexion grating does not show the bands so well as a quartz prism, because metals, e.g. silver-on-glass, do not reflect all this kind of light completely except at grazing incidence. The furthest band shown by a Rowland grating at incidence 45° has wave-length 2200 tenth-metres, whereas the extreme ultra-violet usually quoted (the Fraunhofer line U) is 2948 tenth-metres, and the bright bands shown by quinine are about 3250 and 3830 respectively. With a quartz prism the fluorescence caused by light of shorter wave-length than 2200 can be seen. But it is quite possible that Sir George Stokes in 1852 saw as far as we can see to-day.

OLIVER J. LODGE.

BENJAMIN DAVIES.

November 7.

Osmotic Pressure.

As Mr. Whetham has called attention in your columns to my attempt to give a mechanical hypothesis for osmotic pressure, in the October number of the *Philosophical Magazine*, perhaps you will allow me to give some explanation of my somewhat faulty use of the term "dissociation hypothesis," which, as Prof. Ramsay has pointed out to me, may easily be misunderstood. I have used the term to signify not the separation of the ions in electrolytes, but rather the freedom of the solute molecules in non-electrolytes, and of the atoms in electrolytes—their dissociation, in fact, from the molecules of solvent. It appears to me that in some statements of the facts of osmotic pressure, the idea is strongly suggested that the molecules or atoms of the solute are moving about among the molecules of the solvent, and, as far as pressure at any rate is concerned, dissociated from them, and producing an independent effect, the osmotic pressure being directly due to the solute. My aim is to show that we may more reasonably account for the facts by supposing that the solute molecules or atoms are associated with the solvent molecules, entering into some kind of more or less unstable combination with them, and that the solution is not to be regarded as consisting of two parts producing independent pressures. The extra osmotic pressure is, of course, due to the solute in one sense, in that it would not exist without it; but it is an indirect effect, due to the modified compound molecules formed. The first effect is a decrease in "mobility" of the solution, so that the exchange in the two directions through a semi-permeable membrane is unequal, and it is only rendered equal when the solution is put under the extra pressure which we call osmotic pressure.

Mr. Whetham has shown, in a very simple way, that my hypothesis does not necessarily conflict with the facts of electrolysis, and that the idea of dissociation of the ions from each other may easily be reconciled with it.

J. H. POYNTING.

Mason College, Birmingham, October 31.

"Purple Patches."

I SHOULD be very glad if I could obtain information as to the cause and nature of certain "purple patches" which I have noticed from time to time for many years past, but have been unable to get explained. The patches in question occur during, or immediately after, rain, on the pavement or roadway; dashes of vivid purple, or rather violet, varying in size from small splashes or drops to patches as large as the palm of one's hand, but most commonly they are about the size of a shilling. When quite fresh, sometimes a little clot is observable in the centre of the splash. Sometimes I find one patch completely isolated, sometimes two or three in close proximity; sometimes, again, numerous little drops scattered over a certain space; once I counted twenty or thirty tiny dashes in about ten yards of pavement. When quite wet the violet colour can be rubbed up with a handkerchief or paper, which it stains as with "aniline purple" dye, as it does the pavement, and when once dry it is quite ineradicable, and lasts till it is worn away by exposure, or the feet of passers-by. I observe it to occur chiefly during warm rain after a dry or cold spell; never during dry weather, whether in summer or winter. During the past hot summer there was none to be

found, but directly the weather changed in July, I saw it in various localities. This was also the case in the long cold winter of 1895, when on the breaking up of the frost there were plentiful patches to be seen up and down the streets; there was also a complete absence during the following summer, till the drought gave, and then again I found this appearance recur. I naturally observe it most in Bath, where I live; but it is not at all confined to one place or situation. I have found good specimens at such widely different places as the doorway of a hotel at Oban; the Castle Hill, Edinburgh; railway platform at Morecambe; doorstep at Windermere; in streets and roads at Cambridge, Bude, Penzance, St. Ives, Clevedon; once in a London street (Pall Mall East), and once some was found in a cold water bath.

I have from time to time made inquiries from various people who I thought would know, but have not been fortunate enough to meet any scientific person who has observed it. But one learned professor to whom I described the "patches," suggested whether "purple bacteria" would prove a solution to the mystery, and recommended me to inquire through the medium of your columns. I should be much obliged if some one would enlighten me, or mention some authority to whom I could refer.

A. PEDDER.

13 Somerset Place, Bath, October 27.

Note on "*Plasmodiophora brassicæ*."

ALTHOUGH it is well known that *Plasmodiophora brassicæ* attacks the great majority of cruciferous plants, yet no instance of the common Shepherd's purse (*Capsella bursa-pastoris*, D.C.) being attacked is recorded in this country. Thus Massee (*Proc. Roy. Soc.*, vol. lvii.) quotes the Shepherd's purse as being reported by Halsted to be attacked in America, but says "It has not been observed to be diseased in this country, although one of our commonest weeds." During the past summer my attention was drawn to some plants of *Capsella* with swollen roots, growing in a sandy field near Coventry, on land upon which crops of swedes and turnips were grown in the usual rotation, and I had no difficulty in finding several additional specimens. These roots, on examination, were found to have the characteristic plasmodium in their cells. The Shepherd's purse must now be numbered among the plants in this country which provide a home for *Plasmodiophora*, and probably help it to maintain its existence in the ground from year to year, thus proving a possible source of injury to cruciferous crops.

The length of time for which *Plasmodiophora* can retain its vitality in the soil in the absence of any cruciferous plants, is still a matter of uncertainty. In order to ascertain this, in November 1893, I established a series of experiments (fully described in my annual report to the Newcastle Farmers' Club, 1895), intended to extend over a period of six years. In these experiments, six beds (A-F) and six large 18-inch flower-pots (A-F) were prepared, and each strongly infected with pieces of turnip badly diseased with "Finger-and-Toe," one bed and one flower-pot being sown with turnips each successive spring. The beds and pots acted as duplicate experiments, and the soil in both remained unmanured, and was carefully guarded from the intrusion of *Plasmodiophora*, while all cruciferous plants were rigidly excluded.

In 1894, in both pot A and bed A "Finger-and-Toe" appeared; in 1895, in the pot B "Finger-and-Toe" was found upon four plants out of six; and in the bed B, 8 per cent. were diseased. In 1896 the same result was strikingly shown; after a period of three years, the bed C and pot C were still found to be diseased, four out of five plants being affected in the latter, and 10 per cent. in the former. Massee had previously shown that the germs of disease retain their vitality for two years, and my experiment this year shows that this period can be increased to at least three years.

M. C. POTTER.

Durham College of Science, Newcastle-upon-Tyne, October 24.

Sparrows and Wheat.

IN vol. vii. part iii. p. 522 of the *Journal* of the Royal Agricultural Society it is stated that, in the Leicester district, Rivett's wheat is much grown, the reason being that sparrows do not attack it, while they do other varieties. Can any of your readers assign a cause for the exemption from attack of this particular variety? It is most curious if correct; and the authority quoted apparently is a good one.

F. G. BROOK-FOX.

Port Navis Cove, Penryn, Cornwall, November 1.