Brain in Relation to Mind. By J. Sanderson Christison, M.D. Pp. 142. (Chicago: Dr. Christison, 1899.) "COMPARATIVELY few physicians and fewer laymen," says Dr. Christison, "have a satisfactory view of the relationship of brain to mind," and we are afraid that Dr. Christison is not among the minority. When the writer of a book that purports to be a serious work on psychology commits himself to such statements as that "the use of means to a given end implies the pre-existence of a specific potentiality having a plan in the abstract, for only the pre-existing can be the cause of a necessity"; and that "the evolution idea . . . is absolutely incompatible with law and order," it would be out of place to deal with him in a serious review.

LETTERS TO THE EDITOR.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Is New Zealand a Zoological Region?

As some weeks must elapse before Mr. Farquhar is able to reply to the questions asked (p. 273) by my good friend Mr. Wallace, I would urge your readers to bear in mind that no particular sanctity should be attached to the word "Region." Such a sanctity was once ascribed to the word "Species," and to no one living more than Mr. Wallace are we indebted for the removal of the misconception therein involved. Properly used both are very good words, and it would not be easy to get on without them; but to each may be ascribed a virtue which it does not rightly possess. In the case of New Zealand I would meanwhile refer to the expression of Prof. Huxley's views in the *Proceedings of the Zoological Society for 1868*, which so far as I know have never been refuted. ALFRED NEWTON.

Cambridge, January 20.

Compensation in Weather.

As our climate does not, in a long series of years, undergo permanent change, we are accustomed, in a time of extreme cold, wetness, or other quality of weather, to expect, in a vague way, that we shall have, ere long, a variation in the opposite direction, equalising matters. Is it possible to attain greater definiteness in such expectation, and, utilising the principle of compensation, to forecast, in a measure, the character of an approaching season?

I propose to inquire what sort of relation subsists between the cold of a given winter and that of the thirty winters preceding.

We may conveniently measure the cold of winter seasons by the total number of frost days from September to May. (For brevity, we shall here designate each winter season by the year in which it ends; 1842, e.g. meaning 1841-42.)

Let us, then, begin by adding the numbers of frost days in the thirty winters 1842 to 1871, 1843 to 1872, and so on. We thus get a series of twenty-nine numbers, varying from 1593 to 1717, with an average of 1653.

The following table may now be considered:—

Groups of 30 winters containing, of frost days	Average number of frost days in winter following.
(1) Under 1610 ... 3 cases	68.0
(2) 1610-1629 ... 6 "	66.2
(3) 1630-1649 ... 5 "	60.4
(4) 1650-1669 ... 6 "	44.0
(5) 1670-1689 ... 2 "	51.0 e
(6) 1690 or more ... 6 "	42.0

Thus, in general, the larger the number of frost days in the 30-winter groups, the smaller, on an average, is the number in the winter following. A reversal occurs at No. 5. (But there are only two cases.)

This is, perhaps, very much what we should expect on the principle of compensation.

Analysing further, we must bear in mind that the data considered are not very numerous, and beware of building too much on a slender foundation. The remarks here offered are rather by way of suggesting a method which might perhaps be

found practically helpful, if fuller data were found to point in the same direction. We may, then, note these three facts:—

(1) The six mildest winters (since 1871) were each preceded by a 30-year group having more than the average of frost days.

(2) The six coldest winters were each preceded by a 30-year group having less than the average of frost days.

(3) Of fifteen 30-year groups with excessive cold (i.e. over the average), as many as twelve were followed by mild winters, and only three (i.e. one-fifth) by severe winters.

Applying the same method to frost days in the first half of each year, it might be shown that out of twelve 30-year groups with excessive cold, ten were followed by mild first halves, and only two by severe first halves.

Reverting to the former case (total winter seasons), the diagram represents, in curve A, the total frost days in 30-winter groups ending 1871, 1872, &c. Below is an inverted curve, the first point of which indicates the number of frost days in winter 1872, the second in winter 1873, and so on. (The time scale below, it will be seen, is shifted one year to the left.)

A. Curve of frost-days in 30 winters ending 1871, 1872, &c.
B. Curve of frost-days in winters 1872, 1873, &c. (inverted).

A certain correspondence may be traced between these curves. And taking the dots above the average line in A, it will be seen, how, except in three cases, marked by a small circle (in B), each corresponds with a mild winter in B, a fact already noted.

There is a curious lag in part of the second curve. We find two conspicuous wave-crests in A, viz. in 1881 and 1893, and the highest points (i.e. lowest values) in B are in 1884 and 1896. Further, the lowest point in A is in 1885; the lowest in B in 1888. An interval of three years in each case. It might be worth while to observe whether this is repeated to any extent in the future course of these curves.

Now the last thirty winter seasons (ending 1899) have a total of 1657 frost days, and if we allowed ourselves to speculate as to the quantity of frost in the current winter season (1900) we might at least perhaps fairly anticipate, or might have anticipated, that this would not turn out an extremely severe winter, and that the chances are rather against its being extremely mild either. But the point I would emphasise is that, with copious material, useful hints might possibly be obtained as to the limits within which the character of a coming season (or other portion of time), as to heat or other quality would probably occur. To this end, I may remark, some might find a diagram of dots helpful, in which, taking the case of winters, the abscissæ and ordinates are used for the 30-winter groups, and the following winters respectively, a dot being placed at each crossing point of the lines from the two scales. ALEX. B. MACDOWALL.