CORRIGENDUM

High prevalence of diabetes, obesity and dyslipidemia in urban slum population of northern India

A Misra *et al*

International Journal of Obesity (2002) 26, 1281. doi:10.1038/sj.ijo.0802030

Correction to: International Journal of Obesity (2001) 25, 1722–1729

Following the publication of the above paper, the author has identified errors in Table 2. The correct version of Table 2 is reproduced below.

Table 2 Profile of various measures of obesity

Measures of obesity	Males		Females	
	n	Mean $\pm$ s.d.	n	Mean $\pm$ s.d.
Body mass index	165	20.0 $\pm$ 4.1	359	20.7 $\pm$ 4.5 (NS)
Skinfolds (mm) and ratios				
Waist–hip ratio	170	0.86 $\pm$ 0.11	362	1.02 $\pm$ 0.39 (NS)
Biceps	170	5.5 $\pm$ 3.8	362	8.3 $\pm$ 5.4*
Triceps	170	9.7 $\pm$ 6.6	362	14.8 $\pm$ 7.8*
Subscapular	170	13.1 $\pm$ 7.3	362	17.1 $\pm$ 9.5*
Suprailiac	169	14.3 $\pm$ 7.5	361	19.9 $\pm$ 11.3*
Σ 4SF (sum of four skinfolds)	169	42.3 $\pm$ 21.2	361	60.0 $\pm$ 30.9*
Central skinfolds (sum of subscapular and suprailiac skinfolds)	169	27.2 $\pm$ 13.0	361	36.9 $\pm$ 19.8*
Peripheral skinfolds (sum of triceps and biceps skinfolds)	170	15.2 $\pm$ 9.6	362	23.1 $\pm$ 12.6*
Central–peripheral skinfold ratio	169	2.01 $\pm$ 0.69	361	1.67 $\pm$ 0.59*
SS/TR (Subscapular–triceps skinfold ratio)	170	1.54 $\pm$ 0.64	362	1.23 $\pm$ 0.7*
Body fat (%)	169	17.5 $\pm$ 6.2	361	28.4 $\pm$ 7.6*

NS, statistically not significant; * $P < 0.001$.