CORRIGENDUM

Weight loss and long-term follow-up of severely obese individuals treated with an intense behavioral program

JW Anderson, L Grant, L Gotthelf and LTP Stifler

International Journal of Obesity (2007) 31, 565. doi:10.1038/sj.ijo.0803541

Correction to: *International Journal of Obesity* (2007) 31, 488–493. doi:10.1038/sj.ijo.0803423

next to 26-week weight loss. The correct table is shown below:

Since the publication of the above paper the authors have noticed an error in Table 1. The asterisk should appear

The authors would like to apologise for the errors.

Table 1 Baseline data, weight loss and follow-up weights for different groups. Follow-up data represents the last clinic weight available

	Study group			Completers			100# group		
	Women	Men	All	Women	Men	All	Women	Men	All
<i>Group</i>									
Number	998	533	1531	689	411	1100	121	147	268
Age (years)	43	45	44	44	46	44	43	44	43
s.d.	11.3	11.2	11.3	11.5	11.0	11.3	10.9	10.1	10.5
Initial weight (kg)	128.2	158.6	138.8	129.0	157.9	139.8	139.9	173.7	158.4
s.d.	21.4	29.7	28.5	21.1	29.3	28.2	25.8	33.5	34.6
Initial BMI (kg/m ²)	47.6	49.5	48.3	48.0	49.4	48.5	51.3	53.9	52.7
s.d.	6.9	8.6	7.6	7.0	8.8	7.7	8.8	10.3	9.7
<i>Weight loss phase</i>									
12-week weight loss (kg)	14.5*	22.5*	17.3*	18.2	27.0	21.2	23.9	33.9	29.3
s.e.	0.3	0.5	0.3	0.3	0.6	0.3	0.4	0.7	0.5
26-week weight loss (kg)*	19.4	29.1	22.8	25.8	37.0	29.9	41.7	52.5	47.6
s.e.	0.4	0.8	0.4	0.5	0.8	0.5	0.7	1.1	0.7
Total weight loss (kg)	23.9	36.0	28.1	30.8	42.6	35.2	58.2	65.7	62.3
s.e.	0.6	1.0	0.5	0.6	1.1	0.6	1.2	1.5	1.0
Total weight loss of initial weight (%)	18.5	22.5	19.9	23.9	26.7	24.9	41.5	37.5	39.8
s.e.	0.00	0.01	0.00	0.00	0.01	0.01	0.01	0.01	0.01
Treatment, weeks	28.8	32.6	30.1	38.1	39.0	38.5	64.5	50.8	57.0
s.e.	1.4	1.9	1.1	1.8	2.3	1.4	5.8	4.2	3.5
<i>Follow-up</i>									
Number available	697	406	1103	589	358	947	117	134	251
% available	69.8	76.2	72.0	85.5	87.1	86.1	96.7	91.2	93.7
Weight maintained (kg)	17.5	25.9	20.6	19.4	28.1	22.7	36.6	44.8	41.0
s.e.	0.6	1.1	0.6	0.7	1.2	0.6	1.8	2.1	1.4
% Weight maintained (%)	59.7	63.1	61.0	57.0	62.2	59.0	62.5	67.5	65.1
s.e.	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02
Duration follow-up, weeks	59.3	72.0	64.0	67.9	79.0	72.1	92.3	96.8	94.7
s.e.	3.2	5.0	2.7	3.6	5.5	3.1	8.8	9.9	6.7

Abbreviations: BMI, body mass index; s.d., standard deviation; s.e., standard error. *Last observation carried forward.