is merely attempting to teach chemical arithmetic with no regard for the opportunities which arise of teaching chemistry at the same time. The figures actually resulting from the best experiments are so readily accessible that a book which fails to make use of them and substitutes obvious fictions is scarcely to be recommended.

(5) This book of tables is well compiled, and should prove useful, but the printing and binding are not as good as might be desired in view of the small size of the book and the price at which it is issued.

MINERAL SPRINGS AND WELLS OF ESSEX.

A History of the Mineral Waters and Medicinal Springs of the County of Essex. By Miller Christy and Miss May Thresh, with a critical note by W. H. Dalton. Essex Field Club Special Memoirs, vol. iv. Pp. vi+73. (Stratford, Essex: Essex Field Club; London: Simpkin, Marshall and Co., Ltd., 1910.) Price 2s. 6d. net.

THIS work forms vol. iv. of the Essex Field Club Special Memoirs, and has been reprinted from the Essex Naturalist, with additions. As in most other English counties, there are in Essex a number of springs and wells that have attained notoriety as mineral or medicinal waters, and the authors have done well to prepare a full and precise account of them.

The earliest record is of a spring at Wanstead, which was regarded as a spa in 1619, but has long been lost to sight. Witham, Chigwell Row, and Upminster had mineral waters that were formerly reputed to be of medicinal value. No information is available concerning the particular constituents of the Witham Spa; the water of Chigwell Row was purgative, but of no importance; while that of Upminster contained Epsom salts. Tilbury water, obtained from a well, appears to have been most famous in Essex. but, as the authors remark, the saline ingredients were insufficient to justify its being considered a mineral water. Dr. Richard Russel, however, remarked in 1769 that the water "makes excellent Punch, and is extremely good for Tea." The only genuine mineral waters acknowledged in the present work are those of South Weald, Upminster, and Hockley, which contain as the more prominent ingredient magnesium As the authors admit, every so-called mineral spring in Essex, with one exception, is now neglected, and almost forgotten; and as regards the waters in general they consider "that 'faith' was an important, if not the chief, element in the 'cures' they are credited with." Dovercourt Chalybeate Spa, discovered about 1852, is the sole remaining spa, and in a sample of the water sent in 1897 to Dr. J. C. Thresh, he reported that it contained under one grain of iron salts per gallon.

The authors express their indebtedness to Dr. Thresh for assistance in dealing with the Essex waters from a chemical point of view, and to Mr. W. H. Dalton for notes on the strata whence the waters are derived. Reference should have been made to the Bagshot Sands on p. 63, as the waters of Hockley, as well as those of South Weald, are derived from that

formation or the passage-beds above the mass of London Clay. There are no deep-seated mineral waters in Essex, but the subject, as shown by the authors, is one of considerable interest, and by no means devoid of scientific importance.

H. B. W.

OUR BOOK SHELF.

(1) Edible and Poisonous Fungi. Board of Agriculture and Fisheries. Pp. 28. With 25 coloured plates. (London: His Majesty's Stationery Office, 1910.) Price 16.

(2) Guide to Mr. Worthington Smith's Drawings of Field and Cultivated Mushrooms and Poisonous or Worthless Fungi often Mistaken for Mushrooms, Exhibited in the Department of Botany, British Museum (Natural History). Pp. 24. (London: Printed by Order of the Trustees of the British Museum, Natural History, 1910.) Price 1s.

(1) THE publication of this pamphlet by the Board of Agriculture and Fisheries is intended doubtless to broaden the "mushroom" diet of country dwellers. Whether this object will be attained depends primarily on the doubtful possibility of creating an interest in a lethargic public, and further, in making quite clear the somewhat abstruse differences between the clean and the unclean. With regard to means of discrimination, reliance is placed on coloured plates and short descriptions, to which are added a few hints on preparation for table. What is distinctly lacking is an attractive general account, with information regarding the kinds exposed for sale in foreign market places, where there is often a considerable variety. The list of edible species does not include either the chantarelle or the truffle, while another notable omission is a warning that individuals vary greatly in their power of digesting fungal ferments.

(2) The pamphlet issuing from the British Museum (Natural History) is valuable both as a scientific exposition by one of our most eminent fungologists and also as an authoritative guide for the use of those interested in mushroom cultivation. The descriptions are semi-popular, and the coloured figures are artistic, accurate and well rendered. The setting of the text, as also the plain directions for detecting the poisonous species, add to the practical utility of the pamphlet, which fulfils one of the chief objects of the Trustees, inasmuch as it provides accurate and useful information for the benefit of the general

public.

Fractures and Separated Epiphyses. By A. J. Walton. Pp. vii+288. (London: E. Arnold, 1910.)

In a short preface the author explains that this book is intended for the use of students and those first commencing hospital appointments, but there is every reason to believe it will prove of great value to practitioners in general. Mr. Walton does not confine himself to advising any one method of treatment, but concisely places before the reader the various treatments advocated, with an open-minded criticism of their several points.

The chapters dealing with the etiology and general methods of treatment are, considering the largeness of the subject, both clearly and shortly dealt with, yet nothing of importance has been omitted. In describing the fractures peculiar to each bone, with their treatment, special attention is given to the dates of union in the various epiphyses, and the injuries which they are liable to sustain. The accompanying illustrations, reproduced from radiographs of fractures seen at the London Hospital, are typical and excellent. The book show great care in preparation, and can be recommended to all who need a short, practical work on this subject.

Frank Romer.