to prophesy in individual cases. This, indeed, is at the root of a great deal of the criticism levelled at the claims of education to rank as a science. A parent or an education authority presents a boy to an educator and calls for a prophecy. The educator must decline, since he cannot honestly prophesy in an individual case, though he may be prepared to venture on a reasoned statement of what is likely to occur in the boy's educational career. The educator is, in fact, in precisely the same position as a medical man called in to a case. He can prophesy, but only in general terms. In both cases it is the application of general principles to a particular case.

This raises the whole question of the value of the average in matters of education. Psychologists, in addressing teachers, are beginning to warn them that the average is only an abstraction, and really does not exist. We are told that what the teacher has to concern himself with is "the living child here and now before him," and he is accordingly warned against the insubstantiality of the elusive abstract. But this is to confound two distinct things. It is true that the teacher must always deal with a living pupil here and now before him. But in his dealing with that living pupil he has to apply a paid-up capital of knowledge of men and of boys in general. He must seek to understand the living boy by the aid of knowledge previously acquired, and this knowledge is represented by the average. The master may be unable to prophesy with certainty how Jones minor will act under certain specified conditions. But from a knowledge of third form boys in general he can make a guess that is very likely to hit the mark. The teacher who applies his knowledge of the average third form boy to the minor Jones, without modification to suit Jones's case, acts unintelligently; but the possibility of blunders by a dull master does not reduce the value of the knowledge of the average in the hands of one who is capable. The concept of the average boy as it is developed by experience and study in the mind of the master forms a standard by which other boys may be estimated. This standard is partly subjective, partly objective. In so far as the standard is acquired by the personal experience of the master it is subjective. The unreasoned but very effective knowledge of boy nature that enables an efficient master who is guiltless of any acquaintance with educational theory to know how a boy is likely to act in given circumstances results from the training of experience, and is peculiar to its possessor. On the other hand, the knowledge of boy nature that has been acquired by deliberate study and by experiment is something that has an existence independent of the individual. It is objective, or at any rate has an objective bias.

We must distinguish in practice between the average and the type. The average boy may have no existence in reality, he may be a pure abstraction; but the type is concrete, and may be regarded as the embodiment of all the essentials that go to make up the average, with the addition of certain qualities that must be present in some form or other, though the particular form is immaterial. The average is to the type as the concept is to the generalised image. The type may form a very useful standard for masters whose tendency is strongly towards the concrete; but the average has a special and a different value, and in capable hands is more effectively applied because it is of a wider range. To consider a class as made up of types tends to break up the class feeling, and make the master think of his pupils as a mere group of separate individuals. Undoubtedly the master must in certain connections think of his pupils as individuals, but in other con-

nections he must deal with his class as a whole, as a psychological unit.

This introduces one of the most striking developments of modern educational theory. The older psychologists treated their subject as limited to the study of the mature human individual. The introduction of the idea of development led to the founding of a genetic psychology with its consideration of the individual at his various stages. A further advance is marked by the appearance of collective psychology, which carries the study of the individual into his relations with other individuals. Naturally, both changes were of the greatest advantage to education. The first gave scientific guidance to the popular movement known as Child-Study, the second suggested the scientific study of the class as a collective organism. It is true that this collective psychology is at present in its infancy. But while we owe much to the French psychologists with their dazzling exposition, we are glad to turn to our more solid McDougall for the best scientific basis available for a sound collective psychology. The material he has supplied is waiting to be worked up from the educational side. His statement of the relation between the instincts and the emotions and his manipulation of Mr. Shand's theory of the sentiments provide tempting material for the establishment of an objective standard in connection with the training of the individual character and the interaction of individual characters in groups. Naturally, the results must be expressed in averages, and equally naturally there will be a complaint from certain practical educators. What is the use, it will be asked, of information about how classes in general act? What we want to know is how this particular class before which I stand is going to act. But this is to confound the practice of a science with the science itself. There must always be an intelligent intermediary between the principles of a science and their application to the affairs of life. In this respect the nascent science of education differs in no way from those that are more fully developed. The educator who prides himself on being specially practical is frequently very unreasonable in his demands from educational theory. He is rather apt to complain that it does not supply him with sufficiently detailed instructions. What he wants is a series of recipes which, if scrupulously followed, will inevitably produce certain specified results. But such men take a very humiliating view of their profession. So far from seeking this spoon-feeding, they should rejoice that their work demands the exercise of intelligent initiative. Herein consists, in fact, the dignity of the educator's office. He must be master of the organised knowledge that education has acquired, and must have the power of making the appropriate application of that knowledge to every case as it arises. To assist him in avoiding error he is entitled to look for an objective standard at the hands of those who make education their special study, but for the use of that standard he must himself accept the full responsibility.

GEOLOGY AT THE BRITISH ASSOCIATION.

THE proceedings of the Geological Section at Dundee were of exceptional interest, and the attendances were large up to the end of the meeting. The success of the section was due to two or three special features. Many of the papers dealt with the problems of the Highlands and of the Highland border, questions which are full of knotty points, and the men who are engaged in solving these problems were able to assemble in the meeting-room and there

state their views and face discussion, which was often very lively and always full of vigorous earnestness. Then we had a commodious and quiet meeting-room, and the President (Dr. B. N. Peach) was a host in himself. As Dr. Heim said, in seconding a vote of thanks to him: "When I look in front I always see the sun," and the vivacious countenance and stirring enthusiasm of Dr. Peach were an incentive to all present to rise to their best.

After the delivery of the President's address on "The Relation between the Cambrian Faunas of Scotland and North America," which was a timely and most valuable deliverance giving the evidence for the land connection between North America and North-west Europe in Cambrian times, a lecture was delivered by Dr. T. J. Jehu on the local geology. After a brief reference to the exposures of the Highland Boundary series in Forfar and Kincardine, and to the recent discoveries of rocks of Downtonian age near Stonehaven, the lecturer dealt with the groups of rocks found in the area:—(1) The Highland metamorphic rocks; (2) the Old Red Sandstone; (3) the Carboniferous system of Fife.

(1) The prominent series of metamorphic rocks comprises the Dunkeld slates, schistose slates and micaceous grits included in the Ben Ledi group. With the schists are associated the "Green Beds" of the Geological Survey and certain basic intrusions mapped as hornblende schists.

(2) Two divisions of the Old Red Sandstone are represented, Lower and Upper, separated by a marked unconformity. They consist of two sedimentary groups separated by the great volcanic series of the Sidlaws and the Ochils. The distribution and characteristics of the volcanic beds were described, and the probable positions of the vents from which the materials were ejected were indicated. The Lower Old Red Sandstone rocks were folded and denuded before the deposition of the Upper Beds, which pass conformably up into the Carboniferous System in Fife.

(3) The Carboniferous rocks of Fife consist of the Calciferous Sandstones at the base, overlaid by the Carboniferous Limestone series, which are overlaid by Millstone Grit and Coal Measures. Special attention was directed to the splendid development of workable coal-seams in the middle group of the Carboniferous Limestone series, as well as in the Lower Coal Measures of the district. The remarkable diversity of igneous rocks and the great development of volcanic rocks on the coast of Fife were described, and were subsequently displayed to the members by a field-excursion to Elie.

During the meeting various problems connected with the Pre-Cambrian rocks of the Highlands and the Highland border were thoroughly expounded and discussed.

Mr. E. B. Bailey described the occurrence in eastern Mull of a great breccia-formation with intercalated rhyolitic lavas. The breccia consists of an unbedded assemblage of blunted rocks and fragments of gneiss, granophyre, gabbro and basalt, often associated with rhyolitic *débris* of volcanic origin. The basalt lavas of Mull have been violently folded into a series of anticlines and synclines, and it is in one of these synclines that the main outcrop of the breccia is preserved. It appears that the breccia is a thick layer overlying the basalts and folded with them. Though some parts of the breccia may be of volcanic origin, Mr. Bailey suggests that the greater part has resulted from erosion, which operated during the period of upheaval of the ridges.

Dr. J. S. Flett opened a discussion on the sequence of volcanic rocks in Scotland in relation to the Atlantic-Pacific classification of Suess. He pointed out that the recognition of two great families of igneous rocks, the Atlantic and the Pacific, and their relation to

certain types of earth-movement, which we owe to Mr. Harker, constituted one of the greatest advances in rational petrology. In Scotland the Carboniferous volcanic rocks are typical Atlantic types, associated with collapse and great faults, and the rocks of Lower Old Sandstone age are characteristic of the Pacific group (great crumbling). To the Atlantic group could be added the Permian or late-Carboniferous rocks of Ayrshire and East Fife, and the nepheline-basalts (presumably Tertiary) of Caithness with their associated camptonites and monchequites. The Tertiary volcanic rocks of the Hebrides and the abundant north-west dykes were also ascribed to movements of Atlantic type. The remaining rocks of Scotland Dr. Flett desired to assign to an independent, intermediate group. They are characterised by pillow-lavas and are not connected with movements either of the Pacific or the Atlantic kind, and may thus be placed in a special family.

Dr. Flett's address was followed by a vigorous discussion, in which considerable criticism was expressed of the terms Atlantic and Pacific, and especially of their application to types of rocks in Scotland; but Dr. Flett pointed out that some terms must be used, and these were convenient, but he was not pledged to them in any way. Dr. J. W. Evans (London) thought that Dr. Flett had made out a *prima facie* case, so far as the Scotch volcanic rocks were concerned. He believed that the first differentiation of igneous rocks was a result of the separation of the original magma, on cooling, into two fluids, one consisting mainly of water, silica, alumina and alkalies, the other made up chiefly of magnesia and ferrous oxide with sufficient alumina and silica to form anorthite and the ferric minerals. Subsequent differentiation was principally due to crystallisation. In folded areas the segregation took place at a considerable depth, while the magma still retained its original aqueous contents. The primary differentiation was therefore comparatively complete, and the normal or "Pacific" series of rocks was ultimately formed. In districts of plateau faulting, on the other hand, the differentiation occurred nearer the surface and under conditions which facilitated the loss of water, so that the separation was imperfect, the basic portion retaining more silica and a considerable amount of alkalies, and subsequent segregation by crystallisation gave rise to the alkali or "Atlantic" series. He also suggested that the occurrence of sodium with the basic rocks might be due to the presence of an unusual amount of chlorine, which separated out with the basic constituents and brought the sodium with it.

Dr. Tempest Anderson did not agree with the suggestion that pillow-lavas had been formed in deep water, and gave instances which he had seen of their formation by lavas flowing into shallow waters, and apparently due to sudden cooling by the waves. Mr. G. W. Tyrrell (Glasgow) much preferred the terms "alkalic" and "calcic" for Dr. Flett's main divisions. The differences were mainly chemical and mineralogical, not geographical. Many group names cover a wide diversity of types, and we appreciate the great value of Dr. Flett's recognition of mono- and poly-phyletic rocks, that is, types highly characteristic of one group only and types to be found in all the main divisions of igneous rocks. Dr. Hatch pointed out the advantage of a division which recognises a well-defined suite of rocks in which the alkali and calc-alkali feldspars are developed in equal proportions, a series to which Brögger has given the name Monzonite series, and hoped that such a classification would be generally adopted.

Mr. G. Barrow gave a valuable paper on the older granite in Lower Dee Side, in which he showed that in place of forming large coherent masses, it was

distinguished by *lit-par-lit* intrusions. The granite material in these cases formed minute sills parallel to the foliation of the associated gneiss into which it had been intruded. The granite rises as a dyke and the sills diminish in thickness and extent as they near the surface. In the interior parts the granite is usually grey and contains more biotite than muscovite, and oligoclase is usually abundant. The oligoclase and biotite are found to diminish steadily as the rock is traced towards the taper end of the sills. At this point there is little oligoclase and often no biotite; muscovite is fairly common and the bulk of the felspar is of alkaline composition. It would appear that the fissures in which the dykes occur were filled with igneous material, and that under great pressure the walls were burst open and the still liquid material forced out, and thus separated from that which had already become segregated. This phenomenon may be described as magmatic differentiation intensified by dynamic action. Mr. Barrow also read a paper on buckled folding. Descriptions have been published of areas of regional crystalline metamorphism in which the dip of the bedding is described as at a low angle over a considerable area. Experience is gradually proving that these altered sediments are always intensely folded, and the low dips really represent the most complicated structure, for which the name "Buckled Folding" is suggested. This structure is best seen in the Moine Gneisses, and its development can be studied in the cliff sections between Stonehaven and Muchalls. The grits and shales on the limb of the folds, near Stonehaven, ascend the cliffs in an unbroken course from bottom to top, being isoclinal and unbent. But as we proceed northwards the course of the grit bands up the cliff face is no longer straight, but a small overfold, or "buckle," is developed in it. At first only one is seen in the whole height of the cliff, then two, then three, and so on, until they are so close together that the still straight portion of the fold is no longer than the "buckled" or overfolded portion. If the upward course of each grit band be followed, it will be found that this structure does not alter the dip of the band as a whole. It still descends at much the same angle, but by a zigzag course. The overfolds all face in the same direction right up to Muchalls, and must have been produced after the isoclinal folding was completed. There is no justification for separating the buckled beds from those in which buckling does not occur. A key to the connection is found in the area about Shiehallion, where the quartzose beds forming the margin of the quartzite and containing the boulder bed and the limestone show isoclinal folding; whereas further north the same group comes on again, but this time with buckled folding, or Moine gneiss.

Dr. B. N. Peach and Dr. J. Horne described some interesting investigations which they had recently made on the Archæan rocks of Lewis, in which they showed how closely these beds corresponded to the rocks of the adjoining mainland, without the great series of acid intrusions. The structure is coarsely granulitic, and there is a marked absence of the pyroxene-gneisses with blue quartz, of pyroxene-granulites, and other basic forms, which are so characteristic of the mainland. The remarkable series of basic dykes in the west of Sutherland had not been detected in Lewis. The north-west and south-east strike, referred to by Murchison, was not characteristic of the gneisses of Lewis. The dominant strike is almost north and south, but north-east to south-west or east to west in other areas. The flaggy granulitic gneisses of the Butt of Lewis, which appear to run southwards along the belt of high ground between Stornoway and Barvas, resemble closely the Moine gneisses east of the Moine thrust-plane, but they differ petrologically from the

rocks of sedimentary origin which form the Moine series. The system of overfolding and the direction of the axial planes of the folds approximate to those found in the Moine rocks on the mainland. The platy rocks, or mylonites, occur along definite lines of movement approximately north to south, and thrust-planes have been detected which point to displacement in a westerly direction. Various stages in the development of mylonites from the acid and basic gneisses are represented. In the discussion which followed the value of these investigations was emphasised by Dr. Flett and others.

Important discoveries of fossils in old rocks were announced by Dr. R. Campbell (Edinburgh) and Prof. T. J. Jehu (St. Andrew's). Dr. Campbell described fossil remains found in the Jasper and Green Schist series of the Highland border, at Craigeven Bay, Stonehaven. Crushed spilitic lavas with intercalated black shales, jaspers, and cherts, which in their lithological characters resemble closely the green igneous rocks and associated sediments along the line of the Highland fault, appear on the old Geological Survey maps as of (?) Arenig age. In 1909, in company with Dr. Peach and Dr. W. T. Gordon, the author found several fossils in the black shales. Detailed search by Mr. D. Tait revealed fossils of the forms *Lingulella*, *Obolella*, *Acrotreta*, *Linnarssonella*, and *Siphonotreta*; a bivalve phyllocarid allied to *Caryocaris* and *Lingulocaris*; and cases of a tubuloid worm. The above genera are most commonly found in the Ordovician beds and in the Upper Cambrian, but, owing to the absence of graptolites, Dr. Peach suggested that Upper Cambrian was the most probable age. Whatever may be the ultimate decision as to their stratigraphical horizon, the discovery of these fossils leaves very little doubt that the boundary fault series is *not* pre-Cambrian.

Dr. Jehu followed this paper by one on the discovery of fossils in the boundary-fault series, near Aberfoyle. This series is well exposed between Loch Lomond and Callendar, forming a narrow belt separated by a reversed fault from the Lower Old Red Sandstone on the south-east, and probably by a line of thrust from the Leny Grits on the north-west. It consists of black and grey shales, cherts, grits, and calcareous beds. Remains of *Radiolaria* were discovered by Dr. Peach some years ago in cherts near Gualann. Recently a number of fossils have been found in pale-grey chert bands near Arndrum. These fossils occur in muddy films in the chert belt. They are almost all hingeless brachiopods, and the following forms have been determined by Dr. Peach:—*Acrotreta*, *Lingulella*, *Obolus*, *Obolella*; also the flattened *chaetæ* of polychæte worms. These fossils were regarded as indicating an Upper Cambrian age. Dr. Horne, in the discussion, regarded the collection of fossils from the chert and green-schist series at Stonehaven and Aberfoyle as the most important palæontological "find" affecting Highland geology since the discovery of *Olenellus* in the west of Ross-shire. The strata containing these fossils, which had been provisionally referred to the Upper Cambrian by Dr. Peach and Dr. Walcott, could no longer be considered as of pre-Cambrian age, like those of similar rocks in Anglesey. At a subsequent meeting Dr. Horne read a letter from Dr. Ami (Toronto), who had examined Dr. Jehu's collection of fossils from Aberfoyle, and found them closely resembling those obtained from Upper Cambrian beds belonging to the Quebec group. His opinion was based partly on the brachiopods and partly on the occurrence of an obscurely preserved graptolite resembling *Retiolites ensiformis*, Hall—a type eminently characteristic of the Silurian Sandstones of the Quebec group of Upper Cambrian age. Dr. Horne further stated that this form had been shown

to Miss Ellis (Cambridge), who recognised it as a graptolite with Ordovician affinities.

Dr. A. W. Gibb (Aberdeen) read a paper on an actinolite-bearing rock allied to serpentine associated with the basic intrusion of Belhelvie. Towards the northern end of this mass, which consists of troctolites, serpentines, and allied types, there is a rock which shows a large number of dark-green rounded spots, set in a fine felt of paler green colour, full of glancing needles. Under the microscope the spots are seen to represent olivine, partly unaltered, partly serpentinised, as well as granulitised and drawn out. The rest of the rock is largely made up of actinolite in small crystal flakes, and green spinel and abundant magnetite are present. The exposure has recently been blasted away.

Dr. Wm. Mackie described the volcanic rocks round the Ord Hill of Rhynie, Aberdeenshire. The group embraces at least three lava flows, with associated tuffs and interbedded and overlying sedimentary rocks. Flow brecciation is frequent, and the tuffs are rhyolitic. The sedimentary rocks of the group consist of hard siliceous grits, which, under the microscope, show volcanic fragments. The whole of the group is characterised by fine, secondary quartz infiltration veins. They overlie the basic rocks of the younger Grampian granite series, but are probably considerably older than the oldest Old Red beds of the adjoining area. The lavas, being of an acid type, cannot be correlated with the interbedded andesites of the Old Red Sandstone.

Mr. G. W. Tyrrell (Glasgow) described the alkaline igneous rocks of Ayrshire. Recent work of the Scottish petrologists shows that one of the greatest developments of rocks characterised by primary analcite is contained in the midland valley of Scotland, one centre being in Ayrshire, the other in the area surrounding the Firth of Forth. Geologically these analcite rocks are Carboniferous in age, having a time-range from Carboniferous Limestone to Early Permian. They occur in the form of stratiform sills, lenticular intrusive masses, volcanic plugs, and as a series of lava flows. No masses of true plutonic habit are known.

Passing to Wales, a paper was read by Mr. Edward Greenly on the origin of some of the mica-schists of Anglesey. The mica-schists of southern and central Anglesey are holocrystalline rocks with strong parallel structure, and composed essentially of quartz, alkali-feldspars, and white mica. Dr. Teall regards them as broken-down and partially reconstructed porphyritic felsites. Twenty-five years ago Dr. Callaway recognised their felsitic origin, which the present investigations confirm. These schists, therefore, may be looked upon in general as derived from acid igneous rocks. In some areas mica schists in continuity with them are found in intimate relations to schists of sedimentary origin, so that, probably, pyroclastic material was present in the original igneous series.

Dr. Robert Campbell (Edinburgh) gave a valuable *résumé* of his recent investigations on the Lower Old Red beds of Kincardineshire, showing that a thickness of 3000 ft. ought now to be transferred to the Downtonian, the uppermost group of the Silurian system. At the base of the series near Stonehaven there is 200 ft. of breccias interbedded with fine red mudstones made up mainly of fragments of the underlying (?) Upper Cambrian rocks, and resting unconformably on them. Near Cowie Harbour there occurs a thick belt of grey and greenish mudstones which yield *Dictyocaris* in abundance. *Ceratiocaris*, *Archidesmus*, *Eurypterus*, and other forms have been obtained from the same horizon, and in another bed numerous plates of a new *Cyathaspis* have been found. These interesting fossil finds, which in other areas do not

occur in rocks younger than the Upper Silurian, and the lithological characters recall the typical Downtonian of the south of Scotland. These beds pass upwards into the micaceous sandstone and conglomerates of Stonehaven Harbour, which may be considered as the base of the Lower Old Red Sandstone. The fish remains were examined by Dr. R. H. Traquair, who reported that the small scutes were referable to the category of Cephalaspidian scutes, that the species to which they belong is pretty certain, but additional material is required before proceeding further with identification. He described several specimens of a beautiful new *Cyathaspis*, which he dedicated to Dr. Campbell.

Mr. C. J. Gardiner gave an account of the Silurian inlier of Usk. It is an oval area, eight miles by four, crossed by an important east and west fault. The southern half is composed of two anticlines separated by a fault. The axes of these folds run north and south, and dip southwards. The western anticline is the larger of the two, and shows Wenlock Shales and Limestone and Ludlow beds. The northern half of the inlier is far less simple than the southern in its structure, and is more concealed by drift. The Wenlock Limestone is not met with in this half. The simplest explanation is that the Wenlock Shale is faulted against the Ludlow beds on both sides. As the Aymestry Limestone is absent from the area, it is impossible to separate the Ludlow beds into an upper and lower division. The main boundary faults are crossed at several spots by minor east and west faults which cause small lateral displacements.

There were three important contributions from the palaeobotanical side. Dr. Marie C. Stopes gave an outline of a controversy in New Brunswick between the palaeobotanists and the stratigraphers. In the "Fern Ledges" of St. John there is a rich fossil flora, but no animal remains. Sir W. Dawson described the plants as Devonian. Recent attempts have been made to include the beds in the Silurian for stratigraphic reasons. The author's work in the field indicated considerable overthrust. The unique series had recently been redetermined. Type specimens had been lent by the Canadian museums and brought to London and Paris for comparison, resulting in the identification of a large proportion of well-known European types in the "Fern Ledges" flora as Carboniferous and mostly typical of the Westphalian division of the Coal Measures.

Dr. W. T. Gordon read a paper on the fossil flora of the Pettycur Limestone in relation to botanical evolution. The flora of the Pettycur Limestone (Lower Carboniferous) has a double interest. These forms constitute fragments of the oldest known flora, as Lower Carboniferous plants do not differ markedly from the Upper Devonian forms. Although the Devonian flora is distinct from that of the Permian-Carboniferous epoch, yet the organisation does not indicate that the plants were primitive. The result of Dr. Gordon's investigations was that the flora represented in the Pettycur Limestone appears, on the whole, to contain more generalised and simpler types than occur in the Coal Measures and later strata, and these types may be arranged in order so as to suggest certain possible lines of evolution.

Mr. W. R. Don (Dundee) read a paper on the nature of *Parka decipiens*. This fossil is the only common and quite the most characteristic fossil of the Lower Old Red Sandstone of the Kincardine-Forfar-Perth area. A re-investigation, chiefly microscopical, has been attempted with the aid of Schultz's solution (strong nitric acid and potassium chlorate). Most previous investigators had pronounced it to be vegetable, though Mantell, Lyell, and others considered it as an

egg-packet of *Pterygotus*. The conclusions of the author were confirmatory of those of Sir W. Dawson and Prof. Penhallow that it was vegetable, and after boiling in nitric acid, the presence of spores within the carbonised tissue was demonstrated. In attempting to form some conception of the original structure and shape of Parka, the author concluded that the original spore-containing tissue was almost flat, not spherical, and unlike any known sporangia of to-day. There was, certainly, intimately connected with it a so-called "indusium." In the discussion, Dr. G. Hickling stated that his independent observations on Parka were in very close agreement with those of Mr. Don. He questioned, however, the nature of the cells, hitherto regarded as spores, and preferred to consider them as constituting simply a parenchymatous mass. These masses possibly served as reproductive buds analogous to the gemmæ of the *Hepaticæ*. Dr. Newell Arber (Cambridge) remarked on the extreme interest of Mr. Don's careful work on Parka, and was quite prepared to allow that the organs termed spores were undoubtedly of that nature. He differed from Dr. Hickling's criticisms, and pointed out that if these bodies were not spores, but parenchymatous cells, they certainly could not have survived the extremely severe chemical treatment to which they had been subjected. Dr. Hickling appeared to have examined chiefly spore sacs in which the spores were not yet mature, a condition also commonly noticed in some of Mr. Don's specimens, and this appeared to have been the chief basis of his criticism. Dr. Arber agreed with Mr. Don that Parka was undoubtedly a member of the *Thallophytæ*, and very possibly an alga.

The first results of investigations on the contents of the Millstone Grit of Yorkshire were communicated by Mr. Albert Gilligan (Leeds). Following up the work of Dr. Sorby, the late Mr. A. Longbottom had collected some very large pebbles from the Middle Grits of Silsden. These had been examined by the author, who had extended his researches to other beds of the series in Yorkshire. Some of the pebbles were of considerable size and showed a remarkable assemblage of rocks. By far the most common were acid igneous rocks. Only one specimen of a basic igneous rock had been found. The metamorphic rocks were quartz-schist and mica-schist, with a few fragments of gneiss. Numerous pebbles were found to be perfectly fresh microcline. Pieces of pegmatite were common. Some fragments obtained from the Plompton Grit proved to be a peculiar silicified oolitic rock. A few pebbles showed traces of organisms such as sponge spicules. The heavy minerals of the grits were not numerous, zircon and garnet being the most plentiful. The feldspars in the grit were quite fresh, and this suggests either disintegration of the parent rock by differences of temperature and rapid transportation, or comparative absence of carbonic dioxide in the atmosphere. The author had been much impressed by the many points of similarity between the Millstone Grit and the Torridon Sandstone, and was disposed to think that areas of similar rock types were laid under contribution for each.

Mr. T. O. Bosworth gave an account of some investigations into the heavy mineral grains in the sands of the Scottish Carboniferous. The chief heavy minerals found were garnet, zircon, magnetite, tourmaline, rutile, staurolite, anatase, barytes. The sands containing an extraordinary amount of angular garnet were probably derived from the Highland schists of the north and north-west. The sands devoid of garnet probably came from the north-east, east, or south.

Mr. J. S. Owens communicated the results of some experiments on the settlement and transport of sand in water. It was shown that there is a definite rela-

tion between the rate of settlement and the temperature of the water. The curves show that velocity of fall varies almost with the water temperature, the rate being always increased by rise of temperature, but that as the diameter of the grains increases the temperature effect becomes less, until for grains more than one-tenth inch in diameter the effect is practically negligible.

Mr. Edward Greenly contributed a theory of the Menai Strait, in which he accepted Ramsay's view of the Strait as a glacial furrow, but the middle of the Strait cannot be explained in that way. Evidence was adduced to show that this reach was excavated by glacial waters during the recession of the ice at a time when the mutual relations of the ice of the mountain land and of the sea-basin admitted of the accumulation of a temporary lake. Post-glacial erosion and subsequent changes of level have completed the bed of the Strait as it now exists.

The origin of kopjes and inselberge was dealt with by Dr. J. D. Falconer. It had been suggested that a landscape with inselberge was of desert origin, but the various phenomena could be explained more readily as the result of weathering and erosion during successive small oscillatory movements of a regional character in the neighbourhood of base-level.

Mr. G. W. Grabham gave notes of an exploration of the country north of Lake Albert. This area extends west from Rejaf to the watershed and southwards as far as the lake. It is entirely composed of gneisses, which are for the most part affected by a north-south foliation. The only member of apparently sedimentary origin consists of a band of quartzite which is traceable for some distance. Among the gneisses, a group characterised by graphitic pegmatite was recognised. Some intrusions occur among these gneisses, and form the only intervening link in time between them and the surface deposits. In the country north of Lake Albert extensive deposits belonging to an extinct lake were found. Lake Ismail, as it is proposed to call this lake, stood about 600 ft. above the present river-level, and its site is marked by gravels and beds of clay. Its limits are uncertain, but it did not extend into the area now occupied by Lake Albert. In later times the crust fractures of the Rift Valley occurred, and the present lake and river system was initiated. In more recent times still the river has again been modified by crustal movements.

The usual grants were recommended for the continuance of committees of research, and a new committee was formed to investigate the fish beds of Dura Glen. The meeting was attended by a large number of foreign geologists, who took an active interest in the discussions of the section. Among them were Prof. C. Barrois (Lille), Dr. Tschernyschew (St. Petersburg), Prof. A. Heim (Zurich), Dr. H. Reusch (Christiania), Prof. J. Welsh (Poitiers), Prof. E. Tietze (Vienna), Dr. Pirsson (Yale), Dr. Leith (Wisconsin), Dr. Ami (Toronto). The annual geological dinner was held at the Royal Hotel on Friday evening (September 6). Dr. Peach made a most genial chairman, and delighted the audience by rendering the "Song of the Seraphim," specially written for the Red Lion Club dinner held during the Dundee meeting in 1867, and then sung by the author, Dr. Henry Woodward, to the tune of "Bonnie Dundee." The menu card was graced by a photographic copy of the picture of Sir Charles Lyell painted in 1870 by Lowes Dickenson, when Sir Charles was seventy-three years of age. The picture was on view in the loan collection. Dr. Jehu and his assistants arranged three enjoyable excursions for the afternoons, and a whole day to see the wonderful coast sections of Carboniferous py eruptions near Elie. At the close of the meeting two extended field excursions were arranged, one

to Loch Assynt, under the leadership of Dr. Peach and Dr. Horne, and another to the country between Aberdeen and Arbroath, under the leadership of Mr. Barrow, Dr. Campbell, and Dr. Hickling. These were very enjoyable and instructive, and proved great attractions to the foreign geologists and a large number of their British *compères*.

W. LOWER CARTER.

UNIVERSITY AND EDUCATIONAL INTELLIGENCE.

CAMBRIDGE.—The John Winbolt prize has been awarded to R. V. Southwell, of Trinity College, for an essay on "The Failure of Thin Tubes by Instability or Secondary Flexure."

BIRMINGHAM.—During the winter and spring terms two courses of lectures on "Civic Design and Town Planning" are to be given, by Mr. Raymond Unwin, in the department of civil engineering.

The following appointments have been made:—Mr. C. Walker as lecturer in physiology; Mr. Laurence Ball as assistant lecturer in pathology and bacteriology; Mr. P. M. Chadwick, assistant lecturer and demonstrator in civil engineering; Mr. A. Clubb, demonstrator in mining (as successor to Mr. C. D. Mottram); Mr. H. I. Coe, assistant lecturer and demonstrator in metallurgy. Mr. Percy May has resigned his post as assistant lecturer and demonstrator in chemistry, and Mr. Frederick Challenger has been nominated to the vacancy.

A COURSE of free lectures to teachers on "The Past Around Us," a series of brief studies introductory to the folk-culture of Britain, is being given by Mr. Walter W. Skeat, at the Horniman Museum, Forest Hill, S.E., on Saturday mornings, from October 12 to December 14. Admission is by ticket only, to be obtained from the Clerk of the London County Council.

LECTURES on volcanic action, earth movements, the geological action of water, and the evolution of scenery and life on the globe are to be delivered by Dr. Werner Marchand on October 17, 24, and 31, in the meeting rooms of the British Esperanto Association, 133 High Holborn (Museum Station Buildings), W.C. They will commence at 7.30 p.m., and will be delivered in Esperanto.

THE winter meetings of the Child Study Society begin this evening at the Royal Sanitary Institute, when Dr. T. P. Nunn will lecture on the psychological development of the school subjects. The list of lectures and discussions to be held this year provides many subjects of interest to students and teachers concerned with the education of children. Particulars as to membership may be obtained from the honorary secretary of the society, Mr. W. J. Durrie Mulford, 90 Buckingham Palace Road, London, S.W.

THE University College (London) Committee will shortly proceed to fill the vacancy in the Quain studentship in biology which has been created by the resignation of Mr. E. J. Salisbury, on his appointment as lecturer in botany at the East London College. Any student of the college is eligible for the studentship who has for at least three terms attended one or more classes in the special study in respect of which the studentship is awarded. Applications should be received on or before Saturday, October 26.

FOR some time articles have been appearing at regular intervals in the Journal of the Department of Agriculture and Technical Instruction for Ireland describing recently established Irish technical schools. These articles have been published afterwards as

separate pamphlets for distribution by the department. The twelfth and thirteenth contributions to the series have been received in booklet form. The former is called "Technical Instruction in Limerick," and has been written by Mr. J. Comerston, the principal of the Limerick Technical Institute; the other deals similarly with Cork, and is by Dr. John H. Grindley, principal of the Crawford Municipal Technical Institute, Cork. The accounts of the work done in technical education in these important Irish industrial centres provide excellent evidence of the success which is attending the department's efforts to meet the educational needs in different parts of Ireland.

THE distinguishing characteristic of the calendar for the present session of the City of Bradford Technical College is the excellent series of thirty plates, which chiefly illustrate the very complete arrangements made for the practical study of the branches of technology on which the industries of the district depend. This college awards certificates, diplomas, and an associateship. The diploma of the college is awarded to each day student who has been in attendance for three complete sessions, subsequent to passing an entrance examination, and has passed the college examinations in all subjects of the diploma course taken. The diploma is awarded to evening students under the same regulations as to day students, except that an evening student who has been at least three years in attendance, and has obtained the ordinary certificate, is exempt from the first-year diploma course. To become an associate a candidate must be twenty-one years of age and have had at least one year's practical experience with a firm engaged in his trade or profession, subsequent to obtaining the diploma. Some of the subjects in which diplomas may be obtained are: preparing, combing and spinning, weaving and cloth structure, chemistry and dyeing, and power production and transmission.

ON Wednesday, October 9, at Bradfield College, Berks, a new block of science rooms was opened by Sir William Osler, F.R.S. A large proportion of the boys at the college have studied science during the last twenty years, and some fifty to seventy pupils work in the mechanical shops added in 1898. The new science schools have this year been added to deal more effectively with the growing demand, and mainly through the efforts of the present headmaster, the Rev. H. Costley White. Among the assembly present at the ceremony were the warden (Mr. Edward Armstrong), the Right Hon. G. W. Palmer, Mr. R. Dyke Acland, K.C., Sir Arthur Rücker, F.R.S., and Mr. J. H. Benyon. Sir William Osler, in a speech after the ceremony, dwelt on what he considered to be an ideal education for those suited to and seeking scientific pursuits in after life. He would have a thorough knowledge of Latin and Greek; he believed in the optimistic Greek outlook on life for boys; during his last two years the boy should specialise in science, which should occupy most of his school hours. The speaker objected very strongly to the use of the term "stinks" as applied to science study. He said that that one word had done more harm in implying discredit to, and in keeping back pupils from the study of, the subject than any other factor. The new science block is detached from the rest of the school buildings, and has an attractive exterior. The entrance lobby leads into two chemical laboratories on the right and two physics rooms on the left, each easily accommodating twenty boys. In the chemical laboratories each room is adequately fitted with fume cupboards, balance slabs and store cupboards, and has a raised demonstration bench at one end of the room. The two physics rooms are each sup-