CORRIGENDUM

The impact of one or two missed doses on the duration of action of combined perindopril and indapamide

MG Myers¹ and FHH Leenen², on behalf of the General Practice Research Group

¹Schulich Heart Centre, Sunnybrook Health Sciences Centre, Department of Medicine, University of Toronto, Toronto, Ontario, Canada and ²Hypertension Unit, University of Ottawa Heart Institute, Ottawa, Ontario, Canada

Journal of Human Hypertension (2008) 22, 516; doi:10.1038/sj.jhh.1002307

Correction to: *Journal of Human Hypertension* (2007) 21, 86–93. doi:10.1038/sj.jhh.1002107

Following the publication of the above paper, the authors have identified an error in Figure 2 and would like to apologize for any confusion

this might have caused. The correct figure is reproduced below. The numerical values for the fall in blood pressure 0–24 h after dosing and after one (24–48 h) and two missed doses (48–72 h) were correctly stated in the text of the original article.

Figure 2 Changes in mean $\pm$ s.d. 24-h ambulatory systolic and diastolic blood pressure versus baseline during the first 24 h after dosing with perindopril 4 mg + indapamide 1.25 mg (P 4 + I 1.25) and after one (24–48 h) and two missed doses (48–72 h). Mean values for 0–24 h after dosing are shown separately for the one and two missed-dose sub-groups. All changes in mean blood pressure versus baseline were significant ($P < 0.001$).