ductions from excellent photographs of timber operations and forest scenes in many parts of the British Empire. The inclusion in the work (pp. 328-84) of the tables of strengths of woods, which were published by Laslett, is probably justified by the fact that Laslett's book is out of print. These tables are of considerable value, but as Laslett had not at his disposal apparatus for determining the moisture contents of the woods examined, the figures are not so trustworthy as they seem. It would certainly be preferable in future, as Mr. Howard admits, to conduct experiments on woods "with some named and specific percentage of moisture." Moreover, we are now aware, thanks to the numerous tests carried out of late years at the Forest Products Laboratory, Madison (U.S.A.), that there is a remarkable variation in the properties of timbers of the same species from different logs and from different localities. That timbers are in no way comparable with metals in the uniformity of their physical characteristics is of course well known.

A chapter on the artificial seasoning of timber, with a note on laboratory and working tests, contributed by Mr. S. Fitzgerald, will prove useful.

There are errata and misstatements in this book, but they are not of a kind to detract much from its real value, which consists in the thoroughly practical nature of the information given on the uses, conversion, and utilisation of so many kinds of timber. Some of the inaccuracies may, however, be pointed out. It is, unfortunately, not true that, as stated on p. 118, "larch disease has practically disappeared." The explanation of the name "sycamore" on p. 266 is entirely erroneous, and is an instance of the strange fascination that wrong etymology has for many people. The word "sycamore" was originally applied in Greece to a species of *Ficus*, and is now transferred in England to the large-leaf maple, *Acer pseudoplatanus*, and in America to the Western plane, *Platanus occidentalis*. A whole page—p. 164—is devoted to the etymology of the term "wainscot oak," about which there is no doubt. It could have been expressed in three lines. "Wainscot" is derived from a Dutch word (fourteenth century) which means "oak wood with a wavy grain"; in other words, "wainscot oak" is "figured oak." Mr. Howard misquotes Skeat, who ("Concise Etymological Dictionary," p. 597) expressly warns the reader against the wrong derivation attributed to him by Mr. Howard. A regrettable error occurs on p. 211, where *Abies pectinata*, the European silver fir, is

called "silver spruce." As is pointed out on pp. 262-63, the latter name can be applied only to the Sitka spruce (*Picea sitchensis*). Endless

Fine example of African mahogany curl. (From "A Manual of the Timbers of the World.")

embarrassment to the landowning class resulted during the war from the confusion between the names "silver spruce" and "silver fir."

The Structure of the Atom.¹

By C. G. DARWIN.

II.—Atomic Number (continued.)

IT would involve very prolonged work to get the atomic numbers accurately by direct experiment with scattered α -particles. Fortunately, this is not necessary, as there exist much more con-

venient indirect means of determining them. The invention of a powerful method of studying X-ray spectra enabled Moseley to examine the spectra from a sequence of elements. Whereas visual spectra have a highly complicated structure, and exhibit the same periodicity as do chemical pro-

¹ Continued from p. 54.

erties, for the X-rays this periodicity disappears, and is replaced by a perfectly orderly progression from element to element. Each spectrum can be derived from the last by a reduction of the wavelengths according to a simple rule. Thus an examination of a sequence of spectra reveals the true order of the elements, and gaps are shown by a missing term in the sequence. It will be seen that Moseley's method only gives differences of atomic number, and not the numbers themselves. To determine these we must find one of them in some other way. This is not difficult, for we know from observation on α -particles that helium must have atomic number 2; and though X-rays have not been observed for the

sion of an α - or a β -particle from the nucleus, and this emission transmutes the atom into a new element. From a study of the chemical behaviour of these successive elements it was concluded that the emission of an α -ray shifts the element two places to the left in the periodic table, while the emission of a β -ray shifts it one place to the right. This fits in exactly with the conception of atomic number, for when the nucleus loses an α -particle (that is, a helium nucleus of atomic number 2) its own number must be reduced by two, whereas the loss of a β -particle out of the nucleus must raise the number by one, since the β -particle is an electron and has a negative charge. By a study of all the radio-elements we can work out

FIG. 1.—The Radio-active Transformations. In every case a step two downwards is accompanied by the emission of an α -particle, and one upwards by a β -particle. Actinium is derived from uranium or an isotope, but the connection shown is rather conjectural. Thorium C has a third branch about which little is known. The final products in all cases are believed to be lead, but as they are not radio-active it is uncertain how many of the isotopes are really different. In the Table of Atomic Numbers they were counted as all different, and one was added on to allow for the existence of common lead. One was also added there for 81 and 83 to represent the non-radio-active types of thallium and bismuth.

first dozen elements, the periodic table is perfectly regular in that region, and it is impossible to believe that it could contain any gaps. More remarkable still, if the formula for the wavelength of the X-ray spectrum is extrapolated right back to hydrogen, it reproduces the fundamental line of the famous Balmer series. The numbers which were given in the table are therefore certainly correct.

In formulating the principle of atomic number we said that it, and not the weight, completely determines the chemical character of the element. This statement is justified by a line of evidence which we must now consider. The process of radio-active transformation consists in the emis-

sion of an α - or a β -particle from the nucleus, and this emission transmutes the atom into a new element. From a study of the chemical behaviour of these successive elements it was concluded that the emission of an α -ray shifts the element two places to the left in the periodic table, while the emission of a β -ray shifts it one place to the right. This fits in exactly with the conception of atomic number, for when the nucleus loses an α -particle (that is, a helium nucleus of atomic number 2) its own number must be reduced by two, whereas the loss of a β -particle out of the nucleus must raise the number by one, since the β -particle is an electron and has a negative charge. By a study of all the radio-elements we can work out

The existence of isotopes was first proved from radio-active evidence. But besides possessing the radio-active characteristics of the atom, the nucleus also carries its mass, and this suggests that isotopes may differ in mass. The first evidence for this can be derived from the study of the figure. For example, uranium at 92 loses an α -particle and two β -particles, and thus returns to 92; and the loss of the particles must have reduced its weight from 238 to 234. The same principle has been experimentally verified by finding that the atomic weight of lead from radium is perceptibly different from that of ordinary lead. We shall see in the next article that ordinary elements also have isotopes; for convenience, these have been given in the table, in so far as they have been yet discovered. It should perhaps be here mentioned

that the mere difference of atomic weight theoretically implies certain minute differences in chemical behaviour, but it can be shown that these are out of all proportion less than those between substances of different atomic number, so much so that it is an almost impossibly laborious task to separate isotopes by ordinary chemical or physical processes.

The general outcome of our argument has therefore been that atomic number, and not atomic weight, is the determining factor in the behaviour of an element. There are exceptions to this rule—for example, specific heat depends directly on the weight of the atom—but apart from these exceptions the importance which in the past the chemist and spectroscopist have given to atomic weight must in the future be attached to atomic number.

(To be continued.)

Obituary.

PROF. WILHELM WUNDT.

WITH the death on September 1, at the advanced age of eighty-eight years, of Prof. W. Wundt a remarkable and striking personality passes away from the scientific world. If, as a philosophic thinker, he did not possess either the speculative genius or the insight of a Herbart or a Lotze, he was yet a mind of extraordinary versatility, whose comprehensive acquaintance with vast fields of knowledge has rarely, if ever, been rivalled. His amazing activity as a writer has been for long a source of wonder to his contemporaries; year after year books, pamphlets, and articles have issued from his pen in steady succession, and there was no department of philosophy which he thus left untouched. Naturally, this tremendous literary output is not all of equal worth, but almost everything he wrote exhibits a surprising mastery of detail and power of turning it to account in reaching theoretical conclusions. As a teacher, too, his influence has been extremely wide and far-reaching; students from all parts of the world met in his class-room, and worked in the Institute of Experimental Psychology at Leipzig, the foundation of which was due to him.

Wilhelm Wundt was born on August 16, 1832, at Neckarau, near Mannheim. In 1851 he began the study of medicine at Heidelberg, and in subsequent semesters pursued his medical studies further in Tübingen and Berlin. Fr. Arnold and E. Hasse, of both of whom he always spoke with great respect, were his teachers in Heidelberg; while in Berlin Johannes Müller was then at the height of his fame, and in the great man's laboratory Wundt was for some time busily engaged. On the completion of his career as an undergraduate, he turned his attention first of all to pathological anatomy, and took his degree in Heidelberg in 1886, his *Arbeit* being a thesis on the condition of the nerves in inflamed and de-

generated organs. In the following year he habilitated in the Faculty of Physiology of the same university, and remained in Heidelberg for some years as Helmholtz's assistant in the physiological laboratory. During that period he published two monographs on physiological subjects—the “*Beiträge zur Lehre von den Muskelbewegungen*” (1858) and the “*Beiträge zur Theorie der Sinneswahrnehmung*” (1859-62)—in the long Introduction to the latter of which he first outlined his conception of the scope of empirical psychology as a natural science, and insisted upon the necessity of using experimental methods in the observation of mental processes. It was in consequence of his being occupied with the problems of sense-perception that the young physiologist was induced to enter the domain of metaphysical inquiry, and started to read, “*ziemlich ziel- und planlos*,” as he tells us, the works of Kant, Herbart, and Leibniz. The first fruits of these and of his more strictly psychological labours saw the light in the “*Vorlesungen über Menschen- und Thierseele*” of 1863—a volume in which, as he afterwards expressed it, he set about his task with more zeal than discretion, and which he came to look upon as containing the wild oats of his youthful days. There followed in 1865 a text-book of “*Human Physiology*” (which apparently had a large circulation, a fourth edition appearing in 1878), and in 1867, when he was already a professor in Heidelberg, a voluminous treatise on what he called “*Medizinische Physik*,” intended to acquaint medical students with the exact physical methods needful in medical investigation. Probably the most elaborate piece of experimental research undertaken by himself was that of which account is given in his “*Untersuchungen zur Mechanik der Nerven und Nervencentren*,” the first part of which was published in 1871 and the second in 1876. The bold and ingenious hypothesis, which he here