that the content of the frame is the content of the fabric in absolute rest, or to make any other deduction he logically can. He does not pretend to explain what energy is or what it may become if reduced to absolute rest. He does not assert that there is no absolute rest, but that it escapes his and all experience.

JOHN G. McHARDY,
Commr. R.N. (Retired).

16 Ebury Street, S.W.1, May 6.

The Production of Metallic Zinc.

IN the issue of NATURE for April 28 I observe under "Notes" (p. 279) a reference to the small volume on zinc recently issued by the Imperial Mineral Resources Bureau. In this reference it is pointed out that the figures relative to the production of metallic zinc in the United Kingdom for 1913 do not harmonise with the figures of production and imports of zinc-ore. Naturally so, for there are other factors involved in the production of metallic zinc in any given year. The output of metallic zinc is not necessarily derived entirely from the ores produced at home or imported in that particular year; the part played by "secondary" production—that is, metal obtained from hard zinc—is of importance in this connection.

As regards the use of the expression "long ton," to which the writer of the note objects, preferring the words "statute ton," it has been made abundantly clear in the prefaces to the Bureau's publications that "the weights are expressed in long tons—that is to say, the British statute ton of 2240 lb." The ton of 2240 lb., though the "statute" ton in the United Kingdom, is not necessarily the "statute" ton in other countries. The expression "long ton" has not only the advantage of conciseness, but it is also well understood throughout the mining and metallurgical world.

R. A. S. REDMAYNE,
Chairman of the Imperial Mineral
Resources Bureau.

2 Queen Anne's Gate Buildings,
Westminster, London, S.W.1.
May 4.

SIR RICHARD REDMAYNE puts forward two explanations to account for the discrepancy in the statistics published by the Imperial Mineral Resources Bureau. The first of these, namely, that stocks of ore may be carried over from year to year, is, in view of the relatively small differences from year to year, inadequate to account for the great discrepancy noted. The second is, in fact, the true explanation. Secondary zinc accounts for about one-half of the so-called zinc output of the country, and thus seriously affects the statistics.

THE WRITER OF THE NOTE.

The Theory of Vision.

PROF. JOLY's papers on vision are very interesting. He adopts the visual purple as the visual substance, but there is no evidence that the rods are percipient elements. The view that they are percipient elements is based on errors, as, for instance, that certain animals—the tortoise is the most quoted—possess only cones; that the periphery of the retina is colour-blind; and that the Purkinje phenomenon is not found with the fovea. The tortoise has the rods and cones as definitely marked and distinct from each other as in man. Has any reader seen a retina in which there

are only rods or only cones in any animal? The periphery of the retina is not colour-blind. Red of sufficient luminosity can be seen to the extreme periphery. The Purkinje phenomenon is found with the fovea, and is a photochemical phenomenon. It is very improbable that the rods are percipient elements. An elaborate nervous mechanism is required to regulate the sensitiveness of the photochemical film, and this appears to be the function of the rods.

The stimulus in vision is undoubtedly liquid, as shown by the movement of positive after-images.

The decomposition of the visual purple stimulates the ends of the cones. The ends of the cones consist of a series of discs varying in diameter.

F. W. EDRIDGE-GREEN.

May 7.

A New British Land Planarian.

MR. MORISON's discovery of the interesting planarian worm (*Rhynchodemus Scharffi*) in a garden at Chiswick, as described in Prof. Dendy's letter in NATURE of May 5 (p. 298), shows that this species has a wider range than was at first anticipated. As Prof. Dendy states, it was first discovered in a Dublin garden in 1894, but since that date it has turned up in the Royal Botanic Gardens at Glasnevin, Dublin. I thought it had probably been introduced into both localities, but that nevertheless it was indigenous to Ireland.

It seemed to me most likely to have been brought from the country with a load of turf. This view was confirmed when, in April, 1901, I found several specimens of this planarian worm in the open country under a fallen tree-trunk near Ballymote, Co. Sligo (see *Irish Naturalist*, vol. x., 1901, p. 133).

R. F. SCHARFF.

National Museum, Dublin, May 12.

Cutting Sections of Cotton Hairs.

IN our laboratory we have now, for some months, utilised Mr. H. J. Denham's plan for celloidin-paraffin embedding of the cotton hairs, on the lines of Kultschitzky's and other processes (Worden: "The Nitro-cellulose Industry," p. 805), described in NATURE of May 5, p. 299, which Mr. Denham kindly communicated to us when he first suggested it, and we have found it most satisfactory. We immerse the hairs in dilute celloidin, which is then boiled down to a syrup (Gilson's process); the hairs are next transferred to paraffin-chloroform, and thence to 60° C. paraffin (Ide's process); this makes a very rapid technique, cut sections being available within two hours. We have also tried the method of Willows and Alexander, but find it cytologically inferior to this celloidin-paraffin technique, which gives us excellent sections at 2.5 μ setting on a Leitz sliding microtome, with accidental sections even thinner.

W. LAWRENCE BALLS.
H. A. HANCOCK.

Experimental Department,
The Fine Cotton Spinners' and
Doublers' Association, Ltd.,
Manchester, May 13.

British Scientific Instruments.

IN the review of the "Dictionary of British Scientific Instruments" published in NATURE of May 12, p. 324, it is stated that the British Optical Instrument Manu-