Letters to the Editor.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Periodicities.

THE recent paper by Sir William Beveridge on "Wheat Prices and Rainfall" (Journal of the Royal Statistical Society, vol. 85, pp. 412-478, 1922) raises a rather important question of principle which is involved not only in discussions over the existence of periodicities, but also over relationships between different variables.

Before Schuster's papers on the periodogram it was customary for a period to be accepted as real provided that it had an amplitude comparable with that of the original figures under analysis; and he revolutionised the treatment of the subject by showing that if the squares of the intensities of the various periodic terms are plotted in a periodogram, and if the data are those of an entirely chance distribution, then the average value of an ordinate being a , the probability that a particular ordinate will equal or exceed ka is e^{-k} . Sir William Beveridge is accordingly perfectly justified in taking Schuster's sunspot period of 11.125 years, or Brückner's 34.8 year period, and deciding that these periods probably occur in his wheat prices if the corresponding intensities are three or four times the average. But he, like many other investigators, goes a stage further, and after picking out the largest from a large number of intensities he applies the same criterion as if no selection had occurred. It is, however, clear that if we have a hundred intensities the average of which, a , is derived from a number of random figures, then the probable value of the largest of these chance intensities will not be a but will be considerably greater, and it is only when the largest amplitude actually derived materially exceeds the theoretical chance value thus obtained that reality can be inferred.

Taking the periodicities of wheat prices on pp. 457-459 between 5 years and 40 years,¹ I estimate that the "width of a line" ranges from 0.1 year for a 5 years' period, through 0.5 at 12 years to 4 years at 33 years; and accordingly that the number of independent periods between 5 years and 40 is in this case about 51. The value of a , the average intensity, being 5.898, it is easily seen that the chance of all the 51 random intensities being less than $3a$ is $(1 - e^{-3})^{51}$, or 0.074, so that the chance of at least one intensity greater than $3a$ is 0.926, not e^{-3} or 0.050, as is habitually assumed. Instead of the chance of an occurrence of $3a$ "making a *prima facie* case for enquiry" (p. 424), the odds are 12 to 1 in favour of its production by mere chance. The chance of at least two intensities above $3a$ is 0.728, of three it is 0.470, of four 0.248, of five 0.109, of six 0.0403, of seven 0.0127, of nine 0.00085, and of eleven 0.00003.

Thus it is not until six intensities over $3a$ are found that the chance of production by pure luck is less than 1 in 20. It is also easily found that if the chance of all the 51 intensities being less than na is to be 19/20, n is 6.9; i.e. the greatest intensity for wheat price fluctuations must be 41, not 18, before the probability of its being due to luck is reduced to 1/20;

¹ Sir William Beveridge points out on pp. 423-424 that amplitudes for periods of less than 5 years are inevitably diminished, while those above 31 are diminished by the process employed for eliminating secular trend: I calculate that the intensity at 35 years should be multiplied by $(0.87)^{-2}$ or 1.3, and that at 34 by 3.8.

and if the likelihood is to be 1/100 we must have $n = 8.5$, the corresponding wheat-price intensity being 50. Of intensities greater than 41 Sir William Beveridge found four, and greater than 50 only two.

At first sight it might seem that the agreement between Sir William Beveridge's forecasted synthesis rainfall curve and the actual rainfall was too great to be explained by a few harmonic terms; but the correlation co-efficient of 0.38 (see p. 475) indicates that while 0.38 of the rainfall variations are accounted for, only $(0.38)^2$, or about a seventh, of the independent factors which control these variations have been ascertained.

As pointed out in a paper "On the Criterion for the Reality of Relationships or Periodicities," in the Indian Meteorological Memoirs (vol. 21, No. 9, 1914), the same principle is valid when discussing relationships. If we are examining the effect of rainfall on temperature and ascertain that the correlation co-efficient between the rainfall and temperature of the same month in a particular English county is four times the probable error, we may infer that the effect is highly probable. But if we work out the co-efficients of that temperature with a hundred factors taken at random, e.g. with the monthly rainfall of Tashkend 5.8 years previously, and pick out the largest co-efficient, it would be wrong to compare it with the average co-efficient produced by mere chance; as shown in the paper referred to, the probable value of the largest of 100 co-efficients is 4.01 times as great as the probable value of one taken at random.

GILBERT T. WALKER.

Meteorological Office, Simla, August 24.

DR. WALKER's note contains, I think, a valid and valuable criticism of the procedure commonly adopted hitherto in comparing individual intensities with the average intensity in harmonic analysis. It would lead me now to modify in several ways my general discussion of the "test of intensity" (pp. 422-424 of my paper in the Journal of the Royal Statistical Society. I was particularly careful, however, in that paper to avoid laying stress on intensity as such. The net result of Dr. Walker's calculations is not to weaken but to confirm my main thesis, that a number of real periodicities exist in European wheat prices from 1550 to 1850.

According to these calculations, the chance of my getting by pure luck between five and forty years one intensity as great as $3a$ is 0.926, but the chance of my getting seven such intensities is 0.0127, and that of getting eleven is 0.00003. Actually I have, between five and forty years, fifteen intensities above $3a$ ($=17.69$); the odds are therefore 80 to 1 that at least nine of these intensities, and 33,000 to 1 that at least five of them, are not due to luck. Obviously every such intensity does, in the circumstances, present a *prima facie* case for further inquiry, the object of the inquiry being to determine which of the 15 intensities have the strongest probabilities of being due to real periods.

In that inquiry the actual height of the intensity in any case (the "test of intensity") is only one and not necessarily the most important point for consideration. As Dr. Walker shows, an intensity in my periodogram of nearly seven times the average might well be due to pure luck (the odds being only 20 to 1 against it). On the other hand, a much lower intensity might represent a true and perfectly regular but weak periodicity, just as a quite small correlation co-efficient may prove a real though weak connexion, if the number of cases compared is very large. Indication of the same period in each half of a sequence when analysed separately (the "test of