On one occasion when developing a photographic plate at the Observatory I received a slight shock from a lightning discharge nearby, or more probably from an induced charge in the lead lining of the developing table. Since then I have hesitated to go up to the Observatory during thunderstorms, and have been accustomed to look out for the "all clear" signal which these mammato-cumulus clouds give us.

Another remarkable fact connected with local thunderstorms at Kodaikanal, and probably elsewhere, is the curious roaring sound emanating from the cloud before a storm begins. At first I considered this was due to heavy rain approaching, but concluded that this could not be so. The situation at Kodaikanal is such that one may find oneself very near to a cloud mass rising over the steep sides of the mountains, and the sound always appears to come from the cloud itself, and not from the ground or from trees. Possibly Dr. Simpson can explain this; he would probably have heard it at Simla.

J. EVERSHERD.

Ewhurst, Surrey,
November 25.

Consumption of Fish by Porpoises.

In the course of our cruises, we have often harpooned porpoises of various species, and occasionally investigated the contents of the stomach. Sometimes the stomach was found to be empty, but in most cases it contained remains of fish, though these were, as a rule, so decomposed by the digestive fluids that identification was impossible. Now and again, however, it could be done. In the vicinity of the Continental Slope, for example, west of the English Channel, where porpoises are nearly always found in abundance, we found great bundles of the pelagic pipe-fish, *Entelurus æquoreus* L., in the stomachs of porpoises taken. But as a rule, the porpoise evidently prefers fish of somewhat more fleshy build than the pipe-fish.

The present note is occasioned by the recent preliminary investigation of a sample from one of the cruises of the *Thor*. On June 24, 1910, being then off the south coast of Spain, in the Mediterranean (between 36° 10' N., 4° 42' W., and 36° 19' N., 4° 06' W.), we harpooned a female specimen of the common long-nosed porpoise (*Delphinus delphis* L.). The stomach contents consisted of fish-residue: more or less dissolved soft parts, crumbling backbones, otoliths, and eye-lenses. I noted that most of the fish-bones were green, but no identification was attempted. The most interesting feature was the great number of otoliths, or ear-bones, of fish. When these were sorted out and counted, there were no fewer than 15,191 of different sizes, though mostly small. Several species were represented, about five; but, owing to lack of material for comparison, I cannot give any further determination at present. Some are presumably those of Scomberesox, clupeoids and scomberoids, possibly also scopolids.

The sample is interesting, inasmuch as it gives some slight idea of the porpoise's enormous consumption of fish: in the stomach of this one specimen we found remains of no fewer than 7596 fish. How long the porpoise took to collect the whole 7596 it is impossible to say, since we do not know how long the otoliths remain in the stomach before being dissolved or passed out. The fact that otoliths are not always found in the stomachs of porpoises seems rather to suggest that they do not remain there very long.

JOHS. SCHMIDT.

Crystallisation of Cementite in Steel.

WITH reference to the particularly interesting article by H. C. H. C. in NATURE of November 17, p. 728, might I mention the following amongst many other examples which have come under my notice illustrating the tendency of cementite to form cell walls or a network under conditions where the occurrence of pearlite is more commonly anticipated? In

FIG. 1.—Cementite network in mild steel. $\times 300$.

dead mild steels the occurrence of cementite in either network or comparatively massive formation has been recognised by a number of investigators. Fig. 1 illustrates an exceptional case in which isolated cell walls were found only near the edge of a dead mild-steel plate, in a region otherwise microscopically carbonless. The only apparent explanation of this occurrence was that the plate must have become carburised locally during the processes of manufacture.

FIG. 2.—Cementite in ferrite grain junctions, in nickel steel. $\times 1500$.

In the alloy steels, the simultaneous occurrence of structurally free ferrite and juxtaposed carbide network would be regarded as uncommon, but Fig. 2 shows an instance of this occurrence in a large oil-hardened nickel-steel forging. At the magnification of 2500 with which the original photograph was taken by means of the super-microscope, the carbon-containing constituent can readily be seen to be lamellar pearlite, which makes easy the identification of the carbide plates in the ferrite boundaries.

F. ROGERS.

64A Westbar, Sheffield.