D₂ alone shows polarisation. In addition, it may be pointed out that the $2p_1-2s$ and $2p_2-2s$ thallium lines show no polarisation in resonance radiation, as would be expected, since the s levels are above, while $2p_2-2d_2$ shows strong polarisation. The first and second doublets of the principal series of caesium and potassium show polarisation in resonance radiation, due presumably to the more refrangible member of the doublet in each case.

For certain types of spectral lines, the theory of the anomalous Zeeman effect shows that resonance radiation excited in a magnetic field by plane polarised light must itself be partially polarised, because of the absence of certain components of the Zeeman pattern (Foote, Ruark and Mohler, J.O.S.A. and R.S.I., 7, p. 415, 1923.) I have found that in the absence of a magnetic field, lines of this type, e.g. $1s-2p_1$ of sodium, $1s-2p_2$ of mercury, $2p_2-3d_2$ of thallium, show polarisation; while those in which all components of the Zeeman pattern are to be expected, as $1s-2p_2$ of sodium, $2p_1-2s$ and $2p_2-2s$ of thallium show no polarisation either with or without the field.

This work was done at the Johns Hopkins University, and is being continued at the Bureau of Standards.

A. ELLETT
(National Research Fellow).

Bureau of Standards,
Washington, D.C.,
November 15.

Heterogeneous Catalysis.

PROF. H. S. TAYLOR has pointed out to us that it is not clear from our paper on the heats of adsorption of oxygen on charcoal (Blench and Garner, J.C.S., 1924, 125, 1288) that the values were expressed in kgm.-cal. and also that we had not discussed the bearing of these results on the mechanism of heterogeneous catalysis. The latter point had, however, been reserved until further and more accurate data had been obtained, but, as it appears to be a matter of general interest, I propose to discuss it briefly in this letter.

The position, mode of orientation, and energy content of the atoms forming the surface of a homogeneous catalyst will very largely determine the part they play as individuals in the catalytic process, and it is not unlikely that a comparatively small number of the total atoms on the surface are the active agents in promoting the chemical change. This view is supported by the very large differences in energy content which have been shown to exist among the atoms on a carbon surface.

On admitting oxygen to an active carbon surface, heat is liberated which is due partly to the formation of carbon monoxide and dioxide and partly to the formation of an adsorption complex, C_xO_y . By allowing for the heats of formation of the two gases formed, it is possible to obtain a value for the "heat of formation" of the complex C_xO_y , which does not necessarily represent solely the heat liberated in its formation, but also includes any decrease in energy of the surface due to the liberation of the two gases. This heat of reaction, expressed in kgm.-cal./gm. mol. O₂ adsorbed, was often greater than the heat of formation of carbon dioxide. It decreased as the amount of gas admitted increased, and increased with the temperature of adsorption. The heat liberated ranged from 52 kgm.-cal. at 18° to 224 kgm.-cal. at 450° C. for the first small quantities of oxygen admitted. On subsequent admission of oxygen lower values than these were always obtained. These heats of reaction are of the same order as the energy liberated in the formation of the C-C linking as deduced from the

heat of sublimation of carbon. Thus the valency of the surface carbon atoms is very far from being saturated by neighbouring carbon atoms.

The ratio of the oxygen atoms adsorbed to those giving rise to carbon monoxide and dioxide decreases with increase in temperature; in other words, a proportion of the atoms adsorbed at the low temperatures is liberated as gases at the high temperatures. Thus the differences between the heats of adsorption at two temperatures must be due to different selections made from the carbon atoms on the surface, and hence are an indication of the very widely differing energy contents of these atoms. The oxygen atoms which remain on the surface at any one temperature will be to a large extent those held by a number of carbon atoms, while those adsorbed on exposed carbon atoms, which are less tightly bound to the surface, will give rise to the oxides of carbon. Thus the high heats of adsorption are not due solely to the energy liberated on combination of the oxygen atoms with the surface, but also to that liberated on the removal of the exposed carbon atoms, i.e. carbon atoms with a high energy content, from the surface. Both energies are undoubtedly very high compared with the heat of formation of carbon monoxide, but without further work it is difficult to decide between their relative importance. The most active atoms are, however, removed on the first admission of oxygen.

A point of difference between the reactions of carbon and oxygen and a catalytic reaction lies in the fact that the atoms of the catalyst do not leave the surface although their relative positions with respect to one another may change during the reaction. It is, however, clear that those atoms lying in an exposed position on the surface, by virtue of their high energy content, will play a large part in the chemical process. They will be the most likely to possess the energy necessary for promoting the trigger action of catalysis. Also it can be seen why the surface becomes increasingly effective with use. If part of the heat liberated during the heterogeneous reaction is absorbed by the surface, this may result not only in an increase in the total surface but also in an increase in the proportion of exposed atoms on the surface, i.e. an increased free energy of surface per sq. cm.

W. E. GARNER.

Chemical Laboratories,
University College,
Gower Street, London, W.C.1.

Seed Inoculation of Lucerne (*Medicago sativa*) and its Relation to the Motility of the Nodule Organism in Soil.

THE successful results that are being obtained in other countries from the inoculation of lucerne seed with the nodule organism (*Bacillus radicola*, Beyerinck) have stimulated investigation of this subject in England. In this practice the details of the method used in the inoculation are of great importance. The method consists essentially in making a suspension of the bacterial culture in some liquid, and in using this suspension to wet the seed. In this way each seed is coated with a film of liquid in which the bacteria lie. There is evidence that after penetrating the tissues of the root the bacteria are unable to pass for any significant distance along it. Consequently, when the seed has germinated, the bacteria must progress or be carried through the soil in order to reach from the seed-coat to the various parts of the root system, where nodules are to be formed. A study of the factors affecting the motility