issued in 1899. Up to that time I had been closely connected with the Cavendish for nearly twenty-five years and have written of things I knew. I must conclude now and leave to others more intimately acquainted than myself with the progress of the last quarter of a century to tell in detail what that progress has been. Thomson continued his work until 1918, through the troubled period of the War, when the researches of the Cavendish assisted to the full in the solution of many problems. In that year Dr. Butler, the revered Master of Trinity, died, and Thomson

was chosen by the Crown as his successor. He resigned the Cavendish professorship with its charge of the laboratory, and the University created for him a professorship of physics which will cease with his tenure of the post. Rutherford followed as professor, and under his genial direction he and his 'boys' from day to day increase the sum of natural knowledge by their amazing results and add in no small degree to the fame and influence of the Cavendish Laboratory, thus verifying the forecast of its founder more than fifty years ago.

Research Work in the Cavendish Laboratory in 1900-1918.

By Prof. J. A. CROWTHER.

THE progress of science and the life of a laboratory have little connexion with the artificialities of chronology, and the beginning of the century does not coincide with any natural division in the work of the Cavendish Laboratory. If, for the sake of clearness, we must make a division in what was fundamentally a natural and continuous growth, our most convenient date will be the year 1903, and our obvious landmark the publication of the first edition of Thomson's monumental book, "Conduction of Electricity through Gases." Here we find for the first time a complete and logical account of the solid achievements of the previous eight or ten years of remarkable activity. Here also we find the foreshadowings of the ideas which were to direct and regulate a large part of the work of the succeeding years.

In the main the problems attacked in the years before 1903 were those of the mechanism of the conduction of electricity through gases. Some of these, it is true, were carried over into the succeeding period. Some, perhaps, have not yet been fully solved, but by 1903 the foundations of the subject had been well and truly laid. It had been proved, beyond doubt, that the conductivity imparted to gases by various agents, such as X-rays, the rays of radium, and ultra-violet light was due to the formation of charged particles or gaseous ions in the gas, and many of the properties of these ions had been determined. Rutherford had determined their mobility; Townsend had determined their rate of diffusion through various gases; C. T. R. Wilson had shown that they could serve as nuclei for the condensation of water in a supersaturated space. Thomson had discussed their motion, and shown that in all probability they consisted of single molecules of the gas in which they were formed.

In a closely allied subject Thomson had demonstrated experimentally the existence of the electron, or negative corpuscle as we then called it, and had determined its few simple properties. The measurement of the electronic charge actually falls within the present century, though

preliminary experiments had been made somewhat earlier. Determinations were made independently by Thomson and by H. A. Wilson, by somewhat different methods, and were published in 1903. During the same period Zeleny made new measurements of ionic mobilities, which when combined with further measurements on the diffusion of ions by Townsend established conclusively the identity of the charge on a gaseous ion with that on a monovalent ion in solution. These experiments mark the culmination of the first epoch.

When I commenced research work in 1905 the second epoch of activity was already well under way. With the exception of C. T. R. Wilson, who in the advanced laboratory was engaged in imparting to a new generation of students those experimental arts of which he is a master, the first group of workers had all dispersed. Their work, as collected in the pages of "Conduction through Gases," formed our text-book, and was the foundation on which we hoped to build.

It was a very happy, though crowded, community to which I had the honour of being admitted. The space available for research had not been extended since the laboratory was built, and what had been adequate for ten or twelve, the average number of research workers before 1903, was becoming a tight fit now that the numbers had risen to an average of twenty-five. A competent observer records that there was more physics to the square centimetre in the Cavendish than anywhere else in the world, and the resources of the laboratory in the way of apparatus were sometimes severely tested, though I cannot recall that they ever broke down. The oft-repeated legend that we worked entirely with sealing wax and string, though flattering to our technique, has little foundation in fact. The apparatus for the new experiments which we wished to make did not exist. It had to be invented and made on the spot. An adequate workshop, staffed with extraordinarily competent and helpful mechanics, with F. Lincoln as chief, carried us through the worst of our difficulties; while Everett, the professor's private

assistant for so many years, was always willing to lend a hand in the worst intricacies of glass-blowing. For the rest there was the spirit of cheery fellowship and good comradeship which extended from the professor

FIG. 11.—Prof. C. T. R. Wilson's original expansion apparatus for making visible the tracks of ionising particles. It was made in the Cavendish Laboratory in 1911 and is the apparatus with which his photographs of the tracks have been obtained.

to the rawest recruit, and bound us all into a genuine community.

It is clearly impossible to attempt to give any adequate account of the activities of so many workers, extending for so many years, within the bounds of a single article. A mere list of publications would fill more space than I have at my disposal. I can only outline the general trend of discovery in the laboratory as I saw it, and many excellent pieces of research such as those of Burton on colloids, of Wellisch on ionic mobilities, of Vegard on osmosis, and of C. T. R. Wilson on atmospheric electricity, must pass unnoticed. This is the more unfortunate in that it may tend to accentuate the erroneous impression that the research work of the Cavendish has been entirely specialised and one-sided. This has never been true. Workers with ideas in any branch of physics have always received a warm welcome, and the Cavendish has rarely been without them. The majority of the research workers, however, came to the laboratory as young students, to learn their job, and it was natural and inevitable that their work should be inspired and directed by the genius of Thomson, and should be correlated to the subjects which he had most in mind.

The dominant problem from 1903 onwards was that of the structure of matter, to which the way had been opened by the discovery of the universality of the electron. Thomson's main theoretical contributions to the subject appeared in 1903, 1904, and 1906, and dealt with possible arrangements of electrons to form atomic structures, and the properties—electrical, magnetic, and optical—which might be expected to result from these arrangements. The substance of these papers reappeared, in a somewhat more popular form, in "Electricity and Matter" and the "Corpuscular Theory of Matter." To bring these ideas to the touchstone of experiment was, perhaps, the main pre-occupation of the laboratory until the year 1912.

A study of electrical conductors seemed to offer a hopeful line of approach. If the current through a metallic conductor is carried by electrons, which share in the thermal agitation of the molecules, some of them should escape from the surface if the temperature is sufficiently raised. Preliminary experiments on the subject were made by O. W. Richardson so early as 1901, and some of the theoretical consequences were discussed by Thomson in the following year. The subject was hotly pursued by O. W. Richardson in the Cavendish until 1906, and afterwards at Princeton, and has developed into the important branch of physics known as thermionics. Thomson, Horton,

FIG. 12.—The first mass-spectrograph set up by Aston in 1919. This instrument was capable of comparing the masses of atoms and molecules with an accuracy of 1 in 1000. By its means, in five years, more than fifty elements were resolved into isotopes or shown to be simple.

and others also made contributions to the subject. The preliminary work necessary to disentangle the genuine effect from the many spurious phenomena which accompany it was tedious and harassing, but important and valuable results were eventually obtained.

The question of the radioactivity of ordinary materials, which was being investigated in the laboratory about this time by N. R. Campbell and A. Wood, also had its bearings on the main problem. Difficulties were met with in eliminating the effects of penetrating radiations from outside sources and the possibility of small traces of radium in the materials employed, and the results were indeterminate, except in the case of some of the alkali metals, which were found to emit appreciable quantities of β -radiation. Experiments made by Kleeman on the relative ionisation produced by different ionising radiations in gases, by Laby on the total number of ions produced by the complete absorption of the α -rays from uranium in different gases, and by Kaye on the output of X-radiation from targets of various metals, while important in themselves, also had a more or less direct bearing on the main problem of the atom.

In 1906 Thomson returned to the attack and gave the theory of two important methods of estimating the number of electrons in the atom of a given element. One method suggested was to measure the diffusion of a parallel beam of β -particles in their passage through thin sheets of matter. Assuming that the deflexions are due to collisions with the electrons in the atoms through which they pass, a determination of the scattering of the particles should provide an accurate estimate of the number of scattering centres. A second method consisted in measuring the intensity of the scattered X-radiation from a given radiator when subjected to a beam of primary X-radiation. Here again, assuming the correctness of the classical theory of the phenomenon which had already been given by Thomson, it should be possible to deduce the number of scattering centres from the intensity of the radiation scattered. The experimental working out of both these methods fell to my lot, and kept me occupied from 1906 to 1911. The results showed that the number of electrons per atom was quite small, approximately of the order of the atomic weight, and that the main mass of the atom must be associated, not with its electrons, but with its positive charge.

In the meantime Thomson had himself been investigating the positive charge on the atom by various modifications of the method which had proved so successful with the electron. The results of these experiments on positive rays were published in various papers from 1908 to 1914. The importance of the new method lies in the fact that for the first time atoms could be dealt with as individuals, and not merely in aggregations of many millions. The results of these experiments, together with those of Aston (1919-1925) which led to the discovery of the existence of isotopes and the whole number law for the masses of the

atoms, are too fresh in the memory to need further description.

In the meantime new ideas were afloat. The phenomena of secondary corpuscular radiation excited by X-rays, as studied by Innes, and of photo-emission (Hughes and others) were becoming more and more impossible to reconcile with the classical wave theory of light. The theory that the wave front was discontinuous was put forward by Thomson so early as 1907, and experiments to test the matter were made by N. R. Campbell, and by G. I. Taylor in 1909. Laue's discovery in 1912 of the diffraction of X-rays by crystals placed a new weapon in the hands of physicists, and by 1913 W. L. Bragg, working in the Cavendish, had already employed it to measure the distance apart of the atoms in a rock-salt crystal, and to deduce an absolute value for the wave-lengths of various X-radiations. A little earlier (1911-1912), C. T. R. Wilson, by an ingenious modification of his early experiments on the deposition of water vapour on charged ions, had evolved a method of making visible the actual tracks of ionising particles in gases, a method the potency of which has not yet been fully exploited, although it is now in use in many laboratories in all parts of the world. Whiddington, by his experiments on the excitation of characteristic X-radiation by cathode rays (1912), was already approaching the quantum theory of X-ray phenomena. At the commencement of 1914 the stage seemed fully set for a new outburst of activity which should rival even the early days of the laboratory under Thomson.

Alas, however, the world stage was set for quite a different drama, and its research workers found problems for solution far different from those which they had been expecting to attempt. When the laboratory began to fill up again in 1919, it was to learn that its professor had found the double duty of directing at one and the same time a great laboratory and a great college too much for one man's strength and time, and the Cavendish professorship was resigned. This is not the end of the story. Sir J. J. Thomson still continues his theoretical and experimental researches with unabated vigour and success, and under its new chief, Sir Ernest Rutherford, the Cavendish goes on to new problems and new triumphs. But these changes may conveniently mark the end of the present article, which has already exceeded its allotted span, without doing anything like justice to the many old companions and friends by whose side I have worked in the Cavendish laboratory, and whose faces look down upon me from the framed groups upon my study walls. Having worked in the Cavendish they will understand the impossibility of the task I have undertaken, and being Cavendish men they will forgive.