equilibrium had been reached. However, instead of remaining quiescent, as one would expect, after a few seconds a sudden re-evolution of gas took place, which was of short duration, and then a state of equilibrium was apparently attained.

I have recently confirmed this observation, and some experiments have been carried out to see whether this phenomenon is of general occurrence with metals, or whether it is specific for copper. It has only been possible, so far, to examine the following metals: lead, iron, tin, platinum and aluminium, and in these cases no such action was observed under similar conditions.

It appears from these experiments that either:

(1) Copper undergoes an allotropic change with heat evolution at a temperature slightly above that of liquid air. One is reminded of the phenomenon of recalescence in the case of iron at higher temperatures.

Or (2) a surface action peculiar to copper comes into play.

Such an effect must cause a considerable loss in filling copper Dewar vessels with liquid air.

I have so far been unable to trace any record of this peculiar behaviour of copper in the literature.

GEORGE JAMES ALEXANDER.

Rock Ferry, Cheshire.

The Reaction to Flea Bites.

It does not seem to be generally appreciated that the irritating wheals which may follow the bites of various insects are anaphylactic in origin. The first time that a person is bitten by a species of insect which has never bitten him before, nothing or next to nothing may happen, but when he has become sensitive to the proteid in the liquid which the insect injects in the process of biting, he develops the local irritable swelling which is familiar to most people.

Some years ago (*J. Path. Bact.*, 17 (1912), 110) I showed that my wife, who gave a violent reaction to the bite of the human flea (*Pulex irritans*), did not respond at all to bites of the rat flea (*Xenopsylla cheopis*) until they had been fed on her five times during six weeks, when she developed extensive wheals. We have recently got the same results with the rabbit flea (*Spilopsyllus cuniculi*). Single fleas were fed on June 6 and June 18, and four fleas on July 11, without any result; after feeding two on August 13 red papules developed seven days later; single fleas were fed on August 22 and September 1, and in each case a moderate reaction developed about 36 hours later. It is rather remarkable that the minute quantity of foreign proteid injected by the fleas in four bites should be enough to make a person sensitive. Lack of sensitisation is not, of course, the only reason why people do not react. Some persons who are extensively flea bitten seem to become immune. Others are naturally immune: however often I am bitten by them, I give no response to fleas of any kind, though midges and some mosquitoes irritate me extremely. At least that is so in Hertfordshire; from my only experience of Scotch midges, in which I was bitten very freely, I had no trouble; they were probably a different species.

The rabbit flea does not in my experience bite man very readily; in some trials I have quite failed to get them to do so even after several days' starvation. When they do, they take a long time to get their feed, and if undisturbed may remain attached for so long as an hour. Brumpt ("Précis de parasitologie," 3rd ed., 1922, p. 829) notices this peculiarity, which seems to apply to their feeding on rabbits as well as man.

A. E. BOYCOTT.

October 2.

NO. 2973, VOL. 118]

Zoological Nomenclature: Hübner's (1806) 'Tentamen.'

THE secretary of the International Commission on Zoological Nomenclature has the honour to invite attention of the zoological profession to the fact that application has been made for the suspension of the International Rules, in the case of Hübner's (1806) 'Tentamen' in order to establish its nomenclatorial availability.

Briefly summarised: The formal nomenclatorial status of this document, involving about 100 names admitted by some authors as of generic rank, has been under controversy for many years, and opinion of specialists in Lepidoptera is still divided.

The arguments, as submitted, in favour of suspension of rules, maintain that: (1) there are sound reasons both for admitting and for denying recognition to the 'Tentamen' from the viewpoint of interpreting the Rules; (2) the evidence *pro* and *con* is not sufficiently conclusive to remove the question from debate; (3) the rejection of the 'Tentamen' will produce greater confusion than uniformity, will necessitate a vast amount of undesirable labour and economic loss of time and work; (4) if, on the ground of expediency, the rules can be suspended in this case, the nomenclature of the Lepidoptera, as used for the past thirty years, can be largely maintained.

The 'Tentamen' is one of the most important and most controversial cases ever submitted to the Commission. A discussion, with essential bibliographic references, will be found in *Smithsonian Misc. Coll.*, v. 73 (4) (now in press).

The Commission will delay announcement of vote, on the requested suspension, at least until September 1, 1927, in order to give interested authors, and especially entomological societies, opportunity to study the premises and to present to the Commission their views and arguments, *pro* or *con*, regarding the action requested.

In order to protect groups other than Lepidoptera, a prerequisite to suspension of rules would be that representative specialists in Lepidoptera agree upon and furnish to the Commission definite bibliographic references to the 107 names which they view as genotypes.

C. W. STILES,

Secretary to Commission.

U.S. Hygienic Laboratory,
Washington, D.C.,
August 21.

The Problem of the Origin of Species as it appeared to Darwin in 1859 and as it appears to us To-day.

I HAVE discovered two errors in my recent paper which appeared under this title in *NATURE* of August 21, 1926. These errors were due to the hurried assemblage of facts and citations under the heading "Geographic Isolation and Speciation"; they will be carefully corrected in a sequel to this paper which is now in preparation for the *American Naturalist*, to be published under the title "The Origin of Species, V.: Speciation and Mutation." Meanwhile, I desire to point out as inaccurate the following statement in my paper:

"As an example of wide isolation without speciation, the house wren of Florida, *T. ædon ædon*, exhibits the same characters as those of Tierra del Fuego, but in the intermediate regions another species, *T. musculus*, exhibits a large amount of subspeciation and several complete intergradations."

On the contrary, the house wren is not known to breed in Florida and does not occur in Tierra del