

Arsenic trioxide, a therapeutic agent for APL

Ting-Dong Zhang^{1,3}, Guo-Qiang Chen^{2,3}, Zhu-Gang Wang², Zhen-Yi Wang², Sai-Juan Chen² and Zhu Chen^{*,2}

¹The First Hospital affiliated to Harbin Medical University, 23 You Zheng Road, Nangang District, Harbin, 150001, PR China;

²Shanghai Institute of Hematology, Rui Jin Hospital affiliated to Shanghai Second Medical University, 197, Rui Jin Road II, Shanghai 200025, PR China

Acute promyelocytic leukemia (APL) is an interesting model in cancer research, because it can respond to the differentiation/apoptosis induction therapy using all-trans retinoic acid (ATRA) and arsenic trioxide (As₂O₃). Over the past 5 years, it has been well demonstrated that As₂O₃ induces a high complete remission (CR) rate in both primary and relapsed APL patients (around 85~90%). The side effects are mild to moderate in relapsed patients, while severe hepatic lesions have been found in some primary cases. After CR obtained in relapsed patients, chemotherapy in combination with As₂O₃ as post-remission therapy has given better survival than those treated with As₂O₃ alone. The effect of As₂O₃ has been shown to be related to the expression of APL-specific PML–RAR α oncoprotein, and there is a synergistic effect between As₂O₃ and ATRA in an APL mouse model. Cell biology studies have revealed that As₂O₃ exerts dose-dependent dual effects on APL cells. Apoptosis is evident when cells are treated with 0.5~2.0 μ M of As₂O₃ while partial differentiation is observed using low concentrations (0.1~0.5 μ M) of the drug. The apoptosis-inducing effect is associated with the collapse of mitochondrial transmembrane potentials in a thiol-dependent manner, whereas the mechanisms underlying APL cell differentiation induced by low dose arsenic remain to be explored. Interestingly, As₂O₃ over a wide range of concentration (0.1~2.0 μ M) induces degradation of a key leukemogenic protein, PML–RAR α , as well as the wild-type PML, thus setting up a good example of targeting therapy for human cancers. *Oncogene* (2001) 20, 7146–7153.

Keywords: arsenic trioxide (As₂O₃); acute promyelocytic leukemia (APL); apoptosis; differentiation; PML–RAR α

Introduction

Arsenic has long been known to act as a carcinogen, involved in human skin, lung, liver, kidney and urinary bladder cancers (Huff *et al.*, 2000). Paradoxically, arsenic has also long been demonstrated to have anti-cancer activity in some cases. In traditional Chinese medicine, arsenic trioxide (As₂O₃) was recorded in the Compendium of Materia Medica by Mr Li Shi-Zhen (1518–1593). In Western medicine, arsenous oxide (Fowler's solution) was used as a treatment of choice for chronic myeloid leukemia (CML) in the 19th century. As Osler stated in his 1892 textbook of medicine, 'There are certain remedies that have an influence upon the disease. Of these, arsenic, given in large doses, is the best. I have repeatedly seen improvement under its use.' However, due to toxic side effects of long-term heroic-dose of oral arsenic in most patients, and with the advent of modern radiotherapy and chemotherapy, the arsenic treatment for CML was given up in Western medicine (Kwong and Todd 1997; Tamm *et al.*, 1998).

Recently, a discovery on the therapeutic effect of As₂O₃ in acute promyelocytic leukemia (APL) has revived this ancient drug (Chen *et al.*, 1996; Mervis, 1996; Look, 1998; Gallagher, 1998; Conrad, 1999; Warrell, 1999; Slack and Rusiniak, 2000; Wang and Chen, 2000; Chen *et al.*, 2001). APL is a special subtype of acute myeloid leukemia (AML), characterized by the specific chromosomal translocation t(15;17) with resultant leukemogenic PML–RAR α fusion gene, and the clinical response to differentiation therapy using *all-trans* retinoic acid (ATRA) (Melnick and Licht, 2000). Over the past several years, As₂O₃ as an effective salvage treatment for relapsed and/or refractory APL patients has been confirmed worldwide. Recently, this drug was formally approved by the State Drug Administration (SDA) in China (1999) and the Food and Drug Administration (FDA) of the United States of America (Cohen *et al.*, 2001). Encouraged by such a bedside discovery, mechanisms of action of As₂O₃ have been actively addressed by many groups. The potentials of As₂O₃ in the treatment of cancers other than APL are also under exploration. Here, we shall discuss the recent advances in the clinical practices and the mechanisms of action of As₂O₃ in APL.

*Correspondence: Z Chen, Shanghai Institute of Hematology, Rui Jin Hospital affiliated to Shanghai Second Medical University, 197, Rui Jin Road II, Shanghai 200025, PR China;
E-mail: zchen@ms.stn.sh.cn

³Ting-Dong Zhang and Guo-Qiang Chen contributed equally to this work.

Clinical practice of As₂O₃ in the treatment of APL

Clinical efficacy and pharmacokinetics

The recent clinical trial of As₂O₃ was started in 1971. A group from the First Hospital affiliated with the Harbin Medical University, China, used this compound through intravenous administration in more than 1000 cases of different types of cancer. Therapeutic effects were observed in several cancer types, including CML, lymphoma, esophageal cancer, and particularly APL. The preliminary result in APL was reported in 1992 (Sun *et al.*, 1992). A preparation containing arsenic with a trace amount of mercury chloride was administered at a dose of 10 mg As₂O₃/day and could induce clinical complete remission (CR) in 21 out of 32 (65.6%) patients without major toxic side effects. In 1996, two reports were made from China using purified As₂O₃ (0.16 mg/kg/day). The Harbin group (Zhang *et al.*, 1996) showed that As₂O₃ treatment achieved CR in 52.4% (22/42) relapsed APL patients. The Shanghai Institute of Hematology (SIH) reported an even better result among 15 patients relapsed after ATRA and chemotherapy, with CR achieved in nine of 10 patients treated with As₂O₃ alone and five of five patients treated with As₂O₃/chemotherapy combination. The treatment duration for induction of remission was 28–54 days. Of note, the only non-responder in the SIH series had leukemic cells losing t(15;17) and PML–RAR α gene expression at relapse, although the disease was a typical APL at initial diagnosis, and the first CR was induced by ATRA (Chen *et al.*, 1996; Shen *et al.*, 1997).

The pharmacokinetic studies were performed among eight relapsed patients by measuring plasma drug levels with gas chromatography. The results showed that after a peak level of 6.85 (5.54–7.30) μ mol/L, plasma arsenic was rapidly eliminated, with a $t_{1/2\alpha}$ of 0.89 (SD: 0.29) h and a $t_{1/2\beta}$ of 12.13 (SD: 3.31) h. Hence, over most times during As₂O₃ treatment, plasma arsenic levels fluctuated between 0.1–0.5 μ M. Such a pharmacokinetic behavior did not alter after continuous use of As₂O₃. Increased amounts of arsenic appeared in the urine with a daily excretion accounting for approximately 1–8% of the total daily dose used. Arsenic contents in hair and nails were increased during treatment and declined after withdrawal of the drug (Chen *et al.*, 1996; Shen *et al.*, 1997).

The Memorial Sloan-Kettering Cancer Center of New York was the first to use As₂O₃ in the treatment of APL patients in Western countries. In the first report (Soignet *et al.*, 1998), eleven of the 12 patients had a CR after treatment with 0.06–0.2 mg As₂O₃/kg/day for 12–39 days. A more recent U.S. multicenter study performed in 40 patients achieved a CR rate of 85% (Cohen *et al.*, 2001). In addition, the effectiveness of As₂O₃ in the treatment of APL was also reported by some small-scale studies (Huang *et al.*, 1998; Agis *et al.*, 1999; Galimberti *et al.*, 1999; Ohnishi *et al.*, 2000; Zhang *et al.*, 2000; Lin *et al.*, 2000a).

Recently, a trial using low dose As₂O₃ (0.08 mg/kg/day) was carried out in SIH (Shen *et al.*, 2001). Among

20 cases studied, a similar CR rate (80%) was achieved as compared to that attained with a conventional dose. Pharmacokinetic study was conducted in six patients given low dose As₂O₃. The mean C_{pmax} was 2.63 (range, 1.54–3.42) μ mol/L, nearly half of the levels with a standard dose, and the ranges of $t_{1/2\alpha}$ and $t_{1/2\beta}$ were 1.2–2.7 h and 6.23–14.9 h, respectively. Of note, the arsenic concentrations in the plasma of bone marrow (BM) in five patients 24 h after administration of the drug at low dose were 0.061–0.49 μ mol/L.

Improvement of bleeding diathesis with As₂O₃ treatment

Clinically, in patients with APL, the disease is frequently accompanied by a bleeding syndrome, which is often exacerbated by chemotherapy and causes death at the early stage of treatment. It was suggested that the aberrant expression of tissue factor (TF) in APL cells is at least one of the mechanisms underlying the pathogenesis of the APL coagulopathy (Barbui *et al.*, 1998; Cheng *et al.*, 1999; Tallman, 1999). In APL patients receiving As₂O₃ treatment, the improvement of disseminated intravascular coagulation (DIC) or hyperfibrinolysis paralleled the amelioration of bleeding symptoms, the correction of plasma fibrinogen level, and the decrease in membrane procoagulant activity (PCA) as well as TF contents of APL blasts (Zhu *et al.*, 1999; Agis *et al.*, 1999). Consistent with the *in vivo* findings, the membrane PCA and TF at both protein and mRNA levels in APL cell line NB4 were rapidly down-regulated by 1 μ M As₂O₃ (Zhu *et al.*, 1999; Ohsawa *et al.*, 2000).

Toxic side effects

During As₂O₃ treatment in APL patients, there exists no BM depression. Instead, As₂O₃ induces leukocytosis in about 50% of patients (Shen *et al.*, 1997; Soignet *et al.*, 1998; Niu *et al.*, 1999; Camacho *et al.*, 2000). These patients can even develop retinoic acid syndrome (RAS)-like symptoms such as fever, skin rash, edema, which were quickly relieved by steroid administration (Lin *et al.*, 2000b). The leukocytosis could resolve in all cases without chemotherapy (Camacho *et al.*, 2000). Other mild side effects were encountered in about 40–50% of relapsed patients, such as fatigue, fever, edema, nausea, anorexia, diarrhea, emesis, headache, insomnia, cough, dyspnea, dermatitis, tachycardia, pain, hypokalemia, hypomagnesemia, and hyperglycemia. The most common (>10%) Grade 3 or 4 adverse events were abdominal pain, epistaxis, dyspnea, hypoxia, bone pain, thrombocytopenia, neutropenia, hypokalemia, and hyperglycemia (Shen *et al.*, 1997; Soignet *et al.*, 1998; Cohen *et al.*, 2001). Of note, evident polyneuropathy compatible with chronic arsenic toxicity was noted in a few patients who received As₂O₃ maintenance therapy and one even had marked distal muscular atrophy (Huang *et al.*, 1998). Cardiac dysrhythmias were reported with arsenic use (Huang *et al.*, 1999). In a recent report from Japan, prolonged QT intervals were observed in all patients during

induction therapy with As₂O₃ and ventricular premature contractions were noticed during eight of 12 courses of therapy (Ohnishi *et al.*, 2000). QT prolongation was also seen in 16 (38%) patients in the study from FDA (Cohen *et al.*, 2001), which could return to baseline following cessation of As₂O₃.

In a recent clinical study conducted in SIH (Niu *et al.*, 1999), 11 newly diagnosed APL patients received As₂O₃ treatment with informed consent. The CR was obtained in eight (72.7%) of them. However, As₂O₃ treatment resulted in elevated plasma liver transaminase levels in seven cases and two died of severe hepatic toxicity, in contrast to the mild liver dysfunction in about one third of the patients treated for relapsed APL. It was thus recommended that ATRA should be used as the first line drug for remission induction in newly diagnosed APL cases, while As₂O₃ can be either included into multi-drug consolidation/maintenance clinical trials after the first CR or used as a rescue for relapsed cases.

Post-remission treatment among relapsed APL patients

In a report by SIH, 33 patients subject to distinct post-remission treatment protocols after arsenic-induced CR were followed-up for 8–48 months (Niu *et al.*, 1999). Three groups were formed, including four cases treated with chemotherapy alone (group A), 18 with As₂O₃ alone (group B) and 11 with both As₂O₃ and chemotherapy (group C). Although the estimated disease-free survival (DSF) rates at 1 and 2 years in all patients were 63.6% and 41.6%, respectively, recurrence of leukemia developed in 3/4 cases in group A, 12/18 in group B and 2/11 in group C. Combining chemotherapy and As₂O₃ seems able to decrease the relapse and to yield a statistically significant better survival status compared to arsenic alone as post-remission treatment (Niu *et al.*, 1999). Recent data showed that the estimated 2-year DFS (49.1%) was not different in the low dose As₂O₃ treatment group as compared with that of the standard dose (Shen *et al.*, 2001).

Molecular remission

RT-PCR analysis in both newly diagnosed and relapsed groups showed that application of As₂O₃ for a short period of time was not sufficient to eliminate minimal residual disease. Long-term use of the drug could, nevertheless, lead to a molecular remission at least in some patients (Niu *et al.*, 1999). Of five patients reported by Zhang *et al.* (2000), two achieved molecular remission (negative RT-PCR for *PML-RARα* fusion transcripts) by the end of the second and third cycles of As₂O₃ therapy.

Dual activities of As₂O₃ on APL cells *in vivo* and *in vitro*: induction of apoptosis and differentiation

The clinical effectiveness of As₂O₃ in APL has stimulated research activities aiming at an under-

standing of its mechanisms of action. Although many questions remain to be understood, *in vivo* and *in vitro* data suggested that induction of apoptosis and partial differentiation of APL cells is likely to constitute the cellular basis of the effects of As₂O₃ (Figure 1).

Evidences from APL patients

Initial clinical observations showed that during daily continuous intravenous infusion of As₂O₃ in APL, a large amount of myelocyte-like cells and degenerative, dying cells with condensed nuclei losing cytoplasm appeared in both BM and peripheral blood, with the gradual reduction of leukemic promyelocytes. Interestingly, the myelocyte-like cells were derived from abnormal promyelocytes because they still carried the *PML-RARα* gene as revealed by the fluorescence *in situ* hybridization (FISH) analysis. Those degenerative cells were positively labeled by *in situ* terminal deoxynucleotidyl transferase, which is characteristic of programmed cell death (Chen *et al.*, 1997). The presence of leukocytosis and RAS-like syndrome during arsenic treatment also supports the idea that As₂O₃ could induce differentiation. A similar finding was described by Huang *et al.* (1998). Soignet *et al.* (1998) performed a further *in vivo* analysis on the mechanisms of action of As₂O₃. They reported that As₂O₃ therapy induced a progressive decrease in the proportion of cells expressing CD33, an antigen associated with primitive myeloid cells, along with an increase in the proportion of cells expressing CD11b, an antigen restricted to mature myeloid elements. As₂O₃ also induced the increment of cells that simultaneously expressed both antigens and displayed a hybrid signal of *PML* and *RARα* genes on FISH analysis. Interestingly, serial Western blot analysis of BM mononuclear cells during As₂O₃ treatment showed induced expression of the precursor forms of caspase 2 and caspase 3 and activation of both caspase 1 and caspase 3.

Evidences from APL models and *in vitro* studies

Using syngenic grafts of leukemic blasts from *PML-RARα* transgenic mice, Lallemand-Breitenbach *et al.* (1999) demonstrated that As₂O₃ induced apoptosis and modest differentiation of APL cells *in vivo*, and significantly prolonged the survival of the diseased animals. In subcutaneous tumors which were formed by implantation of ATRA-resistant UF-1 APL cells into hGM-CSF-producing transgenic SCID mice (Kinjo *et al.*, 2000), tumor size was decreased to half the initial one after As₂O₃ treatment, with the appearance of cells showing typical apoptotic morphology. Meanwhile, one of the As₂O₃-treated mice showed mature granulocytes in the diminished tumor.

In order to approach the complex mechanisms, *in vitro* effects of pharmacologic concentrations of As₂O₃ on fresh APL cells and APL cell lines including ATRA-sensitive NB4 cells, and ATRA-resistant NB4-derived sublines MR2, R1 and R2, were studied. The initial study showed that As₂O₃ at 1.0–2.0 μM could sig-




Figure 1 Working hypothesis for the mechanism of action of As_2O_3 in remission induction of APL. As_2O_3 induces remission of APL patients via two independent pathways: i.e., by triggering partial differentiation and inducing apoptosis. High-dose As_2O_3 -induced apoptosis involves mainly the -SH groups-related $\Delta\Psi_m$ collapse due to the opening of the PT pore, which leads to the release of pre-apoptotic factors from mitochondria to cytoplasm, followed by caspase activation and degradation of specific substrates. On the other hand, low-dose As_2O_3 -induced differentiation might be related to cyclic AMP pathway, nuclear receptor activity and/or regulation of histone acetylation. As_2O_3 at both high- and low-dose triggers the degradation of PML-RAR α . PML-RAR α oncoprotein may therefore represent the common target for both ATRA and As_2O_3 . However, the underlying mechanism seems to be different, the former targeting RAR α through specific binding whereas the latter targeting PML moiety through SUMO-1-conjugation in a yet unknown manner. \rightarrow : Stimulation. \neg : Inhibition

nificantly induce apoptosis in APL cells regardless of their sensitivity to ATRA (Chen *et al.*, 1996, 1997), which was widely confirmed by other groups (Kitamura *et al.*, 1997, and see later). In order to establish an *in vitro* model of differentiation, NB4 cells were treated with 0.1–0.25 μM of As_2O_3 over a long time of culture. Indeed, after 6–9 days of treatment, cells started to present morphologic differentiation with CD11b expression and differentiation-associated cytochemical features. The differentiation seems only a partial one since most cells were blocked at the myelocyte to metamyelocyte stage of maturation and no significant increase of nitroblue-tetrazolium (NBT) reduction, a classical marker of granulocytic maturation, was observed in NB4 cells (Chen *et al.*, 1997). Later, Gianni *et al.* (1998) isolated an As_2O_3 -resistant NB4 subline (NB4-AsR). These cells failed to undergo apoptosis, but maintained the partial differentiation response to arsenic. Of note, ATRA-triggered differentiation and apoptosis were accelerated in As_2O_3 -treated NB4-AsR cells.

Mechanisms of As_2O_3 -induced apoptosis

A relatively wide spectrum of As_2O_3 -induced apoptosis

Several groups reported that HL-60 and U937 cell lines, both are of myeloid lineage, had weak responses to

1 μM As_2O_3 in terms of growth and survival inhibition (Chen *et al.*, 1996; Kitamura *et al.*, 1997; Zhang *et al.*, 1998), but a few groups observed that As_2O_3 could induce apoptosis in these cells (Wang *et al.*, 1998). Although different culture situations and/or variations of cell lines might be responsible for such a discrepancy (McCabe *et al.*, 2000), it may suggest that the cell type specificity of As_2O_3 -induced apoptosis is limited. Indeed, over the past few years, several studies found that As_2O_3 could also inhibit cell growth and/or induce apoptosis among some lymphoma and solid tumor cell lines in a dose-dependent way. As_2O_3 could also cause apoptosis in proliferating layers of human umbilical vein endothelial cells and prevent capillary tubule and branch formation under *in vivo* and *in vitro* assay conditions (Roboz *et al.*, 2000), raising the possibility of inhibiting metastasis. However, it should be noted that significant induction of apoptosis in cell systems other than APL may sometimes require high concentration As_2O_3 (5–10 μM), a dose difficult to be achieved *in vivo*.

Mitochondria are involved in As_2O_3 -induced apoptosis

The fact that As_2O_3 induces apoptosis in a variety of cell types suggests that some common mechanisms should be involved. It is well established that mitochondria play a major rate-limiting role in apoptosis. Various pro-apoptotic signals converge on

the mitochondria and trigger progressive disruption of mitochondrial transmembrane potentials ($\Delta\Psi_m$) by modulating the permeability transition pore complex (PTPC), a multiprotein complex which interacts with the apoptosis-regulating protein Bcl-2/Bax family (Kroemer and de Thé, 1999; Costantini *et al.*, 2000a). As do most chemotherapeutic drugs, As_2O_3 can also induce an early $\Delta\Psi_m$ collapse in intact APL cells (Cai *et al.*, 2000), malignant lymphocytes (Zhu *et al.*, 1999) and carcinoma cell lines (Shen *et al.*, 1999; Seol *et al.*, 2001). As_2O_3 -induced nuclear apoptosis also required mitochondria in an *in vitro* cell-free system, and As_2O_3 acted on isolated mitochondria to induce PTPC opening and opened the purified, reconstituted MPT *in vitro* in a cyclosporin A- and Bcl-2-inhibitable fashion (Larochette *et al.*, 1999). These data support that mitochondria, PTPC in particular, are important target of As_2O_3 . Besides, in the APL model, glutathione (GSH)-reducing agent dithiothreitol (DTT) could effectively block As_2O_3 -induced $\Delta\Psi_m$ collapse and apoptosis. In contrast, buthionine sulfoximine (BSO), a depleter of cellular GSH, and ascorbic acid substantially enhanced the effect of As_2O_3 (Zhu *et al.*, 1999; Dai *et al.*, 1999; Jing *et al.*, 1999; Cai *et al.*, 2000). BSO could also make As_2O_3 -resistant HL60 and U937 cells to become sensitive to the compound. Therefore, it is presumed that As_2O_3 -induced $\Delta\Psi_m$ collapse and apoptosis are associated with dithiol oxidation or cross-linking, which has been shown to be associated with a higher probability of PTPC opening (Figure 1, lower part). Initially, we speculated that adenine nucleotide translocator (ANT) may be a target of As_2O_3 (Cai *et al.*, 2000), since it is an important component of PTPC and contains the vicinal thiol group. In addition, ANT alone reconstituted into artificial lipid bilayers suffices to confer a membrane permeabilization response to thiol cross-linking agents. However, a recent report showed that As_2O_3 could not cause the oxidation of a critical cysteine residue (Cys⁵⁶) of purified ANT (Costantini *et al.*, 2000a). In addition to the direct effects on PTPC, As_2O_3 could disrupt mitochondria through some indirect mechanisms, such as the induction of reactive oxygen species (ROS). Nevertheless, there are controversies in literature with regard to the role of ROS. It was reported that the increased intracellular peroxide levels were accompanied by As_2O_3 -induced apoptosis, which could be inhibited significantly by N-Acetyl-L-cysteine (a thiol-containing antioxidant), diphenylene iodonium (an inhibitor of NADPH oxidase), 4,5-dihydro-1,3-benzene disulfonic acid (a selective scavenger of O_2^-) and catalase (Chen *et al.*, 1998). However, other data showed that DTT blocked apoptosis induced by As_2O_3 , but it could not antagonize As_2O_3 -induced ROS production (Zhu *et al.*, 1999). In the experiments with butylated hydroxyanisole, Uslu *et al.* (2000) suggested that the apoptotic activity of As_2O_3 in prostate and ovarian cancer cells was not through inducing superoxide generation.

Next to PTPC opening is the release of pro-apoptotic factors from mitochondria into the cytosol,

where they activate sequentially upstream and downstream caspases such as caspase-9 (Costantini *et al.*, 2000b; Brenner and Kroemer, 2000). Indeed, As_2O_3 could induce the release of cytochrome *c* from mitochondria to the cytosol (Chen *et al.*, 1998; Jing *et al.*, 1999). The activation of caspase-3 and cleavage of its specific substrates such as poly (ADP-ribose) polymerase (PARP) could also be seen in most cells treated with As_2O_3 (Akao *et al.*, 1999; Huang *et al.*, 2000; Cai *et al.*, 2000; Jing *et al.*, 1999) (Figure 1, lower part). In addition to caspase-3, Kitamura *et al.* (2000a) showed that caspase 8 and Bid were also activated by As_2O_3 in a GSH concentration-dependent manner in NB4 cells. An inhibitor of caspase 8 blocked not only the activation of caspase 3 but also the loss of $\Delta\Psi_m$. However, Sternsdorf *et al.* (1999) realized that As_2O_3 -induced apoptosis was caspase 3 independent and was only partially blocked by a global caspase inhibitor. In experiments conducted by SIH, caspase-3 was activated in 2 μM As_2O_3 -treated Namalwa and Raji cells, but not in 1 μM of As_2O_3 -treated BJAB, MOLT-4 and SKW-3 cells despite significant apoptosis, indicating that caspase-3 was not always necessarily involved in As_2O_3 -induced apoptosis (Zhu *et al.*, 1999).

Apart from the effect on mitochondria, as a pleiotropic agent, As_2O_3 may have additional actions that also contribute to the induction of cell death, such as the activation of c-Jun NH₂-terminal kinases (Huang *et al.*, 1999) and inactivation of the GTP binding site of monomeric tubulin (Li and Broom, 1999). Although As_2O_3 does not increase p53 level or activity in most cells, a marked increase in p53 protein level was seen during As_2O_3 -induced apoptosis in a human gastric cancer cell line, while co-incubation with p53 anti-sense oligo-nucleotide suppressed both p53 over-expression and apoptosis induced by As_2O_3 (Jiang *et al.*, 2001). Hence, mechanisms of action of As_2O_3 may be very complex, depending on dose, cell type, or cellular environment (McCabe *et al.*, 2000).

Mechanisms of As_2O_3 induced-differentiation of APL

As previously mentioned, during the 0.08–0.16 mg/kg/day i.v. As_2O_3 treatment, plasma As_2O_3 levels were basically in the range of 0.1–0.5 μM , corresponding to the concentrations inducing primarily differentiation of APL cells *in vitro* (Chen *et al.*, 1996, 1997; Shen *et al.*, 1997, 2001). However, the mechanism underlying As_2O_3 -induced differentiation is still poorly understood. The fact that arsenic-resistant NB4-AsR subline failed to undergo apoptosis, but maintained the partial differentiation response to As_2O_3 suggested that distinct pathways may be responsible for the differentiation and apoptosis (Gianni *et al.*, 1998). Recently, we showed that 0.1 μM As_2O_3 did not induce differentiation in ATRA-resistant APL cell lines. Furthermore, low-dose arsenic could induce partial differentiation of ATRA-sensitive, but not ATRA-resistant, HL-60 cells. In addition, the gene expression patterns modified by low-dose As_2O_3 overlapped with

those by physiological concentrations of ATRA (Cai *et al.*, 2000). These data suggested a possible cross-talk between the retinoid signaling and the regulatory pathways modulated by As₂O₃ (Figure 1, upper part). Of note, *in vivo* studies proposed that As₂O₃ and ATRA generate synergistic effect, such as an accelerated tumor regression and prolonged survival of animals (Lallemand-Breitenbach *et al.*, 1999; Kinjo *et al.*, 2000; Rego *et al.*, 2000), although the benefit of their combined use in cell culture is controversial (Chen *et al.*, 1996; Gianni *et al.*, 1998; Shao *et al.*, 1998; Jing *et al.*, 2001). Nevertheless, As₂O₃ showed no direct effect on RA receptors, in that the interaction of wild-type RAR α and PML–RAR α with nuclear receptor co-repressor was not affected, and arsenic showed no obvious effect on the transactivation activity of RAR α in an RARE-luciferase reporter system (Chen *et al.*, 2001 and unpublished data). Interestingly, as will be discussed below, the differentiation of APL cells seems to be closely related to the modulation of PML–RAR α , the key protein in leukemogenesis.

Modulation of PML and PML–RAR α

The unique response of APL to As₂O₃ implies the importance of intrinsic properties of APL cells. Interestingly, both wild-type PML and PML–RAR α fusion protein have been shown to be major targets of arsenic treatment. PML is a tumor suppressor involved in complex functions including growth arrest and apoptosis induction. Normally PML is located in the nucleus on a specific subdomain named PML nuclear body (NB). In APL cells, due to the heterodimerization of PML–RAR α with wild-type PML, PML–NBs are disrupted into nuclear micro-speckle pattern, with loss of PML functions (Melnick and Licht, 2000; Andre *et al.*, 1996). A surprising finding was that 0.1–2.0 μ M of As₂O₃ targets PML and PML–RAR α onto NBs, followed by degradation of these proteins (Chen *et al.*, 1996, 1997; Zhu *et al.*, 1997; Sternsdorf *et al.*, 1999; Gianni and de Thé, 1999). Furthermore, As₂O₃ increases covalent modification of PML by SUMO-1, a small ubiquitin-like protein. The SUMO-1-polymodified forms of PML are compartmentalized exclusively in the PML–NBs in non-APL cells, while the unmodified form of PML is found in the soluble nucleoplasmic fraction (Muller *et al.*, 1998; Sternsdorf *et al.*, 1999). Therefore, As₂O₃-enhanced PML sumoylation constitutes the basis of the restoration and accumulation of the PML in enlarged NBs in As₂O₃-treated APL cells (for more

information, please refer to the review article by Dr H de Thé in the same issue) (Figure 1).

There are controversies on the role of modulation of PML/PML–RAR α proteins in As₂O₃-induced apoptosis in APL cells. Through analysis of U937 cells stably expressing exogenous PML–RAR α gene, Sternsdorf *et al.* (1999) reported that As₂O₃ could induce apoptosis only in U937 cells expressing the PML–RAR α but not in cells expressing PLZF–RAR α , a fusion gene generated as a result of the variant translocation t(11;17)(q23;q21) in APL. However, degradation of PML–RAR α and change of the PML-subcellular localization were similarly induced by As₂O₃ in NB₄ cells and NB₄-derived As₂O₃-resistant subline (Kitamura *et al.*, 2000a). Therefore, the contribution of PML–RAR α and PML modulation to apoptosis under the arsenic treatment seems to be relative even if present. In contrast, the degradation of PML–RAR α may play a crucial role in the differentiation of APL cells, since this protein modulation can occur in the presence of very low concentration (0.1–0.25 μ M) of As₂O₃, while arsenic induced neither differentiation of APL cells with t(11;17) nor degradation of PLZF–RAR α protein (Koken *et al.*, 1999; Kitamura *et al.*, 2000b). The low-dose arsenic shows marked cell type selectivity, since it causes maturation of APL cells but exhibits little effect on other cancer cells, although a recent report suggested that low-dose As₂O₃ can increase the lysis of myeloma cells by lymphokine-activated killer cells (Deaglio *et al.*, 2001). Importantly, there is also a marvelous target selectivity of As₂O₃, in that only the PML–RAR α , but not the wild-type RAR α , is degraded. The relief of the transcriptional repression of PML–RAR α may pave the way for the operation of physiological RA signaling in granulocytic differentiation, particularly at the stage of promyelocytes (Figure 1, upper part). The possible synergistic effect of the arsenic with a number of important regulatory pathways such as RAR/RXR (Cai *et al.*, 2000), Cyclic AMP (Zhang JW, Zhu Q, Tong JH, unpublished data) and histone deacetylase inhibitor may also enhance the restoration of the cell differentiation program.

Acknowledgments

This work is supported in part by the Chinese National Key Project for Basic Research (973), Chinese National High Tech Program (863), National Natural Science Foundation of China, Samuel Waxman Cancer Research Foundation and Clyde Wu Foundation of Shanghai Institute of Hematology.

References

- Agis H, Weltermann A, Mitterbauer G, Thalhammer R, Edelhauser M, Seewann HL, Valent P, Lechner K, Fonatsch C and Geissler K. (1999). *Ann. Hematol.*, **78**, 329–332.
- Akao Y, Nakagawa Y and Akiyama K. (1999). *FEBS Lett.*, **455**, 59–62.
- Andre C, Guillemain MC, Zhu J, Koken MH, Quignon F, Herve L, Chelbi-Alix MK, Dhumeaux D, Wang ZY, Degos L, Chen Z and de Thé H. (1996). *Exp. Cell Res.*, **229**, 253–260.
- Barbui T, Finazzi G and Falanga A. (1998). *Blood*, **91**, 3093–3102.

- Brenner C and Kroemer G. (2000). *Science*, **289**, 1150–1151.
- Cai X, Shen YL, Zhu Q, Jia PM, Yu Y, Zhou L, Huang Y, Zhang JW, Xiong SM, Chen SJ, Wang ZY, Chen Z and Chen GQ. (2000). *Leukemia*, **14**, 262–270.
- Camacho LH, Soignet SL, Chanel S, Ho R, Heller G, Scheinberg DA, Ellison R and Warrell RP. (2000). *J. Clin. Oncol.*, **18**, 2620–2625.
- Chen GQ, Zhu J, Shi XG, Ni JH, Zhong HJ, Si GY, Jin XL, Tang W, Li XS, Xiong SM, Shen ZX, Sun GL, Ma J, Zhang P, Zhang TD, Gazin C, Naoe T, Chen SJ, Wang ZY and Chen Z. (1996). *Blood*, **88**, 1052–1061.
- Chen GQ, Shi XG, Tang W, Xiong SM, Zhu J, Cai X, Han ZG, Ni JH, Shi GY, Jia PM, Liu MM, He KL, Niu C, Ma J, Zhang P, Zhang TD, Paul P, Naoe T, Kitamura K, Miller W, Waxman S, Wang ZY, de Thé H, Chen SJ and Chen Z. (1997). *Blood*, **89**, 3345–3353.
- Chen YC, Lin-Shiau SY and Lin JK. (1998). *J. Cell. Physiol.*, **177**, 324–333.
- Chen Z, Chen GQ, Shen ZX, Chen SJ and Wang ZY. (2001). *Semin. Hematol.*, **38**, 26–36.
- Cheng GX, Zhu XH, Men XQ, Wang L, Huang QH, Jin XL, Xiong SM, Zhu J, Guo WM, Chen JQ, Xu SF, So E, Chan LC, Waxman S, Zelent A, Chen GQ, Dong S, Liu JX and Chen SJ. (1999). *Proc. Natl. Acad. Sci. USA*, **96**, 6218–6223.
- Cohen MH, Hirschfeld S, Susan FH, Ibrahim A, Johnson JR, O'Leary JJ, White RM, Williams GA and Pazdur R. (2001). *Oncologist*, **6**, 4–11.
- Conrad ME. (1999). *N. Engl. J. Med.*, **340**, 1043–1045.
- Costantini P, Jacotot E, Decaudin D and Kroemer G. (2000a). *J. Natl. Cancer Inst.*, **92**, 1042–1053.
- Costantini P, Belzacq AS, Vieira HL, Larochette N, de Pablo MA, Zamzami N, Susin SA, Brenner C and Kroemer G. (2000b). *Oncogene*, **19**, 307–314.
- Dai J, Weinberg RS, Waxman S and Jing Y. (1999). *Blood*, **93**, 268–277.
- Deaglio S, Canella D, Baj G, Arnulfo A, Waxman S and Malavasi F. (2001). *Leukemia Res.*, **25**, 227–235.
- Galimberti S, Papineschi F, Carmignani A, Testi R, Fazzi R and Petrini M. (1999). *Bone Marrow Transplant*, **24**, 345–348.
- Gallagher RE. (1998). *N. Engl. J. Med.*, **339**, 1389–1391.
- Gianni M, Koken MH, Chelbi-Alix MK, Benoit G, Lanotte M, Chen Z and de Thé H. (1998). *Blood*, **91**, 4300–4310.
- Gianni M and de Thé H. (1999). *Leukemia*, **13**, 739–749.
- Huang C, Ma WY, Li J and Dong Z. (1999). *Cancer Res.*, **59**, 3053–3058.
- Huang CH, Chen WJ, Wu CC, Chen YC and Lee YT. (1999). *Pacing Clin. Electrophysiol.*, **22**, 965–967.
- Huang S, Huang CF and Lee T. (2000). *Biochem. Pharmacol.*, **60**, 771–780.
- Huang SY, Chang CS, Tang JL, Tien HF, Kuo TL, Huang SF, Yao YT, Chou WC, Chung CY, Wang CH, Shen MC and Chen YC. (1998). *Br. J. Haematol.*, **103**, 1092–1095.
- Huff J, Chan P and Nyska A. (2000). *Toxicol. Sci.*, **55**, 17–23.
- Jiang XH, Chun-Yu Wong B, Yuen ST, Jiang SH, Cho CH, Lai KC, Lin MC, Kung HF and Lam SK. (2001). *Int. J. Cancer*, **91**, 173–179.
- Jing Y, Dai J, Chalmers-Redman RM, Tatton WG and Waxman S. (1999). *Blood*, **94**, 2102–2111.
- Jing Y, Wang L, Xia L, Chen GQ, Chen Z, Miller WH and Waxman S. (2001). *Blood*, **97**, 264–269.
- Kinjo K, Kizaki M, Muto A, Fukuchi Y, Umezawa A, Yamato K, Nishihara T, Hata J, Ito M, Ueyama Y and Ikeda Y. (2000). *Leukemia*, **14**, 431–438.
- Kitamura K, Yoshida H, Ohno R and Naoe T. (1997). *Int. J. Hematol.*, **65**, 179–185.
- Kitamura K, Minami Y, Yamamoto K, Akao Y, Kiyoi H, Saito H and Naoe T. (2000a). *Leukemia*, **14**, 1743–1750.
- Kitamura K, Hoshi S, Koike M, Kiyoi H, Saito H and Naoe T. (2000b). *Br. J. Haematol.*, **108**, 696–702.
- Koken MH, Daniel MT, Gianni M, Zelent A, Licht J, Buzyn A, Minard P, Degos L, Varet B and de Thé H. (1999). *Oncogene*, **18**, 1113–1118.
- Kroemer G and de Thé H. (1999). *J. Natl. Cancer Inst.*, **91**, 743–745.
- Kwong and Todd D. (1997). *Blood*, **89**, 3487.
- Lallemant-Breitenbach V, Guillemain MC, Janin A, Daniel MT, Degos L, Kogan SC, Bishop JM and de Thé H. (1999). *J. Exp. Med.*, **189**, 1043–1052.
- Larochette N, Decaudin D, Jacotot E, Brenner C, Marzo I, Susin SA, Zamzami N, Xie Z, Reed J and Kroemer G. (1999). *Exp. Cell Res.*, **249**, 413–421.
- Li YM and Broome JD. (1999). *Cancer Res.*, **59**, 776–780.
- Lin CP, Huang MJ, Chang IY, Lin WY and Sheu YT. (2000a). *Leuk. Lymphoma*, **38**, 195–198.
- Lin CP, Huang MJ, Chang IY and Lin WY. (2000b). *Leuk. Lymphoma*, **38**, 191–194.
- Look AT. (1998). *J. Natl. Cancer Inst.*, **90**, 86–88.
- McCabe MJ, Singh KP, Reddy SA, Chelladurai B, Pounds JG, Reiners JJ and States JC. (2000). *J. Pharmacol. Exp. Ther.*, **295**, 724–733.
- Melnick A and Licht JD. (2000). *Blood*, **93**, 3167–3215.
- Mervis J. (1996). *Science*, **273**, 578.
- Muller S, Matunis MJ and Dejean A. (1998). *EMBO J.*, **17**, 61–70.
- Niu C, Yan H, Yu T, Sun HP, Liu JX, Li XS, Wu W, Zhang FQ, Chen Y, Zhou L, Li JM, Zeng XY, Yang RR, Yuan MM, Ren MY, Gu FY, Cao Q, Gu BW, Su XY, Chen GQ, Xiong SM, Zhang TD, Waxman S, Wang ZY and Chen SJ. (1999). *Blood*, **94**, 3315–3324.
- Ohnishi K, Yoshida H, Shigeno K, Nakamura S, Fujisawa S, Naito K, Shinjo K, Fujita Y, Matsui H, Takeshita A, Sugiyama S, Satoh H, Terada H and Ohno R. (2000). *Ann. Intern. Med.*, **133**, 881–885.
- Ohsawa M, Koyama T, Shibakura M, Kamei S and Hirose S. (2000). *Leukemia*, **14**, 941–943.
- Rego EM, He LZ, Warrell RP, Wang ZG and Pandolfi PP. (2000). *Proc. Natl. Acad. Sci. USA*, **97**, 10173–10178.
- Roboz GJ, Dias S, Lam G, Lane WJ, Soignet SL, Warrell RP and Rafii S. (2000). *Blood*, **96**, 1523–1530.
- Seol JG, Rark WH, Kim ES, Junk CW, Hyun JM, Lee YY and Kim Bk. (2001). *Int. J. Oncol.*, **18**, 249–255.
- Shao W, Fanelli M, Ferrara FF, Riccioni R, Rosenauer A, Davison K, Lamph WW, Waxman S, Pelicci PG, Lo Coco F, Avvisati G, Testa U, Peschle C, Gambacorti-Passerini C, Nervi C and Miller WH. (1998). *J. Natl. Cancer Inst.*, **90**, 124–133.
- Shen Y, Shen ZX, Yan H, Chen J, Zeng XY, Li JM, Li XS, Wu W, Xiong SM, Zhao WL, Tang W, Wu F, Liu YF, Niu C, Wang ZY, Chen SJ and Chen Z. (2001). *Leukemia*, **15**, 735–741.
- Shen ZX, Chen GQ, Ni JH, Li XS, Xiong SM, Qiu QY, Zhu J, Tang W, Sun GL, Yang KQ, Chen Y, Zhou L, Fang ZW, Wang YT, Ma J, Zhang P, Zhang TD, Chen SJ, Chen Z and Wang ZY. (1997). *Blood*, **89**, 3354–3360.
- Shen ZY, Tan LJ, Cai WJ, Shen J, Chen C, Tang XM and Zheng MH. (1999). *Int. J. Mol. Med.*, **4**, 33–37.
- Slack JL and Rusiniak ME. (2000). *Ann. Hematol.*, **79**, 227–238.

- Soignet SL, Maslak P, Wang ZG, Jhanwar S, Calleja E, Dardashti LJ, Corso D, DeBlasio A, Gabrilove J, Scheinberg DA, Pandolfi PP and Warrell RP. (1998). *N. Engl. J. Med.*, **339**, 1341–1348.
- Sternsdorf T, Puccetti E, Jensen K, Hoelzer D, Will H, Ottmann OG and Ruthardt M. (1999). *Mol. Cell. Biol.*, **19**, 5170–5178.
- Sun HD, Ma L, Hu XC and Zhang TD. (1992). *Chin. J. Integrat. Chin. Trad. Med. Western Med.*, **12**, 170–171.
- Tallman MS. (1999). *Semin. Thromb. Hemost.*, **25**, 209–215.
- Tamm I, Paternostro G and Zapata JM. (1998). *N. Engl. J. Med.*, **340**, 1043.
- Uslu R, Sanli UA, Sezgin C, Karabulut B, Terzioglu E, Omay SB and Goker E. (2000). *Clin. Cancer Res.*, **6**, 4957–4964.
- Wang ZG, Rivi R, Delva L, Konig A, Scheinberg DA, Gambacorti-Passerini C, Gabrilove JL, Warrell RP and Pandolfi PP. (1998). *Blood*, **92**, 1497–1504.
- Wang ZY and Chen Z. (2000). *Lancet Oncol.*, **1**, 101–106.
- Warrell RP. (1999). *Haematologica*, **84**, 75–77.
- Zhang P, Wang SY and Hu XH. (1996). *Chin. J. Hematol.*, **17**, 58–60.
- Zhang T, Westervelt P and Hess JL. (2000). *Mod. Pathol.*, **13**, 954–961.
- Zhang W, Ohnishi K, Shigeno K, Fujisawa S, Naito K, Nakamura S, Takeshita K, Takeshita A and Ohno R. (1998). *Leukemia*, **12**, 1383–1391.
- Zhu J, Guo WM, Yao YY, Zhao WL, Pan L, Cai X, Ju B, Sun GL, Wang HL, Chen SJ, Chen GQ, Caen J, Chen Z and Wang ZY. (1999). *Leukemia*, **13**, 1062–1070.
- Zhu J, Koken MH, Quignon F, Chelbi-Alix MK, Degos L, Wang ZY, Chen Z and de Thé H. (1997). *Proc. Natl. Acad. Sci. USA*, **94**, 3978–3983.
- Zhu XH, Shen YL, Jing YK, Cai X, Jia PM, Huang Y, Tang W, Shi Sun YP, Dai J, Wang ZY, Chen SJ, Zhang TD, Waxman S, Chen Z and GQ. (1999). *J. Natl. Cancer Inst.*, **91**, 772–778.