hill at 1725 ft., registered 0.0025-0.05 in. in six hours during which periods dense *Nebelreissen* cloaked the hilltop, *no actual rainfall occurring the while*. A control gauge placed well away from tall vegetation during these self-same periods registered not the slightest precipitation.

(2) The taller the vegetation the greater the precipitation induced: occasional observations at the base of a row of 100 ft.-115 ft. high *Eucalyptus globulus* growing on the north side of the research station hill indicate that the catch in six hours of dense *Nebelreissen*—without any actual rainfall—may range from 0.0025 to 0.15 in. The writer, while working at the base of these in misty weather, has had to wear a rainproof, so heavy has been the fall of water from the foliage, whereas, when the wind is of low velocity, he has been able to leave papers lying 20-30 yards away from the trees without the records becoming wet. During wind of high velocity it has been noted that an interrupted fall of moisture occurs to a distance of about 40 yards in the direction of the wind.

A continuous series of observations during misty weather has not been possible, but it seems clear that an increased fall of from 10 in. to 15 in. per annum occurs under the Eucalypts as the result of condensation of the hydrometeors. In mountain macchia and mountain forest on the south-west, south, and south-east slopes of the Outeniqua range and of its highest foothills—sites mist-clad on the aggregate a quarter to a half of the year—the increased fall as the result of condensation of the mists must be very much greater.

JOHN PHILLIPS.

Forest Research Station,
Deepwells, Knysna, S.A.,
Dec. 5, 1927.

Active Nitrogen.

IN the current issue of the *Journal of the American Chemical Society*, Dr. Bernard Lewis criticises certain aspects of the work upon active nitrogen which has recently been published from the Laboratory of Physical Chemistry, Cambridge. He states, *inter alia*, that "the authors (Willey and Rideal) neglect to consider many reactions initiated by active nitrogen which require more energy than two volts"; in spite of this somewhat sweeping statement which, if upheld, would stultify a good deal of the work under discussion, he does not quote or even hint at a single reaction which can be held to justify his contention. His next sentence, "The varied phenomena caused by active nitrogen make it evident that 2-volt level nitrogen molecules do not issue from the discharge," is equally illustrative of its author's tendency to state opinions as facts.

It is difficult to understand our being at variance over the obtaining of ammonia from active hydrogen and inert nitrogen. In the experiments at Cambridge, condenser discharges (identical with those employed in obtaining active nitrogen) were employed and careful precautions were taken against mixing of the gases in the discharges. Three concordant experiments showed that ammonia equivalent to about 0.2 per cent. active hydrogen (calculated as H_3) could be obtained if the gases after mixing were further treated with a little water vapour, admitted from another side tube, and drawn to exhaust through a condensing trap cooled by liquid air. In two runs, also, it was found that when the nitrogen was activated no ammonia was obtained, and the glow appeared to diminish slightly, a result which at the time was interpreted as being due to the decomposition by the active nitrogen of ammonia first

formed from the inert nitrogen and active hydrogen; the quantity of active nitrogen present (1 per cent.) rendered this quite probable. The experiments were carried out only to test Wendt and Landauer's claims as to the properties of active hydrogen, and were part of the attempts to activate this gas by second-type collisions with active nitrogen as described in the second paper.

It is a pity that Lewis did not make a quantitative study, however short, of his ammonia synthesis from active nitrogen and active hydrogen. Lord Rayleigh has shown (*Proc. Roy. Soc.*, **85**, 219; 1911), and Dr. Rideal and the author have confirmed his observation, that ammonia extinguishes the glow of active nitrogen and chemical reaction appears to be traceable. This fact would appear to constitute a grave objection to Lewis's theory of the formation of ammonia from atomic nitrogen and atomic hydrogen, namely:

Spectroscopic evidence, taken in conjunction with the decay of the afterglow being bimolecular with respect to the glow-producing system, leads to the view that the first step at any rate in the development of the luminosity is the recombination of nitrogen atoms, which are the primary reacting system according to his scheme. If, then, they are to take part in the ammonia synthesis, considerable changes—probably extinction—of the glow might be anticipated; conversely, if ammonia were produced in any quantity by any other mechanism, it should also affect the glow, but no mention of such a phenomenon occurs in his paper. The matter will later be discussed more fully in another place.

Recent studies of the decay of the after-glow have, it is believed, led to important clues as to the origin and nature of active nitrogen. It appears that between 1 mm. and 10 mm. pressure the decay process is very complex, although bimolecular with respect to the active nitrogen. Moreover, it appears very likely that the luminosity and the chemical activity are steps in an involved deactivation process; the first of these stages appears to consist of a ternary collision between two atoms and a molecule, as would be expected from the negative temperature coefficient of the decay process as observed long ago by Lord Rayleigh (*Proc. Roy. Soc.*, **A**, **86**, 262; 1912).

A full account of these investigations will shortly be published.

E. J. B. WILLEY.

Laboratory of Physical Chemistry,
Cambridge, Feb. 13.

Units of Energy.

DR. RUSSELL's suggestion (*NATURE*, Feb. 4, p. 170) that the kilowatt-hour is the best unit of energy for practical purposes, ought to be popular with the gas companies. At present, by Act of Parliament, we have to purchase our energy from the public supply companies by two different measures, and the cheaper purveyor does not get full credit for his cheapness. The unit for electricity is of course the kilowatt-hour, 3.6×10^6 joules; the unit for gas energy is the therm—the energy required to heat 1000 lb. of water through $100^\circ F.$, and this is 105×10^6 joules. Thus the therm is approximately 29 kilowatt-hours. In other words, when the price of gas per therm is the same as the price of electricity per kilowatt-hour, electrical energy is 29 times as dear as gas energy.

As to the scientific aspect of the question of units, it would be useful if someone with missionary zeal could provide handy reference tables bringing in all