already reported that the characteristic secondary electron emission inflections from a crystal surface are different from those of ordinary metals between 0 and 40 volts. It is proposed to repeat the soft X-ray experiments with other crystals, as also to investigate the secondary electron emission from crystals further at higher potentials. My sincere thanks are due to Prof. Richardson for his kind help and encouragement, and to Dr. H. H. Potter for the crystal of nickel.

S. RAMACHANDRA RAO.

Wheatstone Laboratory,
King's College, London, W.C.2,
Jan. 29.

Incoherent Scattering.

THERE are several surprising peculiarities of the phenomena of modification of wave-length in scattering, but the most striking is the rareness with which one is able to find an infra-red line to fit even approximately the frequency shifts observed. On the other hand, there are many strong infra-red lines which have no corresponding frequency shift. Moreover, there is scarcely any direct correlation of intensity even when there is a supposed match in frequency.

The idea suggested by Smekal that a molecule may subtract from or add to an incident quantum one of its characteristic energy quanta and scatter the resultant sum or difference in a single quantum is so neat and clear that it is accepted as the explanation of the scattering experiments of Raman and Krishnan and others. The most important characteristics of such a process are those just indicated which are not realised in the experiments.

Kramer's and Heisenberg's correspondence principle treatment is much more successful in accounting for the facts, and Schrödinger's wave mechanics gives an almost identical result, formally, which is on closer inspection in even better accord with experiment. The wave theory for the scattering of light of frequency ν by a system excited in two of its characteristic states—let us call them k and l —leads to terms incoherent with the incident radiation of frequencies $\nu_{kl} \pm \nu$ and with intensities proportional to the square of the quantities

$$A_{kn}A_{ln}\left[\frac{1}{\nu_{kn} \pm \nu} + \frac{1}{\nu_{ln} \mp \nu}\right](\nu_{kl} \pm \nu)^2.$$

The upper signs or the lower signs are used throughout. The A_{kn} , for example, is the matrix element $\int \psi_k \psi_n dx$, which describes the intensity of the transition between states k and n , giving out radiation of

frequency $\nu_{kn} = \nu_k - \nu_n = \frac{E_k - E_n}{h}$.

We see that the modified frequencies $\nu_{kl} \pm \nu$ differ from the incident ν , not by absorption frequencies (although these may also appear) but by differences between these. That is to say, only when two allowed transitions (i.e. A_{kn} and A_{ln} different from zero) have a common level (n) is there any intensity in the scattered frequencies $\nu_{kl} \pm \nu = \pm(\nu_{kn} - \nu_{ln}) \pm \nu$.

Examination of the data shows that the frequency shifts in scattered light can be interpreted in this way. The case of carbon tetrachloride is not the most striking, but it is perhaps the simplest which has so far been tried. Only two out of five frequency shifts coincide with known infra-red lines. These two correspond perhaps to the weakest shifted lines, while the infra-red lines are the strongest. The intensity formula indicates that the strongest frequency shifts should correspond to the frequency differences between the strongest fundamentals and the other fundamentals

which end on the same level. This rule suggested the energy diagram (Fig. 1). It is believed to contain elements of reality, but should not be considered as a final and complete picture. The numbers along the full vertical lines are wave numbers of the infra-red lines due to transition indicated. The numbers by the dotted lines are the differences between levels joined, and indicate frequency shifts to be expected in scattered light. The one of 27 wave numbers would be too close to the exciting line to be observed. The others were obtained from grating and prism plates, and are probably correct to a couple of wave numbers.

The test of the theory is to compare the wave-lengths calculated from this diagram with those observed. The infra-red data are from J. Lecomte, "Le Spectre infrarouge," p. 213, and the references there given. The table gives the wave-lengths which are directly connected with the frequency shifts. All the other known lines are faint, and with one exception

FIG. 1.

of much shorter and more uncertain wave-length. The exception is the weak line observed by Marvin at 15μ . Its interpretation is not yet definite. The others are explicable as overtones or combinations of the levels shown. In fact, just those combinations appear which, according to the wave theory, indicate the possibility of the modified lines actually observed.

TABLE.

Calc. (μ)	13.14	12.69	10.24	9.97	9.30	9.07	8.20	8.02	6.57	6.46
Obs. (μ)	13.10	12.72	10.23	9.96	9.3	9.07	8.26	8.02	6.57	6.45

The agreement is in every case well within the experimental latitude. The significance and power of the scattering experiments in unravelling infra-red spectra is very much greater than if they merely checked infra-red measurements. With their help and with more precise infra-red data it may be hoped that the vibrations of complex molecules will be interpreted. A beginning will be made in a fuller account of this work.

R. M. LANGER
(National Research Fellow).

Bureau of Standards,
Washington, D.C.,
Jan. 4.