That active nitrogen is a molecule of nitrogen in this metastable condition is further supported by some recent experiments which we have carried out on the life of active nitrogen. This can be varied within wide and indefinite limits simply by the regulation of pressure, everything else remaining constant. For a short life of the order of 0.1 sec. the experiment is best performed by drawing our nitrogen at about 7 to 8 mm. pressure from a region of condensed discharge. For very low pressures, say 0.03 mm. of mercury, the active nitrogen is formed with an electrodeless discharge, and its life may be abnormally extended to several minutes. It has been clearly observed by us that for any given specimen of nitrogen the life of active nitrogen increases continuously and regularly with the decrease of pressure. This is a strong evidence for the presence of metastable molecules.

In another series of experiments we have produced the infra-red lines of nitrogen belonging to the electronic configurations $2L_2M_1 \leftarrow 2L_2M_2$ by exciting first nitrogen and then active nitrogen with uncondensed discharge under exactly identical conditions. No change in the relative intensity of lines was observed, which points to the conclusion that there is no appreciable density of atoms present in active nitrogen.

P. K. KICHLU.
S. BASU.

Department of Physics,
Science College, Patna, Mar. 27.

DR. KICHLU and Mr. Basu seem to have overlooked an early investigation (*Proc. Roy. Soc., A*, vol. 86, p. 264), in which it is shown that a given sample of active nitrogen, made active at a low density by the electrodeless discharge, can, after intervals up to several minutes, be made momentarily very bright by compression.

This experiment seems to cover what Dr. Kichlu and Mr. Basu have observed, with the additional point that compression causes the active gas to give up its energy rapidly *in the form of light* (α bands).

This seems clearly to prove that collisions of some kind are the occasion of the emission of α bands (1st positive nitrogen bands).

I am not sure if I understand the views of the authors rightly. But the level which Mulliken concludes is metastable is the lower level concerned in the emission of the α bands. It is not clear to me how the metastability of this level helps us to understand how the gas remains for a long time loaded with the energy necessary for the emission of α bands, which involve a level several volts higher.

RAYLEIGH.

Terling Place,
Chelmsford, Essex, April 24.

Properties of the Terms of the Helium Molecule.

If in a diatomic molecule the influence of the internuclear axis on the valence electron is strong compared with the influence of the nuclear rotation (case I.), the component σ , along that axis of the vector l representing the moment of momentum of the electron, is, as was shown by Hund, a whole multiple of $h/2\pi$, and the rotational energy is, apart from a constant, proportional to $j(j+1)$ (j =total moment of momentum of the molecule). If, on the other hand, the influence of the rotation is predominant (case II.), l is quantised with respect to the axis of rotation, and if its component along this axis is ρ , the rotational energy is proportional to $(j-\rho)(j-\rho+1)$. The energy in the intermediate case

is a complicated function of j , and has been calculated approximately for simple cases by Hill and van Vleck (*Phys. Rev.*, 32, p. 250; 1928).

Case I. is realised in most molecules. The electronic spin usually complicates the problem. In the helium band spectrum we can observe, as already shown in a qualitative way by Weizel (*Zeit. f. Phys.*, 52, p. 175; 1928), all the different stages of transition between the cases I. and II. The terms which are produced by the different orientations of orbits with $l=2$ (δ -complex) are of special interest. The bands which originate from a combination of this complex with the $2p$ -level show a very anomalous behaviour both with respect to the position of the lines and to their intensities. All their properties can, however, be understood, if one follows the transition from case I. to case II. The connexion between the theoretical and empirical term symbols and the values of σ and ρ is given below.

Term	Case I.	$\pi\Sigma$	$\pi\Pi_a$	$\pi\Pi_b$	$\delta\Sigma_a$	$\delta\Pi_b$	$\delta\Pi_a$	$\delta\Delta_b$	$\delta\Delta_a$
Case II.	π	π_0	π_1	δ_2	δ_1	δ_0	δ_1	δ_2	
Empirical designation	σ	p_a	p_b	z	x	y	d_b	d_a	
σ	0	1	1	0	1	1	2	2	
ρ	1	0	-1	2	1	0	-1	-2	

The connexion of the Greek letters used in designating the terms with the values of l and σ is evident (Mulliken uses S, P, D instead of Σ, Π, Δ). In the He_2 -molecule only the terms which are antisymmetric in the nuclei are present. They can only have odd values of $j-\rho$. Therefore, also in case I., j is odd for even values of ρ (index a) and even for odd ρ (index b).

In the 3δ -complex, case I. is realised for small values of j . This means we must use the σ -classification and have the normal type of transitions studied especially by Mulliken. We have here $\Sigma \rightarrow \Pi, P-, Q-,$ and R -branches, $\Pi \rightarrow \Pi$ only P - and R -branches, and $\Delta \rightarrow \Pi, P-, Q-,$ and R -branches. That is in exact agreement with the observations. For larger values of j the coupling of l with the internuclear axis becomes looser, which is shown by a shift in the energy levels and the appearance of Q -branches in the $\Pi \rightarrow \Pi$ band. At the same time the P -branch of the $\Sigma \rightarrow \Pi$ band disappears and the Q -branch becomes much weaker. The behaviour of the separate energy levels is represented by formulæ obtained in the same way as those of Hill and van Vleck.

The $2p-4\delta$ complex, which is also completely known, shows that the 4δ -terms are also for small values of j in the transition stage between cases I. and II. Whereas the energy of the separate terms becomes a complicated function of j , the theory shows that the mean values of δ_1 and δ_{-1} and those of $\delta_2, \delta_0,$ and δ_{-2} behave like the energy of a $\sigma\Sigma$ -term which can always be represented by a simple quadratic expression in j . That is in excellent agreement with the observations, and the molecular constants can easily be calculated in this way.

For the 5δ and 6δ complexes the observational data are not yet entirely complete. But the existing data show that stage II. is reached already for very small values of j . The anomalous energy values have disappeared. The nuclear moment of momentum is again an integration constant and ought therefore only to change 0 and ± 1 in a transition. That means that for the combination of a δ_ρ -term with the $2p$ level which is in stage I., we get the following branches:

Initial term	δ_2	δ_1	δ_0	δ_{-1}	δ_{-2}
Branches	R, Q	R, Q, P	Q, P	Q, P	P
Appearance	P	Q, P	R, Q, P	R, Q	R

So far as the data permit this to be tested, it was found to be in agreement with the facts.

The transition stage of the π -terms is analogous, though much simpler, owing to the fact that there are only three of them, and because the one with $\sigma=1$