Letters to the Editor.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, nor to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Solutions and Heat Engines.

MAY I add a word to the discussion on osmotics? As regards the osmotic pressures of mixed gases I would point out that the reviewer's case (2) (NATURE, April 13, p. 569), where two atmospheres of nitrogen are inside the chamber and one atmosphere of hydrogen is on each side, is not in osmotic equilibrium if there be any interaction between the molecules of nitrogen and hydrogen: the equilibrium will obtain, so it seems to me, only when the partial pressure of the hydrogen equals its pressure outside, and we have, as yet, no means of calculating this effect.

With liquid solutions a little consideration will convince one that there are a multitude of formulæ, functions of the concentration, which will fit the facts for dilute solutions. Among these may I direct attention to one which seems promising?

If the observations of Berkeley, Hartley, and Burton (*Phil. Trans. R.S.*, vol. 218) on the osmotic pressures of cane sugar and α methyl glucoside are tabulated, as below, against the weight concentration multiplied by the density of the solution, squared, the figures agree to about 5 per cent.

CANE SUGAR.									
At 0° C.					At 30° C.				
c_2/c_1	O.P.	Ratio of O.P.'s	d^2	Ratio of $d^2 \times c_2/c_1$	O.P.	Ratio of O.P.'s	d^2	Ratio of $d^2 \times c_2/c_1$	
0-3400	Atm.	1	1	1	Atm.	1	1	1	
0-5650	43-91	1.27	1.357	1.1	26-82	1	1-216	1	
0-8120	67-43	1.54	1-468	1-56	47-25	1-76	1-333	1-83	
1-1200	100-53	2-29	1-580	2-30	72-59	2-71	1-439	2-82	
1-4100	134-86	3-07	1-667	3-08	107-55	4-01	1-548	4-18	
1-8300	186-86	4-26	1-768	4-21	143-33	5-34	1-632	5-56	
2-1750	230-70	5-25	1-835	5-20	198-89	7-42	1-730	7-64	
2-4300	264-46	6-02	1-877	5-93	249-16	9-29	1-796	9-47	
					..	..	..	..	
α METHYL GLUCOSIDE.									
c_2/c_1	O.P.	Ratio of O.P.'s	d^2	Ratio of $d^2 \times c_2/c_1$	O.P.	Ratio of O.P.'s	d^2	Ratio of $d^2 \times c_2/c_1$	
0-3500	Atm.	1	1	1	Atm.	1	1	1	
0-4500	48-29	1	1-199	1	49-42	1	1-179	1	
0-5500	64-22	1-33	1-245	1-34	65-14	1-32	1-222	1-34	
0-6500	80-50	1-67	1-287	1-68	81-73	1-65	1-262	1-68	
0-6400	96-17	1-99	1-310	1-99	96-75	1-96	1-294	2-01	
0-7500	115-74	2-40	1-361	2-44	115-34	2-33	1-331	2-42	
0-9000	142-46	2-95	1-408	3-00	141-66	2-87	1-375	3-00	
1-0500	170-18	3-52	1-451	3-60	168-34	3-41	1-415	3-60	

NOTES.— c_2 and c_1 are the concentrations, i.e. number of grams per gram of solution; d is the density of the solution compressed to its osmotic pressure.

It is easy to see that the osmotic pressures must be a function of the density; for consider two cylinders containing different solutions and furnished at the bottom with semi-permeable membranes which just touch the surface of the solvent. If we neglect the stratification caused by the gravitational field, then, when there is equilibrium across the membrane, $P = hd$ where P is the osmotic pressure, d the density of the solution, and h its height, hence

$$\frac{P_1}{P_2} = \frac{h_1 d_1}{h_2 d_2}$$

An explanation of the weight concentration part of the formula can be put forward. Assume that the solute

takes no part in the bombardment of the membrane, that is, this bombardment is conditioned only by the solvent molecules. It will be necessary, therefore, to put a pressure on the solution to increase the speed of the solvent molecules such that the number striking the membrane per second on the solution side will be equal to the number on the pure solvent side. It is easy to see, if our solution is an ideal one (that is, there is no interaction between the two sets of molecules and therefore there is no change in volume when the liquid substances are mixed), that this pressure will be proportional to c_2 , and, if we remember we are dealing with a defect in bombardment, it will roughly be inversely proportional to c_1 . The c 's are the number of grams of solute (c_2) and solvent (c_1) in one gram of solution—and c_2/c_1 = weight concentration/100.

Obviously this explanation is but a rough approximation to actual conditions, but, if the formula applies to substances other than the sugar type of molecule, we have a rule-of-thumb means of calculating both osmotic pressures—a matter of some importance as they are just as much physical constants as the density or refractive index.

A little thought will make it evident, if we remember that we are still considering an ideal solution, that we could have put $c_2/c_1 = v_2/v_1$ (where v_2 and v_1 are the volumes of the respective components in 1 c.c. of solution), and we should have had a more consistent formula. But with the v 's¹ of the actual solutions the results are not so good—they only agree to 15 per cent; this is not to be wondered at, for we have not taken into consideration the molecular interaction nor the effect due to closeness of packing. I think, however, that these two considerations can be allowed for if we may assume that when one molecule strikes another the rebound is not instantaneous and a 'rest period' ensues; the effect of closeness of packing may turn out to be a function of the density, but I have not the means at hand for calculating this. I hope to return to the matter in another communication.

BERKELEY.

Determination of Crystal Potentials by Diffraction of High Voltage Electrons.

WHEN electrons are diffracted by a crystal cleavage face, Bragg's law, on taking account of the refractive index of the crystal for the electron waves, becomes

$$n\lambda = 2d \sin \theta \sqrt{1 + \frac{\mu^2 - 1}{\sin^2 \theta}}, \text{ or, putting } \mu = \sqrt{1 + \frac{\phi}{V}} \text{ and}$$

$\lambda = \frac{h}{mv} = \sqrt{\frac{150}{V}} \text{ A.},$ where ϕ is the inner potential of the crystal and V is the energy of the electrons in volts, we obtain

$$\sqrt{V} \sin \theta = \frac{n\sqrt{150}}{2d} \sqrt{1 - \frac{4d^2\phi}{150n^2}}. \quad (1)$$

For a spacing 4 A. the first order will thus disappear entirely for ϕ as small as 2.4 volts, whatever the value of V . This wide variation from Bragg's simple law, then, will be quite as marked for high as for low voltages, and since swift electrons are less liable to be deviated by stray fields, etc., the high voltage method ought to be the more suitable for determining ϕ . The surprisingly large effect of refractive index at these high voltages depends on the very small angles of refraction which occur, and is only strongly marked when the reflecting plane is the free surface of the crystal. This effect has been pointed out by Prof. G. P. Thomson (*Phil. Mag.*, 6, p. 939; 1928).

¹ These are derived from Porter's (*Proc. Roy. Soc.*, 1908, p. 460) definition of his s_2 and s_1 .