Diving.*

By Prof. LEONARD HILL, F.R.S.

THE new submarine escape dresses designed by Mr. R. H. Davis have been successfully used at a depth of 130 ft. at Loch Long and have since been adopted by the Admiralty. Every man in the submarine service will have to have one, just as the soldier has his gas mask. In case of accidental sinking of a submarine, each man will at once put on the dress and partly fill the breathing bag from the small oxygen cylinder which is fixed below the bag, and then breathe in and out of the bag through a cartridge of soda lime which is put inside to absorb the exhaled carbon dioxide. Then the submarine will be flooded so as to equalise the air and the water pressure, and allow a hatch to be opened through which the men will float up one by one and reach the surface; the breathing bag, acting as a balloon, will take them there. There is an escape valve provided for preventing over-distension of the bag.

In using the submarine escape dress, men must be trained to fill the bag only partially with oxygen, so as to allow expansion of the gas on coming up. There is no risk of bubbles of nitrogen forming in the body fluids through coming up quickly, even from a depth of 300 feet, because oxygen has been breathed from the time the air pressure was increased by flooding the submarine.

In case of prolonged difficulty in effecting escape from a submarine, surplus apparatus to afford fresh soda-lime cartridges and oxygen will be required. If the water has entered the submarine so that the air is under pressure, the bag should be filled with expired air two-thirds full and only enough oxygen added to keep it comfortably full—this to prevent oxygen poisoning. In this case, just before coming out, the bag must be squeezed empty and then filled with pure oxygen and this breathed for a few minutes so as to prevent bubbling off of nitrogen in the body.

The new self-contained diving dress allows a diver to get an air supply from a cylinder containing a mixture of air and oxygen, an injector device being used to draw the air in the helmet through a canister of soda lime. The diver is free from hose-pipe and air-pump, and merely carries a life-line with connexion for the telephone in his helmet. Arrangement is made so that he can detach this line and leave it fast outside while proceeding into a wreck, using a distance line to find his way out by.

The use of this apparatus is limited to a certain depth by the percentage of oxygen which it is safe to breathe. If 50 per cent oxygen is supplied in the cylinder, the diver can safely do two or three hours' work in shallow water, but he should not exceed half an hour at a depth of 100 feet. For greater depths, air with smaller percentages of oxygen must be used, and experience is required to work out the most suitable method of supply. If

the air is kept pure by absorption of carbon dioxide by means of a soda-lime cartridge, then a small oxygen cylinder will suffice to keep the percentage in the helmet at, say, 10 per cent, for deep work. The reducing valve can be set to give that amount of oxygen which the diver uses per minute, but there is a risk of too little oxygen arising through extra hard work on the part of the diver, and we must bear in mind that want of oxygen does not give warning before a man loses full use of his senses. An additional cylinder of air with 10 per cent of oxygen in it will have to be carried for altering the buoyancy of the dress as needed. Probably the ordinary method of ventilating the helmet by a hose-pipe will be used for deep work, a tank containing compressed air with 10 per cent oxygen being used as the supply. A ready method of partially deoxygenating air will have to be installed on the diving ship.

The new submersible decompression chamber, also designed by Mr. R. H. Davis, was used with great success at trials conducted by the Admiralty at Loch Long last summer. An attendant goes down in this cylindrical chamber, which is fitted with a door top and bottom, and waits to receive the diver at a depth of 66 feet. The lower door is open and water kept out of the chamber by compressed air pumped into it. The chamber is lit with an electric lamp. The shot rope of the diver passes through rings attached to the outside of the chamber. After doing his job on the bottom he ascends rapidly to the chamber, and climbing up by a small ladder thrusts his helmet up in the chamber, so far that the attendant can cast off hose-pipe and life-line and take off the helmet. The diver then comes right in and the lower door is shut. A telephone message is then sent and the chamber is raised to deck level and decompression carried out there. From the time the diver leaves the bottom, all these operations take only seven or eight minutes to carry out; the diving ship can, if need be, then proceed to harbour. This is a great improvement on the old method of the diver hanging on the shot rope in the cold and dark while decompression in stages is carried out. Moreover, when in the chamber the diver can breathe oxygen by means of a small apparatus, and by this means shorten the decompression time by at least one-half.

There are three dangers from which the diver has to be protected. First, he may be poisoned by too much carbon dioxide through insufficient ventilation. If one air-pump suffices to ventilate the helmet at one atmosphere, an extra pump is required for each atmosphere in addition, as pointed out by Dr. J. S. Haldane, so at 300 feet ten pumps are required. Many divers have been made less efficient while at work and damaged in health by insufficient ventilation. In place of so many pumps, a canister of soda lime can be arranged to purify the air in the helmet, and this is an improve-

* From the Friday evening address delivered at the Royal Institution on Feb. 7.

ment which will, I think, be adopted for deep diving. Secondly, at considerable depths the diver is exposed to high tensions of oxygen; for example, at 300 feet, to 10 atmospheres of air, which is equivalent to 2 atmospheres of oxygen; and oxygen at any tension a little greater than 70 per cent of an atmosphere acts as a poison when breathed for more than a certain time, the time becoming shorter as the tension is raised. While exposure for some days to a tension of oxygen greater than 70 per cent of an atmosphere produces pneumonia, experience has shown that the breathing of pure oxygen at a pressure of one atmosphere for a few hours, as in the use of mine-rescue apparatus, has no ill-effect.

Experiments have shown that animals can safely breathe 2 atmospheres of oxygen for 2 hours, but during this period both the use of oxygen by the body and the output of carbonic acid diminish, the body temperature drops and the animals become drowsy. If the exposure is more prolonged, or in shorter times if the oxygen tension is higher, the animals fall into a comatose condition and finally have convulsions and die. Some very important experiments were made by Bornstein at the Elbe tunnel works. He breathed oxygen at 3 atmospheres (+30 lb.) for 48 minutes, and two of the engineers (at the Elbe tunnels) for 30 minutes without harm. He considered periods such as these set the limits to safety.

In diving experiments in the United States of America in connexion with the practice of an American submarine escape apparatus, oxygen has been used without detriment for short periods up to a pressure of about twelve atmospheres, equivalent to 357 feet of water.

Argyll Campbell has shown that normally when breathing air the tension of oxygen in the tissues equals 37 mm. of mercury. The blood in the veins coming from the tissues is normally about 75 per cent saturated with oxygen and carries $2\frac{1}{2}$ to 3 volumes per cent more carbon dioxide than the arterial blood, which is 95 per cent saturated with oxygen. Under high oxygen tensions in the inspired air, the tissue oxygen tensions rise, for example, from 84 mm. to 430 mm. of mercury, and the venous blood, being almost fully saturated with oxygen, can carry very little more carbon dioxide than arterial blood. Moreover, when the tissues are in danger of oxygen poisoning, the circulation may be reduced. Thus carbon dioxide is held back in the tissues and the tension may rise to poisonous heights, as was found to be the case by Argyll Campbell.

It is clear, then, that for deep diving the ventilation of the helmet must be kept adequate enough to prevent even a small rise of carbon dioxide in the air breathed. It must be kept in mind that, at 10 atmospheres, 0.1 per cent of carbon dioxide in the air breathed produces a tension of carbon dioxide of 1 per cent of an atmosphere, and 1 per cent produces a tension of 10 per cent. The poisonous effect depends on the tension, and 10 per cent is enough to anaesthetise a man. If there is difficulty in carbon dioxide being carried from the

tissues owing to high oxygen tension, this difficulty must not be accentuated in the least degree by ill-ventilation of the helmet, otherwise the diver may become drowsy, and even comatose.

To avoid any poisonous effect of oxygen for deep dives the percentage in the air supplied to the helmet can be halved. Mr. R. H. Davis has contrived apparatus for effecting this, that is, for supplying less oxygen during compression, and more during decompression; for diving to greater depths, air with still less oxygen in it will have to be used. As pure oxygen is going to be used in the submersible decompression chamber, it is obviously best to prevent any excess of oxygen while at work on the bottom. At 10 atmospheres, 2 per cent of oxygen in the air would give a diver the natural amount of oxygen. Suppose, for safety sake, that 10 per cent of oxygen is used at 10 atmospheres, the nitrogen will be increased by one atmosphere, and this will have to be allowed for in reckoning the decompression period.

The effect of breathing oxygen in washing out nitrogen can be shown. If diuresis is established by drinking two or three pints of water, the secretion of the kidneys can be used as a measure of the nitrogen dissolved in the blood. The bladder can be emptied every few minutes and samples of the urine collected with precaution to exclude contact with air. The dissolved gas can be extracted by a vacuum pump, and the amount of nitrogen estimated. At ordinary atmospheric pressure, there is approximately 1 per cent of dissolved nitrogen gas, at 2 atmospheres 2 per cent, at 3 atmospheres 3 per cent, and so on. The effect of breathing oxygen for nine minutes during exposure of the body to a pressure of 3 atmospheres (+30 lb.) was to lower the percentage of nitrogen in the urine then secreted to 2.1, and in the next six minutes to 0.9. Urine secreted during decompression after being at +30 lb. pressure, this time with no breathing of oxygen, contained 2.86 per cent of dissolved nitrogen, more than three times as much. It is clear, then, that the breathing of oxygen quickly washes out nitrogen dissolved in the blood and tissues of the kidneys.

The washing out of nitrogen from the body by breathing oxygen has been shown in another way, by Argyll Campbell and myself. After taking three or four deep inspirations from a bag full of oxygen and expiring into the air so as to wash most of the nitrogen out of the lungs, a deep expiration is made and a sample of alveolar air collected, the subject then breathing through a soda-lime cartridge in and out of a spirometer containing a few litres of pure oxygen. A sample is drawn from the spirometer for analysis at 3 min. and 6 min., in each case at the end of a deep expiration, and the volume of gas in the spirometer being on each occasion measured. At the end of the deep expiration a sample of alveolar air is also collected from the tube leading to the spirometer. The volume of the residual air in the lungs can be calculated, and so, too, the use of oxygen in metabolism by the subject during the period.

It is estimated that there are about 960 c.c. of

nitrogen dissolved in the body of a man under ordinary atmospheric pressure, say 30 c.c. in the blood, 510 c.c. in the fat, which dissolves about five times as much water, and 420 c.c. in the other tissues. Our experiments show that about 30 c.c. of nitrogen per minute are washed out in the first few minutes from a man at rest, and about 50 c.c. from one doing stepping exercise. It seems clear, then, that by breathing oxygen, and exercise, the washing out of half of the nitrogen, probably that amount which causes bubbling and severe symptoms on decompression, takes place in about 10 minutes. It is safe for a diver to breathe oxygen during the time required for decompression in the submersible chamber from 3 atmospheres absolute pressure (66 feet approx. depth) if the oxygen breathing allows the times of the Admiralty table set for the safe decompression of divers to be halved.

Further research on animals is now being carried out to settle the point whether the formation of bubbles can be prevented by plus two atmospheres of pressure (66 ft. depth) when a diver is quickly decompressed from 300 ft. to that pressure after a prolonged stay on the bottom. Possibly a stage will have to be given at 99 ft. If the diver is supplied with, say, 5 per cent of oxygen at the great depths, he can safely begin to increase the oxygen in his helmet as he climbs up from the bottom, using a small cylinder of pure oxygen for this purpose, and then arriving at the submersible decompression chamber be ready to breathe pure oxygen at plus two atmospheres and during the decompression from that pressure. When a bottle of champagne is opened in a chamber at plus two atmospheres, it appears flat, as this pressure stops the formation of bubbles and the gas escapes quietly. In the trials at Loch Long, divers came up quickly from 300 ft. to 66 ft. and suffered no harm. It is important that the deep sea divers go down fasting with the least gas in his alimentary canal. Gas formed therein mechanically obstructs the circulation by expanding during decompression and enhances the danger of bubbles forming in the blood. Out of twenty-four well-fed guinea-pigs compressed for 1 hour to plus 100 lb. and decompressed in five minutes, nineteen died; out of twenty-four fasting guinea-pigs only eleven died. None died when oxygen was given during the period of decompression.

The U.S. Navy is experimenting with helium and oxygen; helium has less solubility and greater diffusivity than nitrogen, and good results are reported. We have no available helium.

There are certain other dangers to which the diver is exposed, as were evidenced by the salving of the *S 51* submarine by the U.S. Navy. The inlet valve may become blocked by ice due to freezing of water condensed out of the air pumped down the hose pipe in wintry weather when the salt water at the bottom of the sea is below freezing point. One diver was drawn up just in time, half suffocated; another was far inside the submarine when this happened, but managed to knock his valve free of ice by striking the outside with a spanner. The

air-supply after that had to be freed from water vapour for such wintry diving. It is astonishing that the divers can work with their hands exposed to such cold water even when gloved. A thick, knitted wool glove containing air in its meshes with a rubber glove outside is, of course, the best protection against cold.

Another danger is the outlet valve becoming jammed through sand getting into it when lying down and tunnelling as those American divers did through clay, using a fire hose and high pressure water to cut out the clay. If the outlet valve closes, a diver may have his dress blown out and be spread-eagled before he can shut off his air supply or open his spit cock and use that as an air outlet. If his telephonic message to cut off the air be not heard his dress will be burst. This actually happened to Eddy, a famous diver, but another diver witnessed the accident and, telephoning the news, Eddy was drawn up in time, put in the recompression chamber, and saved. To be blown up is dangerous, first, because of the rapid decompression, and secondly because of the risk of the helmet hitting the bottom of the attendant ship. If a diver be blown up to the surface, he must be at once sent down again after letting out excess of air, or have his helmet taken off and be rushed into the recompression chamber.

The courage and presence of mind of divers is shown by the story of a diver named Smith who, when tunnelling out the clay and getting near the keel of the submarine *S 51*, had the clay walls of the tunnel fall in upon him. He telephoned for help, but managed to turn round the nozzle between his legs and force out the clay and so escape; then, telephoning he was all right, he went back again into the tunnel and completed the job, and this down in the dark at a depth of 130 feet.

The oxyhydrogen flame has been adapted for use under water, and large thicknesses of metal were quickly burnt through by this means in the salving of *S 51*, by Commander Ellsburgh.

The 'iron man' designed in Germany and used in wrecks by Italian divers is an observation chamber with jointed limbs which allow a certain amount of awkward movement. The joints have ball bearings. The diver, shut inside, is at ordinary atmospheric pressure and breathes oxygen, having the exhaled carbon dioxide absorbed by soda lime. He can use the oxygen supply also to alter the buoyancy of the iron man. There are glass discs for observation, and in dark water a powerful electric lamp has to be lowered down for him to see by. At a depth of 100 feet the sunlight appears as moonlight, and at 200 feet as starlight. Tools are affixed to the end of the 'iron man's' arm, and the diver works these by scissor-like handles inside. The diver in the 'iron man' cannot use his sense of touch, and prolonged training and good lighting are necessary for effecting any skilled work. After a dozen trials in it, one diver failed to shackle a rope on to a buoy. The diver in the 'iron man' can guide a charge of guncotton, or a grab for seizing hold of wreckage, but is not capable of very skilled work.