Grassland Research in the British Empire.

THROUGHOUT the British Empire there are vast tracts of natural pastures which produce herbage of an inferior nutritive value. A large body of evidence has been accumulated in recent years to show that this low nutritive value is to be correlated with deficiencies of inorganic constituents, such as calcium, phosphorus, and chlorine, in the herbage, a condition which is the outcome of growth on mineral-deficient soils. Animals ranging over these pastoral areas and subsisting exclusively on such mineral-deficient herbage, display a low rate of growth and production compared with what is possible on good cultivated pastures. With this decreased growth rate are usually associated lowered fertility and susceptibility to various forms of disease.

During 1926, the committee appointed by the Economic Advisory Council to consider the question of the mineral content of natural pastures recommended that practical investigations should be carried out in a suitable colony or dependency with the view of ascertaining whether the nature of such deficiencies could be determined and the diseases due to them prevented. Following this recommendation, a scheme of research was inaugurated which serves as a model of successful co-operative effort. Kenya Colony was selected as the site of the investigations, and Dr. J. B. Orr, director of the Rowett Research Institute, Aberdeen, assumed the general scientific direction. The Empire Marketing Board rendered signal help by placing generous funds at his disposal. Close and enthusiastic co-operation was established between his own staff of research assistants, some of whom worked in Aberdeen and others in Kenya, and the staff of the Agricultural Department in Kenya Colony. In addition, valuable assistance was given by settlers in the colony, many of whom provided animals and facilities for feeding experiments.

The initial outcome of these important investigations is revealed in the Report now before us, which has been drawn up by Dr. Orr in association with Mr.

A. Holm, director of agriculture, Kenya.* Though the main scheme of work is still in its early stages, the results so far obtained are of the highest economic significance and amply justify the trouble which has been taken to secure them.

It has been shown that the most striking deficiency in the natural pastures of Kenya is that of phosphorus. Indeed, in one district, Molo, the deficiency of this element is as great as that found in certain areas of South Africa, where it is the cause of disease in cattle. On the other hand, in another district, Naivasha, the pasturage, apart from deficiency in sodium and chlorine, is as rich as the herbage of good British cultivated pastures. The application of different fertilisers in the most deficient area increased the yield of herbage from 25 per cent, where common salt only was used, to as much as 400 per cent where nitrogenous and phosphatic fertilisers were employed. The provision of small amounts of the deficient minerals to animals grazing these areas was followed by an increase of 30 per cent in the yield of dairy cows, 10 per cent in the rate of growth of lambs, and about 10 per cent in the weight of the fleeces of sheep.

It was discovered that 'Nakuruitis', a local disease resembling the 'bush-sickness' of New Zealand, could be prevented by allowing animals access to a mixture of common salt and iron oxide. This method of prevention is now being applied with successful results on several farms in Kenya, oxen being kept in good condition for at least a year, whereas formerly, without the use of this simple salt lick, it was impossible to keep them free from the disease for longer than six months without sending them away for a period of grazing in another district.

H. E. WOODMAN.

* Economic Advisory Council: Committee on the Mineral Content of Natural Pastures. Sixth Report. Pp. 66. (London: H.M. Stationery Office, 1931.) 1s. net.

Medicinal Cod Liver Oil.*

ALTHOUGH the medicinal value of cod liver oil has been known for 150 years, it is only recently that it has been attributed to the presence of vitamins A and D. It is now known, moreover, that different samples of oil vary widely in their vitamin potency as well as in their more obvious chemical and physical characteristics. The principles underlying the production of a fine medicinal oil of high activity should be established so that consumers can always be certain of obtaining full benefit from its administration. The recently issued Empire Marketing Board Report by Profs. J. C. Drummond and T. P. Hilditch provides specifications for cod liver oil which are based on the examination of a large number of samples over several years, and makes recommendations for suitable methods of manufacture.

The investigators examined oils from Newfoundland, Scotland and the North Sea, Iceland, and Norway, obtained at different seasons, for vitamin potency, colour, and fatty acid content, and estimated their saponification and iodine values: oils from other fish were also compared with cod liver oil. Vitamin A was estimated both by the growth test on rats and by the antimony trichloride colour test: in a series of oils very fair agreement between the two methods was

obtained, so that the quicker and simpler colour test was used during the greater part of the research as an estimate of the vitamin A potency. Using standard conditions, the results can be expressed in Lovibond blue units; it was not found possible to express the results of assays by the biological method in units, owing to the errors of the animal test. Vitamin D was estimated by the degree of deposition of calcium at the ends of the long bones of rachitic rats as observed under X-rays and following staining with silver nitrate (the 'line test'): the assays were carried out against the standard preparation of irradiated ergosterol employed by the Pharmaceutical Society and the results expressed in terms of the Coward antirachitic unit.

The Newfoundland and Icelandic oils had the highest vitamin potency, the Norwegian the lowest, with the Scottish intermediate, but usually of the same order of activity as the latter. The vitamin A and vitamin D potencies were usually parallel.

A detailed chemical examination of the constituent fatty acids of cod liver and other oils was made: the methods and results have been published in full in the *Biochemical Journal*, by Guha, Hilditch, and Lovern (vol. 24, p. 266; 1930) and by Lovern (*ibid.*, p. 866). The general composition of the fatty acids of cod liver oil is: stearic and myristoleic acids, traces only; myristic acid, 3-7 per cent; palmitic acid,

* The Relative Values of Cod Liver Oils from Various Sources. By Prof. J. C. Drummond and Prof. T. P. Hilditch. (E.M.B. 35.) Pp. 129. (London: H.M. Stationery Office, 1930.) 1s. net.