The value of K_2' was measured at a pH of about 9, when the free carbon dioxide may be neglected, $\text{HCO}_3' + \text{CO}_3''$ equals the bound carbon dioxide as found by gas analysis and $\text{HCO}_3' + 2 \text{CO}_3''$ equals the carbon dioxide in combination with 'excess base' as found by titration.

The temperature coefficient of K_1 , for which the values found by previous workers do not agree closely, remains to be investigated. The results will in due course be published in the *Rapports et Procès-verbaux* of the International Council for the Exploration of the Sea.

KURT BUCH.

Helsingfors.

H. WATTENBERG.

Munich.

H. W. HARVEY.

Plymouth.

Hydrolytic Adsorption of Activated Charcoal.

THERE is a difference of opinion regarding the behaviour of charcoal as a gas electrode in explaining its hydrolytic adsorption (Frumkin, Schilow). It has been suggested alternatively that acidic or basic oxides formed during activation are responsible for such adsorption, and that charcoal is a negative adsorbent possessing a negative charge in water (Schilow). On the gas electrode theory, charcoal should be positively charged and adsorb acids. Negatively charged oxygen charcoal which adsorbs alkali is known. Both positively and negatively charged charcoals are possible. A relationship between the electric charge, as measured electrokinetically, and the hydrolytic adsorption of either acid or alkali has been pointed out, and it was suggested¹ that 'acid' adsorbing activated sugar charcoal would be positively charged. Negatively charged charcoal with hydrogen or weakly basic cations in the mobile sheet of the double layer should adsorb alkali and liberate acid from a neutral chloride solution. This relationship between the sign of the charge and hydrolytic adsorption has been confirmed with charcoal.

In view of the fact that it is recognised (Frumkin) that the gas electrode theory does not apply to the observation of one of us, the results obtained recently (Roychoudhury) are of interest as showing (a) that activated charcoals have generally a negative charge but become positive on sustained washing; (b) a relationship between the sign of the electric charge and acid or alkali adsorption; (c) the primary adsorption of H^+ , OH^- , Cl^- , and SO_4^{2-} ions; (d) an anomalous effect of the mass of the adsorbent on the adsorption of acids; (e) that alkaline extracts can be obtained even from activated sugar charcoal; (f) that both cations and anions are possibly formed during activation, their nature and amount depending on the treatment.

Moreover, the facts observed previously (Frumkin, Schilow), that the adsorption of acid is not determined by the activity of hydrogen ions, that both hydrogen and oxygen charcoals adsorb benzoic acid equally, that hydrogen charcoal adsorbs acids, and the part played by the ash constituents, also suggest a limit to the applicability of the gas electrode theory in this instance.

J. N. MUKHERJEE.

S. P. ROYCHOU DHURY.

Department of Chemistry,
University College of Science and Technology,
Calcutta, July 8.

¹ *J. Indian Chem. Soc.*, 2, 219; 1925.

Orientated Deposits of Copper on Bismuth.

WHEN copper is deposited electrolytically from copper sulphate solution on a cleavage surface (0001) of bismuth with extremely low current density, the minute copper crystals show distinct orientations under the microscope (Fig. 1).

The triangular shaped crystals, nearly equilateral, are of light bluish colour and their sides are parallel

FIG. 1.—Crystals of copper deposited on bismuth. Lines have been drawn marking the long axes of the copper crystals and outlining the bismuth crystal.

to the cleavages (02 $\bar{2}$ 1) on the surface (0001) of bismuth. The parallelogrammatic shaped crystals, dark in colour, have their longer axes parallel to any of the three directions.

The reason for this may be explained by the hypothesis of 'epitaxie' of L. Royer.¹ The details of this investigation will be published in the near future, in the *Memoirs of the Faculty of Engineering*, Hokkaido Imperial University, Japan.

YOSIMITI HORI,
Faculty of Engineering,
Hokkaido Imperial University,
Sapporo, Japan,
July 11.

¹ L. Royer, *Bull. soc. franç. minéral.*, 51, 7; 1928; *Compt. rend.*, 182, 326; 1926; 191, 1346; 1930, etc.

Rotation of 'Dust Devils'.

THE direction of rotation of 'dust devils' has been noted in Lower Egypt and Iraq during the past four years, and the reports received in this office indicate that in about 50 per cent of cases the direction of rotation is clockwise, and in the other 50 per cent anti-clockwise. This seems a surprising result, because if a 'dust devil' is due to the raising of dust and sand by the vigorous upward movement of air over a limited area, the rising column should assume an anti-clockwise rotation. I had an opportunity recently of watching the formation of numerous small whirls over a hot sandy space. The whirls did not attain a height of more than 2 ft., and the objects raised were leaves and feathers (not sand); the eye of the observer, being at a height of about 4 ft. 6 in., was well above the rotating column. Thirty disturbances were observed in about twenty minutes, and in no case was the rotation other than anti-clockwise (that is, cyclonic).

It occurred to me then that perhaps a discrepancy was being introduced in observers' reports, because the level of the eye in relation to the part of the column where rotation was observed is important. A clockwise rotation looked at from below is anti-clockwise