The Advancement of Botany.*

By Prof. T. G. HILL.

DOUBTLESS it is a coincidence that the year 1831 marks not only the founding of the British Association but also, so far as a date can mark, the beginning of a threefold epoch in botany: the search for a natural system of classification; the emergence of morphology as a study separate from taxonomy; and the development of physiology.

In the first, Britain gave a full share; but in morphology and physiology, on the other hand, the tale of British activity is dismal indeed: against a battalion of Continental workers we could barely range a section, and British work caused only a ripple on the botanical surface. But there is one exception: the brilliant generalisations of Charles Darwin.

MORPHOLOGY.

In 1831, morphology for the most part had relation to the phanerogams only; the structure and life history of no plant was completely known; the fields of the vascular cryptogams, the bryophytes, and the thallophytes were almost untrodden. It is not surprising, therefore, that when ontogeny came into being the trickle of investigations fast developed into a flood, and for this the greatest honour is due to Schleiden, Nägeli, von Mohl, and Hofmeister, for they showed the way and contributed the most.

A welter of facts was discovered, many of which, especially those relating to the Thallophyta, had to await the arrival of a new technique, that of cytology, before co-ordination was possible; but for the Muscineæ, the vascular cryptogams and the angiosperms, the times produced the man whose patient and detailed work, directed by a keen intellect, gave cosmos to morphological inquiry. I refer to Hofmeister, one of the greatest botanists of all time. He first traced the development of the sexual organs and of the embryo. He was the first correctly to interpret heterospory: he discovered the alternation of generations. Nowadays, alternation of generations is accepted as a matter of course, but when it is realised that Hofmeister was working it out, almost cell for cell, more than eighty years ago without the aid of the microtome and of cytological technique, as we know it, it will be appreciated how great was his work.

Although many contributed to anatomical knowledge, the great advances were associated with the names of relatively few men. Of these, Schleiden and Schwann, the zoologist, are the first to be commemorated since they laid the foundation of the scientific study of anatomy in their theory of the cell, which enabled others, especially von Mohl and Nägeli, to build a knowledge of the anatomy of plants on a sound basis. Von Mohl discriminated the tissue elements and traced the course of the vascular system. He was

the first accurately to trace the development of vessels and gave the first correct account of stomates. His observations on periderm and cambial activity are of first-rate importance.

Nägeli made several well-known anatomical investigations. His investigations on the cell wall and on the starch grain are classical: the recognition of the cell wall as a product of protoplasmic activity, growth by intussusception and the occurrence of granulose and amylose in the starch grain, need only be mentioned. The results he obtained by the use of polarised light were as important as those of Sponser and others by the use of X-rays in our own times.

In the earlier study of cytology, von Mohl and Nägeli were the pioneers. Von Mohl was the first to see cell division. Later, he studied the 'muilage' described by Schleiden in 1838; he recognised it as a distinct component of the cell and gave it its present name, protoplasm. He described the primordial utricle and realised that the streaming movements of the cytoplasm were independent of the cell sap. Further, von Mohl perceived that the matrix of the chloroplast was protoplasmic.

Nägeli more carefully observed the nucleus and appreciated the facts of cell division more truly; and in this he was the first to recognise that the nucleus is the starting-point. With the passing of time it was recognised that the protoplasm of the plant and of the animal is identical and that it is the basis of life, a conclusion to which Payen, Cohn, and Max Schulze contributed much.

The improvement of microscopical technique led to great discoveries in the details of nuclear division. In this the work of Strasburger is outstanding, and to him is due the main credit of firmly establishing that aspect of botany now called cytology. The contributions of Fleming and of Schmitz must, however, not be overlooked, nor those of Guignard, who, unfortunately, prejudiced his own work by his description of structures which had no existence.

PHYSIOLOGY.

Our knowledge of osmotic phenomena is founded on Dutrochet's studies (1827) on differential diffusion through a colloidal membrane. Graham showed (1854) that the rate of this diffusion depends, *inter alia*, on the nature of the membrane. This led to Traube's discovery of the so-called artificial cell produced by dropping a crystal of copper acetate into a solution of potassium ferrocyanide. This observation was used by Pfeffer (1877), who supported the membrane of copper ferrocyanide in the wall of a porous pot and with it made many highly important observations. Thus was born a branch of physical chemistry, and one of the first results was van't Hoff's theory of solutions. At this time de Vries also was busy on the problem and contributed a great deal to it.

* From the presidential address to Section K (Botany) of the British Association in London on Sept. 28.

Hales, Priestley, Senebier, Ingen Housz, and de Saussure were the pioneers in the study of carbon assimilation, with the result that the more obvious facts were known before 1831. Of these men, de Saussure was the greatest: he was the first to strike a balance which indicated the unity of the photosynthetic quotient, and to show that water and salts, as well as carbon dioxide, are essential for the nutrition of the green plant. His work, however, naturally did not convince the upholders of the humus theory; the problem was settled beyond dispute by Boussingault, who, in 1851, introduced the method of water culture.

Dutrochet (1837) correlated carbon assimilation with chlorophyll and was the first to use the 'bubbling method', a method adopted and improved by Sachs and used by him in quantitative experiments.

Von Mohl (1851) considered that a carbohydrate was formed from the carbon dioxide, but it was Sachs who proved that the starch grains in the leaf were produced from carbon dioxide. Sachs considered that starch was the first product of carbon assimilation; it was not until some years later (1885) that Meyer showed that there were sugar leaves as well as starch leaves and thus indicated that sugar preceded starch in the process.

Turning to the pigments associated with photosynthesis, Grew was the first to suggest that chlorophyll, using the word in its loose sense, was made up of more than one pigment, but to Stokes (1864) is due the credit of making the first real advance; he demonstrated the presence of two green and two yellow pigments. The methods of separation of the mixed pigments were in time refined and culminated in the work of Tswett (1906), who, by his chromatographic method, separated two chlorophylls and five carotinoids, of which one was carotin and the others xanthophyll.

Many years were to elapse before the chemical constitution of chlorophyll was to be elucidated, but Hoppe-Seyler was the first to make the significant discovery, in 1879, that there is a chemical relationship between the green pigment of the plant and the hæmoglobin of blood. The formaldehyde hypothesis was originated by Butlerow (1861) and developed by Baeyer (1870).

Before 1831 certain fundamental facts of respiration were known: Ingen Housz had shown that the plant absorbed oxygen and evolved carbon dioxide, and de Saussure had discovered that a continuous supply of oxygen was necessary for the life of the ordinary plant, that the more vigorous the organ the greater was the amount of oxygen consumed. Dutrochet (1837) confirmed the work of de Saussure and showed that responses to stimulation are not made in the absence of oxygen. He distinguished between the evolution of oxygen in carbon assimilation and the evolution of carbon dioxide in respiration, which facts were confused by many subsequent workers.

The reasons for our deficiencies during this epoch are patent. In the first place, the classical tradi-

tions of our old universities dominated teaching, with the result that the approach to biology was, in the main, through medicine. Other reasons lie in the facts that the period was one of exploration, expansion, and development of the Empire, which meant an inflow of great numbers of plants for identification, and the labourers were few. Finally, the systematists did not entirely favour the new movement and, if not actively antagonistic, they looked upon it with amused tolerance.

THE RENAISSANCE OF BOTANY IN BRITAIN.

The botanical renaissance in Great Britain began about 1875 and it happened in the Royal College of Science. The credit for it is due to Thiselton-Dyer, who, as Huxley's demonstrator, instituted and conducted the first laboratory classes in botany. University College, London, immediately followed. At Cambridge was Sydney Howard Vines.

In 1882 in all branches of botany there were masses of well-ascertained facts; the principles appeared to be settled and the problems formulated. Some aspects, such as anatomy, were in an advanced stage and were developing in various directions; other aspects, physiology, for example, were relatively backward.

In Germany especially, causal morphology was more dominant, in which development von Goebel played a pre-eminent part. His work and his teaching on the influence of the environment on the configuration of plants cannot be too highly appraised; he with Wiesner, Haberlandt, Stahl, Bonnier, and other pioneers stripped morphology of its formalism and revealed the plasticity of the plant.

Of the lower groups I shall say but little; one name, however, leaps to mind, that of Klebs, whose work on the conditions governing the reproduction of the green Algae is classical. In mycology, Marshall Ward laid the foundation of the modern British school. Of a later date, Blakeslee's discovery of heterothallism was of the greatest importance and led to far-reaching results.

A great contribution to our knowledge during this period was given by the palæontologists. Stur and Williamson recognised the occurrence of fern and gymnosperm characters in certain fossil plants, and with the increase of evidence a separate phylum was recognised and named Cycadofilices by Potonié in 1899. Later, in 1906, Oliver and Scott made known their observations on the seed-like structures attached to the fronds of *Lyginodendron*. Passing on to the gymnosperms, the work of Strasburger and others was continued and many gaps in our knowledge were filled; but of all these investigations the most sensational was the discovery in 1896 of the motile sperms in *Ginkgo* and *Cycas revoluta* by Hirasé and Ikeno respectively. Mention also must be made of Wieland's important work on Cycadeoidea and other fossil cycads.

This progress was accompanied by that of detailed cytology, which immediately reacted to the conception of Strasburger and Weismann that the

nucleus is the bearer of hereditary qualities (1884). The reduction in the number of chromosomes was discovered in the parasitic worm *Ascaris* by van Beneden in 1883; the next year Strasburger observed it in the Angiosperms, and this was followed by similar observations by Overton and by Farmer in other groups. Thus was initiated a period of intense cytological study.

The condition of anatomy in 1882 is shown by De Bary's "Comparative Anatomy of the Phanerogams and Ferns": the fundamental facts were well ascertained and the application of those facts determined the direction of increasing knowledge. In Germany the trend, initiated by Radlkofer, was taxonomic, and Solereder took the leading part; in Austria, Haberlandt developed physiological anatomy, whilst in France the drifts were various. The study of purely descriptive anatomy was continued by Hovelacque, for example; Sauvageau, Costantin, and others studied structure in relation to environment, and van Tieghem was the father of stelar theories. In Britain the main morphological interest was phylogeny and this gave the bias to anatomical study. Its principles were first applied by Williamson and Scott in their studies of the fossil plants of the coal measures.

In 1883 van Tieghem and Duliott published their theory of the stele. Their deductive conclusions were not checked by ontogenetic studies, with the result that their theories were discredited soon after the intensive investigations on vascular anatomy were begun in Great Britain. In remembering those who contributed most to this aspect of vascular anatomy, Gwynne-Vaughan comes first, not only on account of his sustained work, but also for the fact that he recognised and demonstrated the importance of the leaf trace.

In the year 1900 came the independent discovery of Mendel and the confirmation of his work by de Vries, Correns, and Tschermak. The effect was immediate; for the first time genetics had definite laws which clarified many problems and disclosed others. It is doubtful if any other botanical work caused such an outburst of activity over so wide a field: a period of intense plant breeding set in; the transmission of colour factors stimulated the investigation of the flavones and other petal pigments and also the distribution of oxidases; and a new field of vast area was opened to the cytologist.

This period is remarkable also in that it saw the growth of ecology. The first ecologist was Humboldt, who was the first to realise the units of vegetation and to essay their ordering (1805). His system, however, was unsound. Interest in the subject was resuscitated by Grisebach in 1872.

Ecological study now became very active: in France, Bonnier was at work on the influence of edaphic factors on plant form and Flahault on vegetation surveys. In Belgium, Massart was studying littoral vegetation (1893), and in Switzerland there was Schröter. A few years later saw the entry of the American vanguard—Clements, Cowles, Harshberger, Livingstone, and Pound,

and in New Zealand Cockayne followed his lonely bent (1901). Robert Smith was the pioneer in Great Britain: in 1898 he published his observations on the plant associations of the Tay Basin, and this was succeeded by various other communications. He was followed by his brother, W. G. Smith, Moss, Rankin, Lewis, Tansley, Yapp, Oliver, and others.

Another noteworthy series of observations of this period was those on the ascent of sap. Strasburger's experiments pushed home the idea that the phenomenon was entirely physical, a view which was adopted by Askenasy, and Dixon and Joly (1895), who realised the significance of the tensile strength of water and on it founded their well-known explanation. Concurrently, the splendid lead given by Pfeffer and de Vries was vigorously followed, and the problems of permeability, osmotic pressure, and, in general, the water relations of the cell, were the subject of hundreds of papers.

Mention also must be made of the progress of knowledge of the gaseous diffusion into and out from the ordinary leaf. F. F. Blackman showed that the evolution of carbon dioxide during respiration and its intake during carbon assimilation was, in general, proportional to the number of stomates on the leaf surfaces. These results were confirmed by Brown and Escombe (1905). Their classic work on the static diffusion of gases was published in 1900, the kernel of which is that the rate of diffusion through the pore of a stomate is governed by the law of diameters.

Turning to metabolism, this third period is remarkable for the clearer focusing of the circumstances attending metabolic activity, and especially of the action and interaction of those factors which influence the rate of particular activities. Before 1905 these factors were considered separately and various optima were given for various functions. The insufficiency of this was soon appreciated by F. F. Blackman, and his application of Liebig's law of the minimum to physiological processes led to his conception of limiting factors. The truth of the doctrine is generally admitted, but the observations of Boysen-Jensen, Harder, Lundegardt, Warburg, and others indicate that a particular factor is only strictly limiting when it is very much weaker than the others.

It has been mentioned that Hoppe-Seyler established a chemical relationship between chlorophyll and hæmoglobin. His work was carried much further by Marchlewski, Nencki, Schunck, and Zaleski. Their work paved the way for Willstätter, who with his fellow-workers, by improved methods of experiment, obtained a number of products—the phytychlorins and the phytyrhodins. The fact was established that magnesium is an essential constituent of chlorophyll and that the pigment is an ester of the alcohol phytyl. These facts led to the chemical characterisation of two chlorophylls, and, later, of carotin and xanthophyll. A few years after, Willstätter and his collaborators elucidated the composition of many anthocyanins.

Of the sequence of events in the elaboration of

food stuffs, but little real progress was made. The work of Brown and Morris (1893) on the chemistry and physiology of foliage leaves, in which they recognised sucrose as the sugar first formed in carbon assimilation, is outstanding and was the first of many investigations amongst which the work of Parkin, Davis, Daish and Sawyer, Gast, Kylin and Weevers may be mentioned.

The work of the Darwins, Sachs and Pfeffer on irritability was continued throughout this period, and, indeed, to the present day.

The natural sequence of this tropistic work was an inquiry into the mechanisms of the sense organs. The statolith theory of graviperception, due to Haberlandt and Némec, was, in general, accepted. In addition to those mentioned, Noll, Czapek, Jost, Vöchting, Oltmanns, de Vries, and Rothert contributed much to the elucidation of the problems. There remained the question of the transmission of the stimulus from the perceptive organ to the motive region. Boysen-Jensen (1910) showed that a stimulus could pass through a water-gap but not through a thin plate of mica. Thus arose the idea that definite bodies, hormones, were generated in the perceptive region by the action of the external agents and travelled to the motive organs to activate the visible reaction. These observations were continued, and amongst those who contributed were Nielson, Paal, Purdie, Rothert, Stark, Dreschel, Snow, and especially Went.

PRESENT-DAY BOTANY.

The present period is chiefly remarkable for the great output of work in those branches of botany which have an applied aspect. Thus mycology, plant pathology, genetics and cytology occupy a prominent position. In physiology, but little progress has been made in the elucidation of the serial events of the various metabolic changes. But the closer examination of the governing factors, the conditions of growth and the more detailed analyses of these activities, have secured our knowledge, inadequate though it be, more firmly, and also have disclosed certain relationships of first-rate importance, hitherto unsuspected—photo-periodism, in which Garner and Allard were the pioneers; the carbohydrate-nitrogen ratio, to our knowledge of which Kraus and Kraybill have contributed much; and the principle of predetermination so clearly demonstrated by Balls. The application of the principle of the hydrogen ion concentration has given an instrument of great precision capable of use in the investigation of a wide range of problems. The cell wall constituents have been the subject of intensive investigations, some dictated by the requirements of industry, and important discoveries and elucidations have been made.

A survey of this period shows two well-defined tendencies: specialisation and the obliteration of the artificial margins between botany and cognate studies.

The study of physiology in particular and of many aspects of ecology require the application

of the technique and conceptions of physics and especially of chemistry: this will increase in the future; if the approach to such problems be made from the biological point of view, progress will be more assured.

The stress of the War soon forced lessons home, and amongst other things learnt was the dependence of man on the plant. The experience of the War was, curiously enough, not forgotten when peace came, with the result that the training of young botanists for economic work and for the investigation of definite problems became with time a settled policy. But other nations also have learnt the lesson—low living ever has been an incentive to high thinking—with the result of gross over-production—some prefer to call it under-consumption—of essential commodities such as wheat, sugar, and rubber. This is mostly due to the unconsidered use of the great advances of applied botanical knowledge and of agricultural engineering without regard to economic consequences. The present tendency in States and industries to curtail expenditure on research and expert knowledge is a wrong policy: "They that be whole need not a physician, but they that are sick".

The successful conduct of these and many like activities is impossible without expert advice. In Great Britain this is provided by many research stations, and their development is all to the good. But there are a few dangers. The founders of new research institutions may be unmindful of the fact that results cannot be commanded. This may react on the investigators, who may be tempted to justify themselves by the dissemination of unripe fruit. Again, the promotion of applied research may tend to obscure the value of pure research. Lastly, there is the comparative isolation of the workers from their academic brothers so that both lose a source of stimulation and ideas. These possible dangers will be eliminated if there be some connexion, the closer the better, between the botanical departments of universities and the research institutions, accompanied by an occasional interchange of the workers in both.

The British Commonwealth of Nations is, in the main, an agricultural Empire: the great need for trained botanists for its administrative and technical service is patent; the problem is their supply and their training. The increased demand is slowly having its effect in the universities, and more students are taking botany for their finals, but their numbers are too few. In the university the training of young botanists for the first three years must be in pure science; this is absolutely essential, and should be followed by a period of appropriate specialised training.

This raises the problem of the manning of our botanical departments: universities cannot compete with industry in matters of salary, and this in the past has resulted in good men being lost to academic work, especially in plant physiology. If good material for the service of the State and of industry is to be provided and pure research maintained, an adequate flow of recruits of the highest quality into academic life is essential.