

## Letter to the Editor

## Testing Early-Onset Chronic Atypical Depression Subtype

Franco Benazzi<sup>\*,1,2,3</sup><sup>1</sup>E Hecker Outpatient Psychiatry Center, Ravenna, Italy; <sup>2</sup>Department of Psychology, University of Bologna, Bologna, Italy; <sup>3</sup>Department of Psychiatry, National Health Service, Forlì, Italy

Neuropsychopharmacology (2004) 29, 440–441. doi:10.1038/sj.npp.1300355

Sir

A recent paper in your journal (Stewart *et al*, 2002), about DSM-IV atypical depression (AD), showed that the typical better response to MAOI than to TCA was seen only in the early-onset and chronic subtype, suggesting the need to better study this important subtype. The study aim was to find support for this subtyping of AD. A large database by the author, collected during the last 5 years for other studies (so eliminating any interviewer bias), was scanned for such evidence. Study methods, briefly reported below, are fully explained in previous reports (Benazzi, 1999, 2000a, b, 2002a–c, 2003a, b).

## METHODS

## Study Setting

An outpatient psychiatry private practice, more representative of mood disorders usually seen in clinical practice in Italy.

**Interviewer.** A senior clinical and mood disorder research psychiatrist.

**Patients.** Consecutive 320 bipolar-II (BP-II) and 243 major depressive disorder (MDD) outpatients, presenting voluntarily for MDE treatment. Substance-related and borderline personality disorders were excluded because of confounding diagnosis of BP-II, and rare in the setting. Clinically significant general medical illnesses and cognitive disorders were also excluded.

**Interview methods.** During the assessment visit (off psychoactive drugs for at least 2 weeks, apart from a few cases on small doses of benzodiazepines), the following

instruments were used: (1) Structured Clinical Interview for DSM-IV Axis I Disorders-Clinician Version (SCID-CV) (First *et al*, 1997). (2) Global Assessment of Functioning scale (GAF, in SCID-CV) for index MDE severity. (3) Family History Screen (Weissman *et al*, 2000). Often, family members/close friends supplemented clinical information during interview. DSM-IV criteria for MDE with atypical features specifier (AD) were followed. 'Early onset' was defined as the onset of first MDE before age 21 years (according to Stewart *et al*, 2002), 'chronic' as index MDE symptoms lasting more than 2 years. Variables often reported to distinguish atypical vs nonatypical depression were assessed (see references by Benazzi, Agosti and Stewart, 2001; Angst *et al*, 2002; Matza *et al*, 2003 and Rabkin *et al*, 1996). Early-onset chronic AD (EO-C-AD) was compared to nonearly-onset, nonchronic AD (non-EO-C-AD).

## Statistics

Logistic regression was used to study associations and to control for confounding and interactions (STATA 7). *P*-values were two-tailed,  $\alpha$  level 0.05.

## RESULTS

AD was present in 44.5% (251/563). EO-C-AD was present in 26.2% of AD (66/251). In the AD sample, univariate logistic regression of EO-C-AD vs BP-II, index age, female gender, recurrences, axis I comorbidity, GAF, bipolar family history, DSM-IV atypical symptoms, found significantly associated BP-II (odds ratio = 2.2, 95% CI = 1.2–4.0), age (odds ratio = 0.9, 95% CI = 0.9–0.9), more recurrences (odds ratio = 4.4, 95% CI = 2.3–8.3), more axis I comorbidity (odds ratio = 2.4, 95% CI = 1.4–4.4), more bipolar (I + II) family history (odds ratio = 2.9, 95% CI = 1.4–5.9), and hypersomnia (odds ratio = 1.8, 95% CI = 1.0–3.2). Multivariate logistic regression of EO-C-AD vs variables found significant in the univariate analysis (not including age), including all possible interactions, found that recurrences (odds ratio = 4.3, 95% CI = 1.8–10.0), axis I comorbidity (odds ratio = 2.4, 95% CI = 1.2–5.1), bipolar family history (odds ratio = 2.7, 95% CI = 1.2–6.1), and hypersomnia

\*Correspondence: Dr F Benazzi, Via Pozzetto 17, 48010 Castiglione di Cervia RA, Italy, Tel: +39 335 6191 852, Fax: +39 054 330 069, E-mail: f.benazzi@fo.nettuno.it, FrancoBenazzi@fbenazzi.it  
Received 24 August 2003; accepted 16 October 2003  
Online publication: 20 October 2003 at <http://www.acnp.org/citations/Npp102003387/default.pdf>

(odds ratio = 2.5, 95% CI = 1.1–5.3) were still significant and independently associated.

## DISCUSSION

Findings support, on clinical and family history grounds, a distinction between EO-C-AD and non-EO-C-AD. The findings support the pharmacological distinction found by Stewart *et al* (2002).

## REFERENCES

- Agosti V, Stewart JW (2001). Atypical and non-atypical subtypes of depression: comparison of social functioning, symptoms, course of illness, co-morbidity and demographic features. *J Affect Disord* **65**: 75–79.
- Angst J, Gamma A, Sellaro R, Zhang H, Merikangas K (2002). Toward validation of atypical depression in the community: results of the Zurich cohort study. *J Affect Disord* **72**: 125–138.
- Benazzi F (1999). Chronic atypical major depressive episode in private practice: unipolar and bipolar II. *Acta Psychiatr Scand* **100**: 418–423.
- Benazzi F (2000a). Depression with DSM-IV atypical features: a marker of bipolar II disorder. *Eur Arch Psychiatry Clin Neurosci* **250**: 53–55.
- Benazzi F (2000b). Early-onset versus late-onset atypical depression: unipolar and bipolar II. *J Affect Disord* **61**: 95–99.
- Benazzi F (2002a). Psychomotor changes in melancholic and atypical depression: unipolar and bipolar-II subtypes. *Psychiatry Res* **112**: 211–220.
- Benazzi F (2002b). Can only reversed vegetative symptoms define atypical depression? *Eur Arch Psychiatry Clin Neurosci* **252**: 288–293.
- Benazzi F (2002c). Should mood reactivity be included in DSM-IV atypical features specifier? *Eur Arch Psychiatry Clin Neurosci* **252**: 135–140.
- Benazzi F (2003a). Is there a link between atypical and early-onset ‘unipolar’ depression and bipolar II disorder? *Compr Psychiatry* **44**: 102–109.
- Benazzi F (2003b). Testing DSM-IV definition of atypical depression. *Ann Clin Psychiatry* **15**: 9–16.
- First MB, Spitzer RL, Gibbon M, Williams JBW (1997). *Structured Clinical Interview for DSM-IV Axis I Disorders—Clinician Version (SCID-CV)*. American Psychiatric Press: Washington, DC.
- Matza LS, Revicki DA, Davidson JR, Stewart JW (2003). Depression with atypical features in the National Comorbidity Survey: classification, description, and consequences. *Arch Gen Psychiatry* **60**: 817–826.
- Rabkin JG, Stewart JW, Quitkin FM, McGrath PJ, Harrison WM, Klein DF (1996). Should atypical depression be included in DSM-IV? In: Widiger TA, Frances AJ, Pincus HA, Ross R, First MB, Davis WW. *DSM-IV Sourcebook*, Vol. 2 American Psychiatric Association: Washington, DC. pp 239–260.
- Stewart JW, McGrath PJ, Quitkin FM (2002). Do age of onset and course of illness predict different treatment outcome among DSM IV depressive disorders with atypical features? *Neuropsychopharmacology* **26**: 237–245.
- Weissman MM, Wickramaratne P, Adams P, Wolk S, Verdelli H, Olfson M (2000). Brief screening for family psychiatric history. The family history screen. *Arch Gen Psychiatry* **57**: 675–682.