Chronic Lithium Chloride Administration to Unanesthetized Rats Attenuates Brain Dopamine D₂-Like Receptor-Initiated Signaling Via Arachidonic Acid

Mireille Basselin^{*,1}, Lisa Chang¹, Jane M Bell¹ and Stanley I Rapoport¹

¹Brain Physiology and Metabolism Section, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA

We studied the effect of lithium chloride on dopaminergic neurotransmission via D₂-like receptors coupled to phospholipase A₂ (PLA₂). In unanesthetized rats injected i.v. with radiolabeled arachidonic acid (AA, 20:4 n-6), regional PLA₂ activation was imaged by measuring regional incorporation coefficients k^* of AA (brain radioactivity divided by integrated plasma radioactivity) using quantitative autoradiography, following administration of the D₂-like receptor agonist, quinpirole. In rats fed a control diet, quinpirole at 1 mg/kg i.v. increased k^* for AA significantly in 17 regions with high densities of D₂-like receptors, of 61 regions examined. Increases in k^* were found in the prefrontal cortex, frontal cortex, accumbens nucleus, caudate–putamen, substantia nigra, and ventral tegmental area. Quinpirole, 0.25 mg/kg i.v. enhanced k^* significantly only in the caudate–putamen. In rats fed LiCl for 6 weeks to produce a therapeutically relevant brain lithium concentration, neither 0.25 mg/kg nor 1 mg/kg quinpirole increased k^* significantly in any region. Orofacial movements following quinpirole were modified but not abolished by LiCl feeding. The results suggest that downregulation by lithium of D₂-like receptor signaling involving PLA₂ and AA may contribute to lithium's therapeutic efficacy in bipolar disorder.

Neuropsychopharmacology (2005) **30**, 1064–1075, advance online publication, 6 April 2005; doi:10.1038/sj.npp.1300671

Keywords: lithium; phospholipase A₂; dopamine; quinpirole; arachidonic acid; bipolar disorder

INTRODUCTION

Lithium (Li⁺) has been used to treat bipolar disorder for about 50 years, but its mechanism of action is not agreed upon (Barchas *et al*, 1994; Cade, 1999). One possibility is that it corrects a neurotransmission imbalance that contributes to the disease. Clinical evidence suggests that this imbalance is due to excessive dopaminergic activity but reduced cholinergic activity (Bunney and Garland-Bunney, 1987; Bymaster and Felder, 2002; Janowsky and Overstreet, 1995; Post *et al*, 1980). Thus, cholinomimetics as well as drugs that inhibit dopaminergic transmission (eg olanzapine, haloperidol) have an antimanic action in bipolar disorder (Bhana and Perry, 2001; Bymaster and Felder, 2002), whereas drugs that stimulate dopamine synthesis (levodopa), bind to dopamine receptors (bromocriptine), or reduce dopamine reuptake (amphetamine) often precipitate mania

(Fisher *et al*, 1991; Peet and Peters, 1995; Sultzer and Cummings, 1989).

Some reported effects in rats of chronically administered lithium are consistent with its ameliorating the suggested cholinergic–dopaminergic imbalance of bipolar disorder. Thus, lithium reduces the convulsant threshold to the nonspecific cholinergic muscarinic agonist arecoline and to anticholinesterases (Evans *et al*, 1990; Jope, 1993; Lerer, 1985; Morrisett *et al*, 1987). In this regard, studies in knockout mice indicate that the agonist-induced convulsions are largely mediated by muscarinic M₁ receptors (Bymaster *et al*, 2003; Hamilton *et al*, 1997). However, lithium's effects on dopaminergic neurotransmission are not agreed upon, as lithium is reported to inhibit (Engel and Berggren, 1980; Friedman and Gershon, 1973), increase (Fadda *et al*, 1980; Hesketh *et al*, 1978), or have no effect (Bliss and Ailion, 1970; Ho *et al*, 1970) on brain dopamine turnover. Data on lithium's effects on brain dopaminergic D₂-like receptor density also are conflicting (Dziedzicka-Wasylewska and Wedzony, 1996; Reches *et al*, 1982; Staunton *et al*, 1982b; Wajda *et al*, 1983), as are data on its ability to modify haloperidol-induced supersensitivity (Carvey *et al*, 1990; Gallager *et al*, 1978; Pert *et al*, 1978; Pittman *et al*, 1984; Reches *et al*, 1982; Staunton *et al*, 1982a).

*Correspondence: Dr M Basselin, Brain Physiology and Metabolism Section, National Institute on Aging, National Institutes of Health, Bldg 9, Room 15128, Bethesda, MD 20892, USA, Tel: +1 301 594 5522, Fax: +1 301 402 0074, E-mail: mirvasln@mail.nih.gov

Received 7 July 2004; revised 10 November 2004; accepted 7 December 2004

Online publication: 16 December 2004 at <http://www.acnp.org/citations/NPPI21604040315/default.pdf>

Agonist binding to cholinergic $M_{1,3,5}$ (but not M_2 or M_4) receptors or D_2 -like receptors can be coupled via a G-protein to the activation of phospholipase A_2 (PLA₂) (Axelrod, 1995; Bayon *et al*, 1997; Nilsson *et al*, 1998; Bymaster *et al*, 1999; Vial and Piomelli, 1995), so as to release the second messenger, arachidonic acid (AA, 20:4 n:6), from the stereospecifically numbered (*sn*)-2 position of membrane phospholipids. Nonesterified AA and its bioactive eicosanoid metabolites can influence many physiological processes, including membrane excitability, gene transcription, apoptosis, sleep, and behavior (Fitzpatrick and Soberman, 2001; Shimizu and Wolfe, 1990).

Of the nonesterified AA released by PLA₂ activation, studies in unanesthetized rats show that a fraction is rapidly reincorporated into brain phospholipids, whereas the remainder is lost by metabolism to eicosanoids, cannabinoids, β -oxidation, or other pathways (DeGeorge *et al*, 1991; Rapoport, 2001, 2003). The quantity lost is replaced rapidly by diet-derived nonesterified AA in plasma, since AA cannot be synthesized *de novo* in mammalian tissue and its precursor, linoleic acid (18:2 n-6), is not significantly elongated to AA in the adult brain (DeMar *et al*, 2004; Holman, 1986). Replacement is proportional to PLA₂ activation and can be imaged by injecting radiolabeled AA intravenously and measuring regional brain AA incorporation coefficients k^* (brain radioactivity/integrated plasma radioactivity) using quantitative autoradiography. k^* is independent of changes in cerebral blood flow and reflects only the activation process (Chang *et al*, 1997; DeGeorge *et al*, 1991; Rapoport, 2001, 2003; Robinson *et al*, 1992).

Using the fatty acid method, we reported that chronic LiCl feeding to rats, producing a therapeutically relevant brain Li^+ concentration, potentiated brain k^* responses to arecoline (Basselin *et al*, 2003b; Bayon *et al*, 1997; Bosetti *et al*, 2002b; DeGeorge *et al*, 1991; Jones *et al*, 1996). Such potentiation agreed with the hypothesis that lithium might correct a cholinergic–dopaminergic imbalance in bipolar disorder by upregulating cholinergic $M_{1,3,5}$ signaling via PLA₂. In the present study, we determined whether LiCl feeding would downregulate D_2 -like receptor signaling via PLA₂, which would further support the imbalance hypothesis.

We measured k^* for AA in 61 brain regions of unanesthetized rats administered i.v. saline (controls) or the D_2 -like receptor agonist, quinpirole, at 0.25 mg/kg and 1.0 mg/kg i.v. (Bach *et al*, 1980; Malmberg and Mohell, 1995; Seeman and Van Tol, 1994). We chose both doses, which have been used previously in behavioral studies (Bordi and Meller, 1989; Horvitz *et al*, 2001; Koene *et al*, 1993), to examine the dose–response relation to PLA₂ activation. Quinpirole, at 1 mg/kg i.v., is reported to increase k^* for AA in 24 of 29 brain regions having D_2 -like receptors. The increases can be blocked by prior administration of the preferential D_2 -like antagonist (+)-butaclamol (Bristow *et al*, 1998), or enhanced in rat brain regions ipsilateral to a chronic unilateral lesion of the substantia nigra, in proportion to their elevated D_2 -like receptor density (Cory-Slechta *et al*, 1996; Eilam and Szechtman, 1989; Hayakawa *et al*, 2001). Quinpirole at the two doses chosen also can provoke orofacial activity and locomotion (Bordi and Meller, 1989; Horvitz *et al*, 2001; Koene *et al*, 1993), so we measured these as well.

Briefly, we found that 6 weeks of LiCl feeding completely blocked the quinpirole-induced increments in k^* for AA that were evident in control diet-fed rats, and altered, but did not block the stereotypic behavioral responses to quinpirole. Part of this work has been reported in abstract form (Basselin *et al*, 2003a).

MATERIALS AND METHODS

Animals and Diets

Experiments were conducted following the ‘Guide for the Care and Use of Laboratory Animals’ (National Institute of Health Publication No. 86-23) and were approved by the Animal Care and Use Committee of the National Institute of Child Health and Development (NICHD). Male Fischer CDF (F-344)/CrlBR rats (2 months old) (Charles River Laboratories, Wilmington, MA), weighing 180–200 g, were housed in an animal facility in which temperature, humidity, and light cycle were regulated. One group of rats was fed *ad lib* Purina Rat Chow (Harlan Teklad, Madison, WI) containing 1.70 g LiCl per kg in pelleted form for 4 weeks, followed by chow containing 2.55 g LiCl per kg for 2 weeks (Basselin *et al*, 2003b). This feeding regimen produces ‘therapeutically equivalent’ plasma and brain lithium levels of about 0.7 mM (Bosetti *et al*, 2002b; Chang *et al*, 1996). Control rats were fed lithium-free Purina rat chow under parallel conditions. Water and NaCl solution (0.45 M) were available *ad libitum* to both groups.

Drugs

Rats were injected intravenously with 0.9% NaCl, or with 0.25 or 1.0 mg/kg quinpirole hydrochloride (Sigma Chemical Co., Saint Louis, MO) in 0.30 ml 0.9% NaCl. [$1-^{14}C$]AA in ethanol (53 mCi/mmol, >98% pure; Moravsek Biochemicals, Brea, CA) was evaporated and resuspended in HEPES buffer, pH 7.4, which contained 50 mg/ml of bovine serum albumin, as described elsewhere (Hayakawa *et al*, 2001). To confirm tracer purity, gas chromatography was performed after converting fatty acids to their methyl esters using 1 ml 1% sulfuric acid in anhydrous methanol (Makrides *et al*, 1994).

Surgical Procedures and Tracer Infusion

After 6 weeks on a control or a LiCl diet, a rat was anesthetized with halothane (2–3%) through a facemask connected to a vacuum scavenging system. Polyethylene catheters (PE 50) filled with 3% heparinized 0.9% NaCl were surgically inserted into the right femoral artery and vein. The wound was infiltrated with 1% lidocaine and closed with surgical clips. The rat was wrapped loosely, with its upper body remaining free, in a fast-setting plaster cast that was then taped to a wooden block, and allowed to recover from anesthesia for 3–4 h prior to isotope infusion. Body temperature was maintained at 37°C using a feedback heating element and rectal thermometer. Blood pressure and heart rate also were measured. Orofacial activity—sniffing, rigidity, and head turning (Bordi and Meller, 1989; Koene *et al*, 1993)—was monitored in a nonblinded

manner from the beginning of drug infusion to the end of the experiment.

One min after i.v. saline or quinpirole, the unanesthetized rat was infused for 5 min through the femoral vein with 2 ml [14 C]AA (170 μ Ci/kg), at a rate of 400 μ l/min, using an infusion pump (Harvard Apparatus Model 22, Natick, MA). Timed arterial blood samples were collected during infusion and to the time of death at 20 min. At 20 min, when less than 1% of peak radioactivity remained in the plasma (DeGeorge *et al*, 1989; Zhou *et al*, 2002), the rat was killed with an overdose (50 mg/kg i.v.) of sodium pentobarbital (Richmond Veterinary Supply, Richmond, VA) and decapitated. The brain was removed, quickly frozen in 2-methylbutane at -40°C , and stored at -80°C for autoradiography.

Chemical Analysis of Plasma

Arterial plasma samples (30 μ l) were extracted with 3 ml $\text{CHCl}_3:\text{MeOH}$ (2:1, v/v) and 1.5 ml 0.1 M KCl, following a modified Folch technique (Folch *et al*, 1957). Aliquots of 100 μ l of the lower organic phase were used to determine unesterified labeled AA concentrations by liquid scintillation counting. Percent efficiency of ^{14}C counting was 88%.

Quantitative Autoradiography

Brains frozen at -20°C in embedding medium were cut in 20- μm -thick serial coronal sections in a cryostat (Hacker Instruments, Fairfield, NJ). The sections were picked up on glass coverslips, dried on a hot plate at 55°C for at least 5 min, then placed sequentially in an X-ray cassette together with calibrated [^{14}C]methylmethacrylate standards (Amersham, Arlington Heights, IL). Sections and standards were exposed for 6 weeks to autoradiographic film (EMC1, Eastman Kodak Company, Rochester, NY), which was then developed in a dark room. Specific regions were identified by comparing the autoradiographs with an atlas and with published autoradiographs of rat brain (Hayakawa *et al*, 2001; Paxinos and Watson, 1987), and were sampled in both hemispheres. The average of three bilateral measurements for each region from a given brain was used to calculate the regional radioactivity (nCi/g) by digital quantitative densitometry (public domain NIH Image analysis program (version 1.62)). Regional incorporation coefficients of AA from plasma into brain phospholipids, k^* (ml/s/g brain), were calculated as

$$k^* = \frac{c_{\text{brain}}^*(20 \text{ min})}{\int_0^{20} c_{\text{plasma}}^* dt} \quad (1)$$

where $c_{\text{brain}}^*(20 \text{ min})$ nCi/g equals brain radioactivity at the time of death, c_{plasma}^* nCi/ml equals plasma AA radioactivity determined by scintillation counting, and t equals time after beginning [^{14}C]AA infusion.

Statistical Analysis

One-way ANOVA, with Tukey–Kramer's *post hoc* test with correction for five comparisons, was performed in each of the 61 brain regions, to compare quinpirole and i.v. saline responses between control diet and LiCl-fed rats, as well as between i.v. saline responses in LiCl-fed rats and control

diet rats. Other comparisons were not considered relevant. Corrections for multiple comparisons across regions were not made because the purpose of the study was to identify the regions that were involved in individual drug and group-diet effects. Additionally, unpaired *t*-tests were used to compare the durations of behavioral activity and calm periods between the two diet groups. The GraphPad Prism version 3.0a for Macintosh (GraphPad Software, San Diego, CA, <http://www.graphpad.com>) was used for these comparisons. Data are reported as means $\pm$ SD, with statistical significance taken as $p \leq 0.05$.

RESULTS

Physiological Parameters and Behavioral Responses

LiCl-fed rats gained weight more slowly than did control rats, as has been reported (Pittman *et al*, 1984). At the time of surgery, the LiCl-fed rats weighed about 20% less than the control diet-fed group (254 ± 21 g, $n = 26$ vs 308 ± 11 g, $n = 24$, $p < 0.0001$).

Quinpirole, 0.25 or 1.0 mg/kg i.v., provoked behavioral cycles, each consisting of an 'activity' period (repetitive sniffing, mouth, and head-turning) followed by a 'calm' period, whereas i.v. saline had no effect on activity (data not shown). The quinpirole-induced cycles were repeated during the 20-min period following the drug, and the lengths of each component were averaged during this time. As illustrated in Table 1, there was no significant difference between LiCl-fed and control rats in the mean duration of the 'activity' period at either quinpirole dose, whereas the 'calm' period was prolonged in the LiCl-fed compared with control diet rats at both doses.

LiCl Effect on Baseline Values of k^*

Figure 1 presents autoradiographs from the following: (A) a rat fed a control diet and given saline i.v., (B) a rat fed a control diet and given quinpirole 1 mg/kg i.v., (C) a rat fed a LiCl diet and given saline, (D) a rat fed a LiCl diet and given quinpirole 1 mg/kg. Values for regional incorporation coefficients k^* for AA are color-coded in the autoradiographs. The figure shows that quinpirole compared with saline increased k^* in certain brain regions in the control diet-fed (A and B) but not in the LiCl-fed (C and D) rats.

Table 1 Effect of Quinpirole on Orofacial Activity in Unanesthetized Rats Fed a Control or a LiCl Diet

	Control diet		LiCl diet	
	Quinpirole dose			
	0.25 mg/kg (9) ^a	1.0 mg/kg (9)	0.25 mg/kg (10)	1.0 mg/kg (10)
Duration (s)				
Orofacial activity	12±4 ^b	15±3	15±3	17±4
Calm period	60±11	35±4	110±15***	80±7***

^aNumber of animals.

^bMean $\pm$ SD.

***Differs from respective control mean by unpaired *t*-test, $p < 0.001$.

Figure 1 Effects of quinpirole on regional brain incorporation coefficients k^* of arachidonic acid in autoradiographs of coronal brain sections obtained 20 min after infusion of [$1\text{-}^{14}\text{C}$]arachidonate ($170\text{ }\mu\text{Ci/kg}$, i.v.): (a) a control diet rat; (b) a control diet rat plus quinpirole 1 mg/kg ; (c) a LiCl diet rat; (d) a LiCl diet rat plus quinpirole 1 mg/kg . Acb, accumbens nucleus; Aud ctx, auditory cortex; CP, caudate-putamen; DLG, dorsal lateral geniculate; Fr, frontal cortex; PFr, prefrontal cortex; SNR, substantia nigra pars reticulata; Vis ctx, visual cortex; VTA, ventral tegmental area.

This pattern for all rats is summarized in Table 2. Comparison between data columns 4 and 1 in Table 2 shows that LiCl feeding significantly elevated mean baseline values of k^* (k^* following i.v. saline) in 15 of the 61 brain regions examined, compared with their respective values in control diet-fed rats. Significant baseline increases were evident in central auditory regions—auditory cortex (27–34%), habenular nucleus lateral (30%), dorsal geniculate nucleus (38%), geniculate medial (34%), and colliculus inferior (17%)—and in central visual regions—visual cortex (27–33%), habenular nucleus medial (33%), and colliculus superior (22%) (Brodal, 1981; Matsumura *et al*, 1992). Increases were also evident in the cerebellar flocculus (27%), subthalamic nucleus (44%), interpeduncular nucleus (35%), and substantia nigra (24–36%).

Quinpirole Effects on k^* for AA

Table 2 (data columns 3 and 1) shows that in control diet rats quinpirole, at 1 mg/kg compared with saline, significantly increased k^* in 18 of the 61 brain regions examined. Increases were observed in the prefrontal cortex (50%), frontal cortex (26–31%), primary olfactory cortex (32%),

olfactory tubercle (26%), islands of Calleja (26%), accumbens nucleus (29%), caudate-putamen (40%), substantia nigra (33%), and ventral tegmental area (29%). Quinpirole, at 0.25 mg/kg , significantly increased k^* only in ventral, lateral, and medial areas of the caudate-putamen (data columns 1 and 2). In contrast to its excitatory effects in control diet-fed rats, quinpirole at 0.25 or 1.0 mg/kg , compared with saline, did not increase k^* significantly in any region in the LiCl-fed rats (comparisons between data columns 4 and 5, and data columns 4 and 6).

DISCUSSION

Administration of 1 mg/kg i.v. quinpirole to control diet-fed rats increased mean k^* for AA significantly in 17 of the 61 (28%) brain regions examined (Table 2), many of which are rich in D_2 -like (D_2 , D_3 , and D_4) receptors (Cooper *et al*, 1996; Cory-Slechta *et al*, 1996; Defagot *et al*, 1997; Eilam and Szechtman, 1989; Gehlert, 1993; Levant and DeSouza, 1993; Levey *et al*, 1993; Shafer and Levant, 1998; Vincent *et al*, 1995). A lower mean percent increase was reported in our prior study with the same quinpirole dose (Hayakawa *et al*,

Table 2 Effect of Quinpirole^a on Regional Incorporation Coefficients k^* of [$1-^{14}\text{C}$]Arachidonic Acid^b in LiCl-Fed Rats

Brain region	Control diet			Lithium diet		
	Saline (n = 6)	QUINP 0.25 (n = 9)	QUINP 1.0 (n = 9)	Saline (n = 6)	QUINP 0.25 (n = 10)	QUINP 1.0 (n = 10)
<i>Telencephalon</i>						
Prefrontal cortex layer I	4.07 ± 0.25	4.60 ± 0.64	6.05 ± 0.89***	4.36 ± 0.26	4.45 ± 0.67	4.23 ± 0.67
Prefrontal cortex layer IV	4.21 ± 0.28	4.89 ± 0.74	6.37 ± 1.17***	4.37 ± 0.26	4.39 ± 0.67	4.02 ± 0.62
Primary olfactory cortex	3.93 ± 0.34	4.49 ± 0.57	5.20 ± 1.02**	4.09 ± 0.21	4.16 ± 0.77	3.89 ± 0.56
<i>Frontal cortex (10)</i>						
Layer I	4.17 ± 0.34	4.61 ± 0.65	5.24 ± 0.36**	4.17 ± 0.23	4.43 ± 0.91	4.08 ± 0.48
Layer IV	4.31 ± 0.27	4.74 ± 1.00	5.63 ± 0.25**	4.28 ± 0.23	4.65 ± 0.90	4.18 ± 0.69
<i>Frontal cortex (8)</i>						
Layer I	4.11 ± 0.44	4.46 ± 0.71	5.45 ± 0.99*	4.30 ± 0.35	4.40 ± 0.81	4.18 ± 0.63
Layer IV	4.56 ± 0.29	4.95 ± 1.04	5.75 ± 0.79*	4.56 ± 0.44	4.67 ± 0.72	4.30 ± 0.66
Pyriform cortex	3.41 ± 0.42	3.46 ± 0.78	3.86 ± 0.47	3.37 ± 0.27	3.98 ± 0.68	3.66 ± 0.35
Anterior cingulate cortex	4.87 ± 0.56	5.23 ± 0.42	5.31 ± 0.50	4.68 ± 0.30	5.08 ± 0.55	4.55 ± 0.69
<i>Motor cortex</i>						
Layer I	4.49 ± 0.70	4.66 ± 0.93	4.78 ± 0.55	4.17 ± 0.25	4.51 ± 0.65	3.97 ± 0.46
Layer II–III	4.77 ± 0.90	4.73 ± 0.93	5.09 ± 0.40	4.35 ± 0.26	4.65 ± 0.56	4.03 ± 0.52
Layer IV	4.90 ± 0.66	5.03 ± 0.57	5.23 ± 0.46	4.56 ± 0.28	4.92 ± 0.66	4.28 ± 0.57
Layer V	3.79 ± 0.55	4.02 ± 0.83	4.34 ± 0.46	3.60 ± 0.21	3.91 ± 0.39	3.75 ± 0.42
Layer VI	3.63 ± 0.52	3.81 ± 0.60	4.14 ± 0.48	3.46 ± 0.20	3.76 ± 0.43	3.39 ± 0.48
<i>Somatosensory cortex</i>						
Layer I	4.38 ± 0.65	4.67 ± 0.96	5.00 ± 0.73	4.12 ± 0.40	4.63 ± 0.67	4.13 ± 0.62
Layer II–III	4.75 ± 0.65	4.94 ± 0.92	5.04 ± 0.72	4.44 ± 0.35	4.59 ± 0.52	4.11 ± 0.39
Layer IV	4.75 ± 0.41	5.33 ± 0.87	5.26 ± 0.59	4.72 ± 0.25	5.10 ± 0.69	4.39 ± 0.57
Layer V	4.62 ± 0.61	4.95 ± 1.07	5.07 ± 0.64	4.71 ± 1.11	4.55 ± 0.64	4.11 ± 0.47
Layer VI	4.16 ± 0.29	4.78 ± 0.96	5.07 ± 0.64	4.15 ± 0.62	4.49 ± 0.63	4.14 ± 0.64
<i>Auditory cortex</i>						
Layer I	4.55 ± 0.31	4.65 ± 0.97	4.61 ± 0.56	5.79 ± 0.37*	4.97 ± 0.64	5.34 ± 0.60
Layer IV	4.57 ± 0.36	5.18 ± 0.58	4.98 ± 0.52	6.12 ± 0.59***	5.37 ± 0.48	6.10 ± 0.69
Layer VI	4.44 ± 0.18	4.90 ± 0.71	4.49 ± 0.65	5.19 ± 0.54	4.97 ± 0.42	4.22 ± 0.37
<i>Visual cortex</i>						
Layer I	4.57 ± 0.48	4.47 ± 0.77	4.05 ± 0.51	5.91 ± 0.77*	4.91 ± 0.47	5.43 ± 0.83
Layer IV	4.63 ± 0.20	4.66 ± 0.90	4.14 ± 0.65	5.89 ± 0.26*	5.09 ± 0.48	5.31 ± 0.55
Layer VI	4.22 ± 0.20	4.58 ± 0.96	4.17 ± 0.52	5.63 ± 0.42*	4.83 ± 0.44	5.52 ± 0.88
Olfactory tubercle	4.16 ± 0.29	4.39 ± 0.82	5.26 ± 0.44*	4.51 ± 0.46	4.29 ± 0.75	4.01 ± 0.65
Islands of Callaja	4.12 ± 0.45	4.19 ± 0.56	5.20 ± 0.41*	4.33 ± 0.57	3.90 ± 0.82	3.70 ± 0.49
Accumbens nu	4.25 ± 0.34	4.38 ± 0.75	5.50 ± 0.70*	4.16 ± 0.21	4.44 ± 0.99	3.70 ± 0.83
Ventral pallidum	3.44 ± 0.37	3.93 ± 0.62	3.93 ± 0.75	3.48 ± 0.32	3.62 ± 0.66	3.51 ± 0.69
Globus pallidus	3.35 ± 0.30	3.76 ± 0.63	3.69 ± 0.39	3.39 ± 0.30	3.91 ± 0.66	4.03 ± 0.50
Entopeduncular nu	3.49 ± 0.38	3.63 ± 0.80	3.79 ± 0.35	3.70 ± 0.59	3.68 ± 0.57	3.17 ± 0.59
<i>Caudate putamen</i>						
Dorsal	3.92 ± 0.20	4.43 ± 0.71	4.95 ± 0.38**	4.01 ± 0.21	4.42 ± 0.58	4.09 ± 0.56
Ventral	3.19 ± 0.24	4.41 ± 0.74*	4.79 ± 0.40**	4.02 ± 0.22	4.37 ± 0.57	4.11 ± 0.54

Table 2 Continued

Brain region	Control diet			Lithium diet		
	Saline (n = 6)	QUINP 0.25 (n = 9)	QUINP 1.0 (n = 9)	Saline (n = 6)	QUINP 0.25 (n = 10)	QUINP 1.0 (n = 10)
Lateral	3.16 ± 0.23	4.38 ± 0.65*	4.84 ± 0.47**	4.02 ± 0.20	4.39 ± 0.55	4.31 ± 0.60
Medial	3.14 ± 0.24	4.48 ± 0.85*	4.87 ± 0.44**	3.94 ± 0.12	4.34 ± 0.55	4.26 ± 0.63
<i>Hippocampus</i>						
CA1	3.06 ± 0.52	3.18 ± 0.61	3.31 ± 0.53	3.72 ± 0.70	3.19 ± 0.49	3.87 ± 0.67
CA2	3.23 ± 0.59	3.29 ± 0.58	3.22 ± 0.67	3.87 ± 0.61	3.28 ± 0.46	3.05 ± 0.69
CA3	3.60 ± 0.35	3.38 ± 0.64	3.36 ± 0.71	4.09 ± 0.64	3.44 ± 0.56	3.24 ± 0.73
<i>Diencephalon</i>						
Habenular nucleus lateral	5.08 ± 0.42	5.30 ± 0.80	5.78 ± 0.74	6.61 ± 0.34**	7.33 ± 0.57	6.69 ± 0.75
Habenular nucleus medial	5.15 ± 0.20	5.28 ± 0.86	5.81 ± 0.58	6.87 ± 0.54**	7.06 ± 0.54	6.78 ± 0.63
Dorsal lat. geniculate nu	4.92 ± 0.65	4.82 ± 0.47	5.29 ± 0.54	6.79 ± 0.52**	6.82 ± 0.93	6.34 ± 0.49
Geniculate medial	5.11 ± 0.65	4.97 ± 0.42	5.53 ± 0.42	6.85 ± 0.42**	6.92 ± 0.69	6.69 ± 0.57
<i>Thalamus</i>						
Ventroposterior lat. nu	4.51 ± 0.53	4.79 ± 0.64	4.39 ± 0.89	4.68 ± 0.88	4.15 ± 0.68	4.17 ± 0.75
Ventroposterior med. nu	4.48 ± 0.42	4.64 ± 0.97	4.40 ± 0.89	4.78 ± 0.86	4.13 ± 0.64	3.99 ± 0.83
Ventrolateral	4.64 ± 0.44	4.57 ± 0.93	4.56 ± 0.73	4.70 ± 1.15	4.53 ± 0.63	4.17 ± 0.66
Ventromedial	4.15 ± 0.49	4.79 ± 0.86	4.52 ± 0.79	4.94 ± 1.07	4.54 ± 0.60	4.24 ± 0.72
Parafascicular nu	4.60 ± 0.43	4.76 ± 0.37	4.05 ± 0.48	4.75 ± 0.80	4.27 ± 0.69	4.61 ± 0.71
Subthalamic nu	4.56 ± 0.52	4.54 ± 0.76	5.08 ± 0.79	6.55 ± 0.70***	5.64 ± 0.78	6.18 ± 0.56
<i>Mesencephalon</i>						
Interpeduncular nu	5.93 ± 0.52	5.64 ± 0.90	6.79 ± 0.51	7.98 ± 0.60***	7.20 ± 0.43	7.12 ± 0.55
<i>Substantia nigra</i>						
Pars reticulata	4.36 ± 0.48	5.07 ± 0.46	5.79 ± 0.45*	5.40 ± 0.57*	5.70 ± 0.41	5.18 ± 0.71
Pars compacta	4.33 ± 0.81	5.02 ± 1.01	5.75 ± 0.91*	5.90 ± 0.47***	5.55 ± 0.41	5.28 ± 0.75
Ventral tegmental area	4.24 ± 0.14	4.53 ± 0.67	5.46 ± 0.42*	4.17 ± 0.54	4.22 ± 0.83	4.20 ± 0.52
Colliculus superior	4.63 ± 0.29	4.56 ± 0.93	5.16 ± 0.83	5.65 ± 0.44	5.35 ± 0.81	5.29 ± 0.54
Colliculus inferior	6.59 ± 0.21	6.66 ± 0.69	6.62 ± 0.78	7.72 ± 0.50*	7.03 ± 0.48	7.70 ± 0.45
Deep layers of sup. colliculus	5.14 ± 0.49	4.52 ± 0.70	5.15 ± 0.77	5.61 ± 0.45	5.10 ± 0.70	5.15 ± 0.54
Pedunculopontine tegmental nu	3.92 ± 0.30	4.26 ± 0.75	3.66 ± 0.35	4.09 ± 0.50	4.44 ± 0.55	4.26 ± 1.00
<i>Rhombencephalon</i>						
Flocculus	5.60 ± 0.45	5.40 ± 0.79	5.63 ± 0.74	7.12 ± 0.46***	6.93 ± 0.96	6.95 ± 0.41
Cerebellar gray matter	4.84 ± 0.74	5.19 ± 0.71	4.71 ± 0.75	5.24 ± 0.97	4.89 ± 0.50	5.25 ± 0.16
Molecular layer cerebellar gray matter	5.24 ± 0.68	5.70 ± 0.58	5.64 ± 0.35	6.11 ± 0.45	5.50 ± 0.44	5.66 ± 0.47
<i>White matter</i>						
Corpus callosum	3.04 ± 0.41	3.12 ± 0.50	3.70 ± 0.51	2.93 ± 0.12	3.07 ± 0.64	2.90 ± 0.90
Cerebellar white matter	2.76 ± 0.64	3.07 ± 0.68	3.13 ± 0.63	3.09 ± 0.66	3.44 ± 0.60	2.81 ± 0.34

^aQuinpirole administration: QUINP 0.25 mg/kg and 1.0 mg/kg i.v.

^bk* = (ml/s/g) × 10⁻⁴.

Each value is a mean ± SD.

*p < 0.05; **p < 0.01; ***p < 0.001 mean different from control saline mean.

No mean k* in response to 0.25 mg/kg or 1.0 mg/kg quinpirole in the LiCl-fed rats differed significantly from k* in response to i.v. saline LiCl-fed rats.

nu, nucleus; lat, lateral; med, medial.

2001), but in that study only 29 brain regions largely having D₂-like receptors were analyzed, and they were analyzed in the hemisphere contralateral to a chronic unilateral lesion of the nucleus basalis. Such contralateral regions can have disturbed dopaminergic signaling (Lawler *et al*, 1995).

Quinpirole 1 mg/kg significantly increased k^* by 49–51% in the prefrontal cortex, which is known to receive dopamine fibers from the ventral tegmental area (Brozoski *et al*, 1979) and to have a high density of D₄ receptors (Ariano *et al*, 1997; Mrzljak *et al*, 1996). An increment in k^* in the Islands of Callaja may represent selective activation of D₃ receptors, since this region lacks D₂ receptors (Levant and DeSouza, 1993; Levey *et al*, 1993; Shafer and Levant, 1998). Quinpirole at 0.25 mg/kg significantly increased k^* only in caudate–putamen areas, by 38–43%, consistent with a dose-response relation of D₂-like receptor mediated activation of PLA₂. The caudate–putamen has a high density of D₂-like receptors and receives strong dopaminergic input from the substantia nigra (Levant and DeSouza, 1993). In contrast, in the rats fed LiCl, quinpirole at the two doses did not provoke a statistically significant change in k^* in any of the 61 brain regions.

In the control diet-fed rats, quinpirole at 0.25 and 1.0 mg/kg provoked repeated cycles of orofacial activity followed by a calm period. Such cycles have been ascribed to activation of D₂-like receptors in the striatum and nucleus accumbens (Bordi and Meller, 1989; Koene *et al*, 1993). That the cycles were altered but were not blocked by LiCl feeding (Table 1) suggests that they are mediated by D₂-coupled signaling pathways involving effector enzymes other than or in addition to PLA₂ (Arnt *et al*, 1987; Cooper *et al*, 1996; Delfs and Kelley, 1990; Senogles, 2000).

Chronic lithium's inhibition of quinpirole-induced elevations in k^* for AA contrasts with its ability to potentiate k^* responses in rats given the cholinergic muscarinic agonist arecoline (Basselin *et al*, 2003b), and with its ability to reduce the convulsant threshold in rats to arecoline and other cholinomimetics (Evans *et al*, 1990; Jope, 1993; Lerer, 1985; Morrisett *et al*, 1987). As k^* responses to arecoline and quinpirole represent PLA₂ activation coupled to M_{1,3,5} and D₂-like receptors, respectively (Bayon *et al*, 1997; DeGeorge *et al*, 1991; Hayakawa *et al*, 2001; Nilsson *et al*, 1998; Rapoport, 2001; Vial and Piomelli, 1995), the arecoline and quinpirole data suggest that lithium affects M_{1,3,5} and D₂-like receptor-mediated activation of PLA₂ in opposite ways. In contrast, LiCl has a more complex effect on k^* responses in rats to DOI, an agonist of 5-HT_{2A/2C} receptors that also can be coupled to PLA₂ (Basselin *et al*, 2005; Felder *et al*, 1990; Qu *et al*, 2003).

The hypothesis that bipolar disorder symptoms arise from excess dopaminergic compared with reduced cholinergic signaling is based, in part, on reports that cholinomimetics as well as inhibitors of dopamine signaling can be therapeutic in bipolar disorder (see Introduction) (Bunney and Garland-Bunney, 1987; Bymaster and Felder, 2002; Janowsky and Overstreet, 1995; Post *et al*, 1980). Nevertheless, a hyper-dopaminergic state has not been demonstrated directly in bipolar disorder, and it is not clear if brain dopamine receptor density is altered in the disease (Bowden *et al*, 1997; Ebert *et al*, 1996; Wong *et al*, 1997b). Reports do suggest that the dopamine reuptake transporter is altered (Greenwood *et al*, 2001) and that a D₂ receptor

polymorphism can contribute to the disease (Massat *et al*, 2002).

If lithium's therapeutic efficacy in bipolar disorder depends partly on its normalizing a cholinergic–dopaminergic neurotransmission imbalance, our results suggest that the imbalance involves signaling via PLA₂ rather than via other receptor-coupled effector enzymes (eg adenylate cyclase, phospholipase C (PLC)) (Cooper *et al*, 1996). Supporting this interpretation are observations in rats that lithium feeding reduces arecoline-initiated hydrolysis of phosphatidylinositol-4,5-bisphosphate by PLC in the brain, that lithium is a pro-convulsant with regard to arecoline and other cholinomimetics (Casebolt and Jope, 1989; Ormandy *et al*, 1991; Song and Jope, 1992), and that it does not inhibit dopamine-sensitive accumulation of cyclic AMP in the guinea pig brain (Reches *et al*, 1978).

In our study (Table 2), chronic LiCl blocked the quinpirole-evoked elevation of k^* for AA in the prefrontal cortex and caudate–putamen, regions having D₂-like receptors. In bipolar disorder patients, glucose metabolism has been reported to be reduced in the prefrontal cortex, particularly in unipolar and bipolar depression (Baxter *et al*, 1985; Drevets *et al*, 1997; Ketter *et al*, 2001). Additionally, D₂ receptor density has been reported to be elevated in the caudate–putamen of bipolar disorder patients (Pearlson *et al*, 1995; Wong *et al*, 1997a) but not of nonpsychotic manic patients (Yatham *et al*, 2002). Our finding that chronic lithium downregulated D₂-like receptor-mediated neurotransmission via AA in the caudate–putamen (k^* was elevated even at 0.25 mg/kg i.v. quinpirole) suggests that lithium may be therapeutic in bipolar patients by counteracting signaling effects of increased D₂ receptor density in the caudate–putamen.

Extrapyramidal motor symptoms of Parkinson disease are considered to reflect reduced dopaminergic compared with increased cholinergic neurotransmission (Brooks *et al*, 1995; Katzenschlager *et al*, 2003; LeWitt and Nyholm, 2004; Sovner and DiMascio, 1978). As LiCl feeding to rats potentiates M_{1,3,5} receptor-signaling via AA (Basselin *et al*, 2003b) but suppresses D₂-like receptor-signaling via AA (Table 2), it is not surprising that lithium in humans frequently causes a tremor and, less frequently, severe extrapyramidal symptoms resembling those of Parkinson disease (Ghadirian *et al*, 1996; Rondot and Bathien, 1979; Sansone and Ziegler, 1985), and that lithium can exacerbate extrapyramidal symptoms in Parkinson patients (Dalen and Steg, 1973).

There are three major PLA₂ enzyme classes in the brain, a Ca²⁺-dependent cytosolic PLA₂ selective for AA, a Ca²⁺-dependent secretory PLA₂, and a Ca²⁺-independent PLA₂ that is more selective for docosahexaenoic acid (22:6 n-3) than for AA (Dennis, 1994; Strokin *et al*, 2003). D₂-like receptors can be coupled specifically to cPLA₂ (Nilsson *et al*, 1998; Vial and Piomelli, 1995), which is localized at postsynaptic neuronal sites (Basavarajappa *et al*, 1998; Ong *et al*, 1999; Pardue *et al*, 2003; Tu and Bazan, 2003) as well as at astrocytes (Stephenson *et al*, 1996). Quinpirole-evoked increases in k^* for AA are likely mediated at the neuronal receptors (Cory-Slechta *et al*, 1996; Eilam and Szechtman, 1989; Reus *et al*, 2001), as the increments occur in regions with high neuronal D₂-like receptor densities (Cooper *et al*, 1996; Gerfen *et al*, 1995; Hayakawa *et al*, 2001;

Wedzony *et al*, 2000) and are potentiated where neuronal D₂-like receptor density is elevated in a rat model of Parkinson disease (Hayakawa *et al*, 2001). Additionally, quinpirole is reported to reduce regional glucose metabolism, measured with 2-deoxy-D-glucose, in the caudate/putamen, lateral habenular, and motor cortex of rats (Carpenter *et al*, 2003; Basselin *et al*, unpublished observations; Engber *et al*, 1990). Decreased glucose metabolism is considered to represent reduced firing of axon terminals derived from neuronal cell bodies (Sokoloff, 1999).

Chronic LiCl feeding to rats is reported to transcriptionally downregulate brain cPLA₂ expression, without affecting sPLA₂ or iPLA₂ expression (Bosetti *et al*, 2002a; Rintala *et al*, 1999; Weerasinghe *et al*, 2004). Thus, the blocking by lithium of *k** responses to quinpirole may be due to reduced expression of the cPLA₂ coupled to D₂ receptors (Vial and Piomelli, 1995). Lithium also has been reported to reduce brain levels of the inhibitory G α protein subunits—G α _{i1} and G α _{i2}—that couple D₂-like receptors to PLA₂ or PLC (Carli *et al*, 1994; Di Marzo *et al*, 1993; Sidhu and Niznik, 2000; Winitz *et al*, 1994), and to reduce GTP binding to these subunits (Colin *et al*, 1991; Jakobsen and Wiborg, 1998; Li *et al*, 1991; Wang and Friedman, 1999). These effects also may contribute to its ability to block quinpirole *k** responses.

In contrast, the G α_q subunits of G-proteins that can couple M_{1,3,5} and serotonergic 5-HT receptors to PLC or PLA₂ (Chen *et al*, 1999; Cooper *et al*, 1996; Roth *et al*, 1998; Sidhu and Niznik, 2000) are not reported to be reduced by chronic lithium (Dwivedi and Pandey, 1997; Wang and Friedman, 1999). This may explain why lithium does not reduce *k** responses to the muscarinic receptor agonist arecoline (it even increases them), nor to the 5-HT_{2A/2C} receptor agonist, DOI, in most brain regions (responses are blocked by lithium in visual and auditory areas in which baseline values of *k** are elevated) (Basselin *et al*, 2005, 2003b).

In humans, lithium has been reported to augment the amplitudes of the P1/N1 components of auditory evoked responses and of the 65-P95 and P95-N125 components of visual evoked responses (Fenwick and Robertson, 1983; Hegerl *et al*, 1990; Ulrich *et al*, 1990). The finding that LiCl feeding upregulated baseline values of *k** for AA in central auditory and visual areas of rat brain (see Results), which agrees with our previous observations (Basselin *et al*, 2005, 2003b), may be related to these clinical effects and to lithium's ability to reduce visual and auditory hallucinations in bipolar disorder patients (Goodnick and Meltzer, 1984; Potash *et al*, 2001).

In a rat model of neuroinflammation, elevated baseline values for *k** correlate with increased activities of cPLA₂ and sPLA₂ but not iPLA₂, as well as increased turnover of AA within brain phospholipids (Lee *et al*, 2004; Rosenberger *et al*, 2004). These correlations suggest that the elevated baseline values for *k** in brain visual and auditory regions in LiCl fed rats (Table 2) also reflect increased PLA₂ activity and increased AA turnover, possibly due to increased retinal and cochlear inputs, respectively, to these regions (Jung and Reme, 1994; Pfeilschifter *et al*, 1988).

In summary, chronic LiCl feeding prevents regional increases in brain incorporation coefficients *k** for AA that can be provoked in control diet-fed rats by the D₂-like

receptor-agonist, quinpirole. The increases likely represent augmented quinpirole-induced PLA₂-mediated release of AA from membrane phospholipids. Their inhibition suggests that lithium's efficacy in bipolar disorder is related to its ability to downregulate the D₂ receptor-mediated activation of PLA₂. This suggestion might be tested in human subjects by means of positron emission (Esposito *et al*, 2003; Giovacchini *et al*, 2002).

ACKNOWLEDGEMENTS

We thank Dr Richard Bazinet for his helpful comments on this paper.

REFERENCES

- Ariano MA, Wang J, Noblett KL, Larson ER, Sibley DR (1997). Cellular distribution of the rat D₄ dopamine receptor protein in the CNS using anti-receptor antisera. *Brain Res* 752: 26–34.
- Arnt J, Hyttel J, Perregaard J (1987). Dopamine D-1 receptor agonists combined with the selective D-2 agonist quinpirole facilitate the expression of oral stereotyped behaviour in rats. *Eur J Pharmacol* 133: 137–145.
- Axelrod J (1995). Phospholipase A₂ and G proteins. *Trends Neurosci* 18: 64–65.
- Bach NJ, Kornfeld EC, Jones ND, Chaney MO, Dorman DE, Paschal JW *et al* (1980). Bicyclic and tricyclic ergoline partial structures. Rigid 3-(2-aminoethyl)pyrroles and 3- and 4-(2-aminoethyl)pyrazoles as dopamine agonists. *J Med Chem* 23: 481–491.
- Barchas JD, Hamblin MW, Malenka RC (1994). Biochemical hypotheses of mood and anxiety disorders. In: Siegel GJ, Agranoff BW, Albers RW, Molinoff PB (eds). *Basic Neurochemistry*, 5th edn. Raven Press: New York. pp 979–1001.
- Basavarajappa BS, Cooper TB, Hungund BL (1998). Effect of chronic ethanol exposure on mouse brain arachidonic acid specific phospholipase A₂. *Biochem Pharmacol* 55: 515–521.
- Basselin M, Chang L, Seemann R, Bell JM, Rapoport SI (2003a). Chronic lithium administration increases cholinergic and decreases dopaminergic-agonist induced signal transduction via arachidonic acid in awake rats. *Soc Neurosci Abstr* 33: 799.1.
- Basselin M, Chang L, Seemann R, Bell JM, Rapoport SI (2003b). Chronic lithium administration potentiates brain arachidonic acid signaling at rest and during cholinergic activation in awake rats. *J Neurochem* 85: 1553–1562.
- Basselin M, Chang L, Seemann R, Bell JM, Rapoport SI (2005). Chronic lithium administration to rats selectively modified 5-HT_{2A/2C} receptor-mediated brain signaling via arachidonic acid. *Neuropsychopharmacology* 30: 461–472.
- Baxter Jr LR, Phelps ME, Mazziotta JC, Schwartz JM, Gerner RH, Selin CE *et al* (1985). Cerebral metabolic rates for glucose in mood disorders. Studies with positron emission tomography and fluorodeoxyglucose F 18. *Arch Gen Psychiatry* 42: 441–447.
- Bayon Y, Hernandez M, Alonso A, Nunez L, Garcia-Sancho J, Leslie C *et al* (1997). Cytosolic phospholipase A₂ is coupled to muscarinic receptors in the human astrocytoma cell line 1321N1: characterization of the transducing mechanism. *Biochem J* 323: 281–287.
- Bhana N, Perry CM (2001). Olanzapine: a review of its use in the treatment of bipolar I disorder. *CNS Drugs* 15: 871–904.
- Bliss EL, Ailion J (1970). The effect of lithium upon brain neuroamines. *Brain Res* 24: 305–310.
- Bordi F, Meller E (1989). Enhanced behavioral stereotypies elicited by intrastriatal injection D₁ and D₂ dopamine agonists in intact rats. *Brain Res* 504: 276–283.
- Bosetti F, Rintala J, Rosenberger TA, Contreras MA, Rapoport SI, Chang MCJ (2002a). Chronic lithium downregulates cycloo-

- xygenase-2 and prostaglandin E₂ concentration in rat brain. *Mol Psychiatry* 202: 844–849.
- Bosetti F, Seemann R, Bell JM, Zahorchak R, Friedman E, Rapoport SI et al (2002b). Analysis of gene expression with cDNA microarrays in rat brain after 7 and 42 days of oral lithium administration. *Brain Res Bull* 57: 205–209.
- Bowden C, Theodorou AE, Cheetham SC, Lowther S, Katona CL, Crompton MR et al (1997). Dopamine D₁ and D₂ receptor binding sites in brain samples from depressed suicides and controls. *Brain Res* 752: 227–233.
- Bristow LJ, Cook GP, Patel S, Curtis N, Mawer I, Kulagowski JJ (1998). Discriminative stimulus properties of the putative dopamine D₃ receptor agonist, (+)-PD 128907: role of presynaptic dopamine D₂ autoreceptors. *Neuropharmacology* 37: 793–802.
- Brodal A (1981). *Neurological Anatomy in Relation to Clinical Medicine* 3rd edn. Oxford University Press: Oxford.
- Brooks DJ, Torjanski N, Burn DJ (1995). Ropinirole in the symptomatic treatment of Parkinson's disease. *J Neural Transm Suppl* 45: 231–238.
- Brozoski TJ, Brown RM, Rosvold HE, Goldman PS (1979). Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science* 205: 929–932.
- Bunney WEJ, Garland-Bunney BL (1987). Mechanisms of action of lithium in affective illness: basic and clinical implications. In: Meltzer HY (ed). *Psychopharmacology: The Third Generation of Progress*. Raven: New York. pp 553–565.
- Bymaster FP, Calligaro DO, Falcone JF (1999). Arachidonic acid release in cell lines transfected with muscarinic receptors: a simple functional assay to determine response of agonists. *Cell Signal* 11: 405–413.
- Bymaster FP, Carter PA, Yamada M, Gomeza J, Wess J, Hamilton SE et al (2003). Role of specific muscarinic receptor subtypes in cholinergic parasympathomimetic responses, *in vivo* phosphoinositide hydrolysis, and pilocarpine-induced seizure activity. *Eur J Neurosci* 17: 1403–1410.
- Bymaster FP, Felder CC (2002). Role of the cholinergic muscarinic system in bipolar disorder and related mechanism of action of antipsychotic agents. *Mol Psychiatry* 7(Suppl 1): S57–S63.
- Cade JF (1999). John Frederick Joseph Cade: family memories on the occasion of the 50th anniversary of his discovery of the use of lithium in mania. 1949. *Aust N Z J Psychiatry* 33: 615–622.
- Carli M, Anand-Srivastava MB, Molina-Holgado E, Dewar KM, Reader TA (1994). Effects of chronic lithium treatments on central dopaminergic receptor systems: G proteins as possible targets. *Neurochem Int* 24: 13–22.
- Carpenter TL, Pazdernik TL, Levant B (2003). Differences in quinpirole-induced local cerebral glucose utilization between naive and sensitized rats. *Brain Res* 964: 295–301.
- Carvey PM, Kao LC, Zhang TJ, Amdur RL, Lin DH, Singh R et al (1990). Dopaminergic alterations in cotreatments attenuating haloperidol-induced hypersensitivity. *Pharmacol Biochem Behav* 35: 291–300.
- Casebolt TL, Jope RS (1989). Long-term lithium treatment selectively reduces receptor-coupled inositol phospholipid hydrolysis in rat brain. *Biol Psychiatry* 25: 329–340.
- Chang MCJ, Arai T, Freed LM, Wakabayashi S, Channing MA, Dunn BB et al (1997). Brain incorporation of [1-¹⁴C]-arachidonate in normocapnic and hypercapnic monkeys, measured with positron emission tomography. *Brain Res* 755: 74–83.
- Chang MCJ, Grange E, Rabin O, Bell JM, Allen DD, Rapoport SI (1996). Lithium decreases turnover of arachidonate in several brain phospholipids. *Neurosci Lett* 220: 171–174.
- Chen G, Hasanat KA, Bebchuk JM, Moore GJ, Glitz D, Manji HK (1999). Regulation of signal transduction pathways and gene expression by mood stabilizers and antidepressants. *Psychosom Med* 61: 599–617.
- Colin SF, Chang HC, Mollner S, Pfeuffer T, Reed RR, Duman RS et al (1991). Chronic lithium regulates expression of adenylate cyclase and Gi-protein alpha subunit in rat cerebral cortex. *Proc Natl Acad Sci USA* 88: 10634–10637.
- Cooper JR, Bloom FE, Roth RH (1996). *The Biochemical Basis of Neuropharmacology* 7th edn. Oxford University Press: Oxford.
- Cory-Slechta DA, Zuch CL, Fox RA (1996). Comparison of the stimulus properties of a pre- vs a putative postsynaptic dose of quinpirole. *Pharmacol Biochem Behav* 55: 423–432.
- Dalen P, Steg G (1973). Lithium and levodopa in parkinsonism. *Lancet* 1: 936–937.
- Defagot MC, Malchiodi EL, Villar MJ, Antonelli MC (1997). Distribution of D₄ dopamine receptor in rat brain with sequence-specific antibodies. *Brain Res Mol Brain Res* 45: 1–12.
- DeGeorge JJ, Nariai T, Yamazaki S, Williams WM, Rapoport SI (1991). Arecoline-stimulated brain incorporation of intravenously administered fatty acids in unanesthetized rats. *J Neurochem* 56: 352–355.
- DeGeorge JJ, Noronha JG, Bell JM, Robinson P, Rapoport SI (1989). Intravenous injection of [1-¹⁴C]arachidonate to examine regional brain lipid metabolism in unanesthetized rats. *J Neurosci Res* 24: 413–423.
- Delfs JM, Kelley AE (1990). The role of D₁ and D₂ dopamine receptors in oral stereotypy induced by dopaminergic stimulation of the ventrolateral striatum. *Neuroscience* 39: 59–67.
- DeMar JC, Ma K, Bell J, Rapoport SI (2004). Brain conversion of linoleic to arachidonic acid is not a major source of the arachidonate found in brain phospholipids of adult rats. Abstr International Workshop on Brain Uptake and Utilization of Fatty Acids, Lipids & Lipoproteins Bethesda, MD. October 7–9.
- Dennis EA (1994). Diversity of group types, regulation, and function of phospholipase A₂. *J Biol Chem* 269: 13057–13060.
- Di Marzo V, Vial D, Sokoloff P, Schwartz JC, Piomelli D (1993). Selection of alternative G-mediated signaling pathways at the dopamine D₂ receptor by protein kinase C. *J Neurosci* 13: 4846–4853.
- Drevets WC, Price JL, Simpson Jr JR, Todd RD, Reich T, Vannier M et al (1997). Subgenual prefrontal cortex abnormalities in mood disorders. *Nature* 386: 824–827.
- Dwivedi Y, Pandey GN (1997). Effects of subchronic administration of antidepressants and anxiolytics on levels of the alpha subunits of G proteins in the rat brain. *J Neural Transm* 104: 747–760.
- Dziedzicka-Wasylewska M, Wedzony K (1996). The effect of prolonged administration of lithium on the level of dopamine D₂ receptor mRNA in the rat striatum and nucleus accumbens. *Acta Neurobiol Exp (Wars)* 56: 29–34.
- Ebert D, Feistel H, Loew T, Pirner A (1996). Dopamine and depression—striatal dopamine D₂ receptor SPECT before and after antidepressant therapy. *Psychopharmacology (Berlin)* 126: 91–94.
- Eilam D, Szechtman H (1989). Biphasic effect of D₂ agonist quinpirole on locomotion and movements. *Eur J Pharmacol* 161: 151–157.
- Engber TM, Susel Z, Kuo S, Chase TN (1990). Chronic levodopa treatment alters basal and dopamine agonist-stimulated cerebral glucose utilization. *J Neurosci* 10: 3889–3895.
- Engel J, Berggren U (1980). Effects of lithium on behaviour and central monoamines. *Acta Psychiatr Scand Suppl* 280: 133–143.
- Esposito G, Giovacchini G, Der MG, Vuong BK, Channing MA, Lerner A et al (2003). [¹⁴C]Arachidonic acid PET activation experiments in human subjects: a feasibility study using visual stimulation. *J Cereb Blood Flow Metab* 23(Suppl 1): 711.
- Evans MS, Zorumski CF, Clifford DB (1990). Lithium enhances neuronal muscarinic excitation by presynaptic facilitation. *Neuroscience* 38: 457–468.
- Fadda F, Serra G, Argiolas A, Melis MR, Gessa GL (1980). Effect of lithium on 3, 4-dihydroxyphenylacetic acid (DOPAC) concen-

- trations in different brain areas of rats. *Pharmacol Res Commun* 12: 689–693.
- Felder CC, Kanterman RY, Ma AL, Axelrod J (1990). Serotonin stimulates phospholipase A₂ and the release of arachidonic acid in hippocampal neurons by a type 2 serotonin receptor that is independent of inositolphospholipid hydrolysis. *Proc Natl Acad Sci USA* 87: 2187–2191.
- Fenwick PB, Robertson R (1983). Changes in the visual evoked potential to pattern reversal with lithium medication. *Electroencephalogr Clin Neurophysiol* 55: 538–545.
- Fisher G, Pelonero AL, Ferguson C (1991). Mania precipitated by prednisone and bromocriptine. *Gen Hosp Psychiatry* 13: 345–346.
- Fitzpatrick F, Soberman R (2001). Regulated formation of eicosanoids. *J Clin Invest* 107: 1347–1351.
- Folch J, Lees M, Sloane Stanley GH (1957). A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* 226: 497–509.
- Friedman E, Gershon S (1973). Effect of lithium on brain dopamine. *Nature* 243: 511–520.
- Gallager DW, Pert A, Bunney Jr WE (1978). Haloperidol-induced presynaptic dopamine supersensitivity is blocked by chronic lithium. *Nature* 273: 309–312.
- Gehlert DR (1993). Quantitative autoradiography of Gpp(NH)p sensitive and insensitive [³H]quinpirole binding sites in the rat brain. *Synapse* 14: 113–120.
- Gerfen CR, Keefe KA, Gauda EB (1995). D₁ and D₂ dopamine receptor function in the striatum: coactivation of D₁- and D₂-dopamine receptors on separate populations of neurons results in potentiated immediate early gene response in D₁-containing neurons. *J Neurosci* 15: 8167–8176.
- Ghadrirani AM, Annable L, Belanger MC, Chouinard G (1996). A cross-sectional study of parkinsonism and tardive dyskinesia in lithium-treated affective disordered patients. *J Clin Psychiatry* 57: 22–28.
- Giovacchini G, Chang MC, Channing MA, Toczek M, Mason A, Bokde AL et al (2002). Brain incorporation of [¹¹C]arachidonic acid in young healthy humans measured with positron emission tomography. *J Cereb Blood Flow Metab* 22: 1453–1462.
- Goodnick PJ, Meltzer HY (1984). Treatment of schizoaffective disorders. *Schizophr Bull* 10: 30–48.
- Greenwood TA, Alexander M, Keck PE, McElroy S, Sadovnick AD, Remick RA et al (2001). Evidence for linkage disequilibrium between the dopamine transporter and bipolar disorder. *Am J Med Genet* 105: 145–151.
- Hamilton SE, Loose MD, Qi M, Levey AI, Hille B, McKnight GS et al (1997). Disruption of the m1 receptor gene ablates muscarinic receptor-dependent M current regulation and seizure activity in mice. *Proc Natl Acad Sci USA* 94: 13311–13316.
- Hayakawa T, Chang MC, Rapoport SI, Appel NM (2001). Selective dopamine receptor stimulation differentially affects [³H]arachidonic acid incorporation, a surrogate marker for phospholipase A₂-mediated neurotransmitter signal transduction, in a rodent model of Parkinson's disease. *J Pharmacol Exp Ther* 296: 1074–1084.
- Hegerl U, Herrmann WM, Ulrich G, Muller-Oerlinghausen B (1990). Effects of lithium on auditory evoked potentials in healthy subjects. *Biol Psychiatry* 27: 555–560.
- Hesketh JE, Nicolaou NM, Arbuthnott GW, Wright AK (1978). The effect of chronic lithium administration on dopamine metabolism in rat striatum. *Psychopharmacology (Berlin)* 56: 163–166.
- Ho AK, Loh HH, Craves F, Hitzemann RJ, Gershon S (1970). The effect of prolonged lithium treatment on the synthesis rate and turnover of monoamines in brain regions of rats. *Eur J Pharmacol* 10: 72–78.
- Holman RT (1986). Control of polyunsaturated acids in tissue lipids. *J Am Coll Nutr* 5: 183–211.
- Horvitz JC, Williams G, Joy R (2001). Time-dependent actions of D₂ family agonist quinpirole on spontaneous behavior in the rat: dissociation between sniffing and locomotion. *Psychopharmacology (Berlin)* 154: 350–355.
- Jakobsen SN, Wiborg O (1998). Selective effects of long-term lithium and carbamazepine administration on G-protein subunit expression in rat brain. *Brain Res* 780: 46–55.
- Janowsky DS, Overstreet DH (1995). The role of acetylcholine mechanisms in mood disorders. In: Bloom FE, Kupfer DJ (eds). *Psychopharmacology. The Fourth Generation of Progress*. Raven: New York. pp 945–956.
- Jones CR, Arai T, Bell JM, Rapoport SI (1996). Preferential *in vivo* incorporation of [³H]arachidonic acid from blood into rat brain synaptosomal fractions before and after cholinergic stimulation. *J Neurochem* 67: 822–829.
- Jope RS (1993). Lithium selectively potentiates cholinergic activity in rat brain. *Prog Brain Res* 98: 317–322.
- Jung H, Reme C (1994). Light-evoked arachidonic acid release in the retina: illuminance/duration dependence and the effects of quinacrine, mellitin and lithium. Light-evoked arachidonic acid release. *Graefes Arch, Clin Exp Ophthalmol* 32: 167–175.
- Katzenschlager R, Sampaio C, Costa J, Lees A (2003). Anticholinergics for symptomatic management of Parkinson's disease. *Cochrane Database System Rev* CD003735.
- Ketter TA, Kimbrell TA, George MS, Dunn RT, Speer AM, Benson BE et al (2001). Effects of mood and subtype on cerebral glucose metabolism in treatment-resistant bipolar disorder. *Biol Psychiatry* 49: 97–109.
- Koene P, Prinssen EP, Cools AR (1993). Involvement of the nucleus accumbens in oral behaviour in the freely moving rat. *Eur J Pharmacol* 233: 151–156.
- Lawler CP, Gilmore JH, Watts VJ, Walker QD, Southerland SB, Cook LL et al (1995). Interhemispheric modulation of dopamine receptor interactions in unilateral 6-OHDA rodent model. *Synapse* 21: 299–311.
- Lee H, Villacres NE, Rapoport SI, Rosenberger TA (2004). *In vivo* imaging of increased brain arachidonic acid metabolism in a lipopolysaccharide rat model of neuroinflammation. *J Neurochem* 91: 936–945.
- Lerer B (1985). Studies on the role of brain cholinergic systems in the therapeutic mechanisms and adverse effects of ECT and lithium. *Biol Psychiatry* 20: 20–40.
- Levant B, DeSouza EB (1993). Differential pharmacological profile of striatal and cerebellar dopamine receptors labeled by [³H]quinpirole: identification of a discrete population of putative D3 receptors. *Synapse* 14: 90–95.
- Levey AI, Hersch SM, Rye DB, Sunahara RK, Niznik HB, Kitt CA et al (1993). Localization of D₁ and D₂ dopamine receptors in brain with subtype-specific antibodies. *Proc Natl Acad Sci USA* 90: 8861–8865.
- LeWitt PA, Nyholm D (2004). New developments in levodopa therapy. *Neurology* 62: S9–S16.
- Li PP, Tam YK, Young LT, Warsh JJ (1991). Lithium decreases Gs, Gi-1 and Gi-2 alpha-subunit mRNA levels in rat cortex. *Eur J Pharmacol* 206: 165–166.
- Makrides M, Neumann MA, Byard RW, Simmer K, Gibson RA (1994). Fatty acid composition of brain, retina, and erythrocytes in breast- and formula-fed infants. *Am J Clin Nutr* 60: 189–194.
- Malmberg A, Mohell N (1995). Characterization of [³H]quinpirole binding to human dopamine D_{2A} and D₃ receptors: effects of ions and guanine nucleotides. *J Pharmacol Exp Ther* 274: 790–797.
- Massat I, Souery D, Del-Favero J, Van Gestel S, Serretti A, Macciardi F et al (2002). Positive association of dopamine D₂ receptor polymorphism with bipolar affective disorder in a European Multicenter Association Study of affective disorders. *Am J Med Genet* 114: 177–185.

- Matsumura M, Kojima J, Gardiner TW, Hikosaka O (1992). Visual and oculomotor functions of monkey subthalamic nucleus. *J Neurophysiol* 67: 1615–1632.
- Morrisett RA, Jope RS, Snead III OC (1987). Effects of drugs on the initiation and maintenance of status epilepticus induced by administration of pilocarpine to lithium-pretreated rats. *Exp Neurol* 97: 193–200.
- Mrzljak L, Bergson C, Pappy M, Huff R, Levenson R, Goldman-Rakic PS (1996). Localization of dopamine D₄ receptors in GABAergic neurons of the primate brain. *Nature* 381: 245–248.
- Nilsson CL, Hellstrand M, Ekman A, Eriksson E (1998). Direct dopamine D₂-receptor-mediated modulation of arachidonic acid release in transfected CHO cells without the concomitant administration of a Ca²⁺-mobilizing agent. *Br J Pharmacol* 124: 1651–1658.
- Ong WY, Sandhya TL, Horrocks LA, Farooqui AA (1999). Distribution of cytoplasmic phospholipase A₂ in the normal rat brain. *J Hirnforsch* 39: 391–400.
- Ormandy GC, Song L, Jope RS (1991). Analysis of the convulsant-potentiating effects of lithium in rats. *Exp Neurol* 111: 356–361.
- Pardue S, Rapoport SI, Bosetti F (2003). Co-localization of cytosolic phospholipase A(2) and cyclooxygenase-2 in Rhesus monkey cerebellum. *Brain Res Mol Brain Res* 116: 106–114.
- Paxinos G, Watson C (1987). *The Rat Brain in Stereotaxic Coordinates* 3rd edn. Academic Press: New York.
- Pearlson GD, Wong DF, Tune LE, Ross CA, Chase GA, Links JM et al (1995). *In vivo* D₂ dopamine receptor density in psychotic and nonpsychotic patients with bipolar disorder. *Arch Gen Psychiatry* 52: 471–477.
- Peet M, Peters S (1995). Drug-induced mania. *Drug Saf* 12: 146–153.
- Pert A, Rosenblatt JE, Sivitt C, Pert CB, Bunney Jr WE (1978). Long-term treatment with lithium prevents the development of dopamine receptor supersensitivity. *Science* 201: 171–173.
- Pfeilschifter J, Reme C, Dietrich C (1988). Light-induced phosphoinositide degradation and light-induced structural alterations in the rat retina are enhanced after chronic lithium treatment. *Biochem Biophys Res Commun* 156: 1111–1119.
- Pittman KJ, Jakubovic A, Fibiger HC (1984). The effects of chronic lithium on behavioral and biochemical indices of dopamine receptor supersensitivity in the rat. *Psychopharmacology (Berlin)* 82: 371–377.
- Post RM, Jimerson DC, Bunney Jr WE, Goodwin FK (1980). Dopamine and mania: behavioral and biochemical effects of the dopamine receptor blocker pimozide. *Psychopharmacology (Berlin)* 67: 297–305.
- Potash JB, Willour VL, Chiu YF, Simpson SG, MacKinnon DF, Pearlson GD et al (2001). The familial aggregation of psychotic symptoms in bipolar disorder pedigrees. *Am J Psychiatry* 158: 1258–1264.
- Qu Y, Chang L, Klaff J, Balbo A, Rapoport SI (2003). Imaging brain phospholipase A2 activation in awake rats in response to the 5-HT_{2A/2C} agonist (±)2, 5-dimethoxy-4-iodophenyl-2-aminopropane (DOI). *Neuropsychopharmacology* 28: 244–252.
- Rapoport SI (2001). *In vivo* fatty acid incorporation into brain phospholipids in relation to plasma availability, signal transduction and membrane remodeling. *J Mol Neurosci* 16: 243–261.
- Rapoport SI (2003). *In vivo* approaches to quantifying and imaging brain arachidonic and docosahexaenoic acid metabolism. *J Pediatr* 143: S26–S34.
- Reches A, Ebstein RP, Belmaker RH (1978). The differential effect of lithium on noradrenaline- and dopamine-sensitive accumulation of cyclic AMP in guinea pig brain. *Psychopharmacology (Berlin)* 58: 213–216.
- Reches A, Wagner HR, Jackson V, Fahn S (1982). Chronic lithium administration has no effect on haloperidol-induced supersensitivity of pre- and postsynaptic dopamine receptors in rat brain. *Brain Res* 246: 172–177.
- Reus B, Lorenzen A, Unsicker K (2001). Dopamine and epinephrine, but not serotonin, downregulate dopamine sensitivity in cultured cortical and striatal astroglial cells. *Receptors Channels* 7: 441–451.
- Rintala J, Seemann R, Chandrasekaran K, Rosenberger TA, Chang L, Contreras MA et al (1999). 85 kDa cytosolic phospholipase A₂ is a target for chronic lithium in rat brain. *Neuroreport* 10: 3887–3890.
- Robinson PJ, Noronha J, DeGeorge JJ, Freed LM, Nariai T, Rapoport SI (1992). A quantitative method for measuring regional *in vivo* fatty-acid incorporation into and turnover within brain phospholipids: review and critical analysis. *Brain Res Rev* 17: 187–214.
- Rondot P, Bathien N (1979). Abnormal movements induced by psychotropic drugs. *Encephale* 5: 41–48.
- Rosenberger TA, Villacreses NE, Hovda JT, Bosetti F, Weerasinghe G, Wine RN et al (2004). Rat brain arachidonic acid metabolism is increased by a 6-day intracerebral ventricular infusion of bacterial lipopolysaccharide. *J Neurochem* 88: 1168–1178.
- Roth BL, Willins DL, Kristiansen K, Kroeze WK (1998). 5-Hydroxytryptamine2-family receptors (5-hydroxytryptamine2A, 5-hydroxytryptamine2B, 5-hydroxytryptamine2C) where structure meets function. *Pharmacol Ther* 79: 231–257.
- Sansone ME, Ziegler DK (1985). Lithium toxicity: a review of neurologic complications. *Clin Neuropharmacol* 8: 242–248.
- Seeman P, Van Tol HH (1994). Dopamine receptor pharmacology. *Trends Pharmacol Sci* 15: 264–270.
- Senogles SE (2000). The D₂s dopamine receptor stimulates phospholipase D activity: a novel signaling pathway for dopamine. *Mol Pharmacol* 58: 455–462.
- Shafer RA, Levant B (1998). The D₃ dopamine receptor in cellular and organismal function. *Psychopharmacology (Berlin)* 135: 1–16.
- Shimizu T, Wolfe LS (1990). Arachidonic acid cascade and signal transduction. *J Neurochem* 55: 1–15.
- Sidhu A, Niznik HB (2000). Coupling of dopamine receptor subtypes to multiple and diverse G proteins. *Int J Dev Neurosci* 18: 669–677.
- Sokoloff L (1999). Energetics of functional activation in neural tissues. *Neurochem Res* 24: 321–329.
- Song L, Jope RS (1992). Chronic lithium treatment impairs phosphatidylinositol hydrolysis in membranes from rat brain regions. *J Neurochem* 58: 2200–2206.
- Sovner R, DiMascio A (1978). Extrapyramidal syndromes and other neurologic side effects of psychotropic drugs. In: Lipton MA, DiMascio A, Killam KF (eds). *Psychopharmacology: A Generation of Progress*. Raven Press: New York. pp 1021–1032.
- Staunton DA, Magistretti PJ, Shoemaker WJ, Bloom FE (1982a). Effects of chronic lithium treatment on dopamine receptors in the rat corpus striatum. I. Locomotor activity and behavioral supersensitivity. *Brain Res* 232: 391–400.
- Staunton DA, Magistretti PJ, Shoemaker WJ, Deyo SN, Bloom FE (1982b). Effects of chronic lithium treatment on dopamine receptors in the rat corpus striatum. II. No effect on denervation or neuroleptic-induced supersensitivity. *Brain Res* 232: 401–412.
- Stephenson DT, Lemere CA, Selkoe DJ, Clemens JA (1996). Cytosolic phospholipase A₂ (cPLA₂) immunoreactivity is elevated in Alzheimer's disease brain. *Neurobiol Dis* 3: 51–63.
- Strokin M, Sergeeva M, Reiser G (2003). Docosahexaenoic acid and arachidonic acid release in rat brain astrocytes is mediated by two separate isoforms of phospholipase A₂ and is differently regulated by cyclic AMP and Ca²⁺. *Br J Pharmacol* 139: 1014–1022.
- Sultzer DL, Cummings JL (1989). Drug-induced mania—causative agents, clinical characteristics and management. A retrospective analysis of the literature. *Med Toxicol Adverse Drug Exp* 4: 127–143.

- Tu B, Bazan NG (2003). Hippocampal kindling epileptogenesis upregulates neuronal cyclooxygenase-2 expression in neocortex. *Exp Neurol* 179: 167–175.
- Ulrich G, Herrmann WM, Hegerl U, Muller-Oerlinghausen B (1990). Effect of lithium on the dynamics of electroencephalographic vigilance in healthy subjects. *J Affect Disord* 20: 19–25.
- Vial D, Piomelli D (1995). Dopamine D₂ receptors potentiate arachidonate release via activation of cytosolic, arachidonic-specific phospholipase A₂. *J Neurochem* 64: 2765–2772.
- Vincent SL, Khan Y, Benes FM (1995). Cellular colocalization of dopamine D₁ and D₂ receptors in rat medial prefrontal cortex. *Synapse* 19: 112–120.
- Wajda IJ, Banay-Schwartz M, De Guzman T, Manigault I (1983). Dietary lithium during development: changes in amino acid levels, ionic content, and [³H]spiperone binding in the brain of rats. *Neurochem Res* 8: 637–648.
- Wang HY, Friedman E (1999). Effects of lithium on receptor-mediated activation of G proteins in rat brain cortical membranes. *Neuropharmacology* 38: 403–414.
- Wedzony K, Chocyk A, Mackowiak M, Fijał K, Czyrak A (2000). Cortical localization of dopamine D₄ receptors in the rat brain—immunocytochemical study. *J Physiol Pharmacol* 51: 205–221.
- Weerasinghe GR, Rapoport SI, Bosetti F (2004). The effect of chronic lithium on arachidonic acid release and metabolism in rat brain does not involve secretory phospholipase A₂ or lipoxygenase/cytochrome P450 pathways. *Brain Res Bull* 63: 485–489.
- Winitz S, Gupta SK, Qian NX, Heasley LE, Nemenoff RA, Johnson GL (1994). Expression of a mutant Gi2 alpha subunit inhibits ATP and thrombin stimulation of cytoplasmic phospholipase A₂-mediated arachidonic acid release independent of Ca²⁺ and mitogen-activated protein kinase regulation. *J Biol Chem* 269: 1889–1895.
- Wong DF, Pearlson GD, Tune LE, Young LT, Meltzer CC, Dannals RF et al (1997a). Quantification of neuroreceptors in the living human brain: IV. Effect of aging and elevations of D₂-like receptors in schizophrenia and bipolar illness. *J Cereb Blood Flow Metab* 17: 331–342.
- Wong DF, Pearlson GD, Tune LE, Young LT, Meltzer CC, Dannals RF et al (1997b). Quantification of neuroreceptors in the living human brain: IV. Effect of aging and elevations of D₂-like receptors in schizophrenia and bipolar illness. *J Cereb Blood Flow Metab* 17: 331–342.
- Yatham LN, Liddle PF, Lam RW, Shiah IS, Lane C, Stoessl AJ et al (2002). PET study of the effects of valproate on dopamine D(2) receptors in neuroleptic- and mood-stabilizer-naïve patients with nonpsychotic mania. *Am J Psychiatry* 159: 1718–1723.
- Zhou L, Vessby B, Nilsson A (2002). Quantitative role of plasma free fatty acids in the supply of arachidonic acid to extrahepatic tissues in rats. *J Nutr* 132: 2626–2631.