Occupancy of Striatal and Extrastriatal Dopamine D₂/D₃ Receptors by Olanzapine and Haloperidol

Robert M Kessler^{*,1,2}, Mohammad Sib Ansari¹, Patrizia Riccardi¹, Rui Li^{1,3}, Karuna Jayathilake², Benoit Dawant³ and Herbert Y Meltzer²

¹Department of Radiology, Vanderbilt University School of Medicine, Nashville, TN, USA; ²Department of Psychiatry, Vanderbilt University School of Medicine, Nashville, TN, USA; ³Department of Electrical Engineering and Computer Science, Vanderbilt University School of Engineering, Nashville, TN, USA

There have been conflicting reports as to whether olanzapine produces lower occupancy of striatal dopamine D₂/D₃ receptor than typical antipsychotic drugs and preferential occupancy of extrastriatal dopamine D₂/D₃ receptors. We performed [¹⁸F] fallypride PET studies in six schizophrenic subjects treated with olanzapine and six schizophrenic subjects treated with haloperidol to examine the occupancy of striatal and extrastriatal dopamine receptors by these antipsychotic drugs. [¹⁸F] setoperone PET studies were performed in seven olanzapine-treated subjects to determine 5-HT_{2A} receptor occupancy. Occupancy of dopamine D₂/D₃ receptors by olanzapine was not significantly different from that seen with haloperidol in the putamen, ventral striatum, medial thalamus, amygdala, or temporal cortex, that is, 67.5–78.2% occupancy; olanzapine produced no preferential occupancy of dopamine D₂/D₃ receptors in the ventral striatum, medial thalamus, amygdala, or temporal cortex. There was, however, significantly lower occupancy of substantia nigra/NTA dopamine D₂/D₃ receptors in olanzapine-treated compared to haloperidol-treated subjects, that is, 40.2 vs 59.3% ($p = 0.0014$, corrected for multiple comparisons); in olanzapine-treated subjects, the substantia nigra/NTA was the only region with significantly lower dopamine D₂/D₃ receptor occupancy than the putamen, that is, 40.2 vs 69.2% ($p < 0.001$, corrected for multiple comparison). Occupancy of 5-HT_{2A} receptors was 85–93% in the olanzapine-treated subjects. The results of this study demonstrated that olanzapine does not produce preferential occupancy of extrastriatal dopamine D₂/D₃ receptors but does spare substantia nigra/NTA receptors. Sparing of substantia nigra/NTA dopamine D₂/D₃ receptor occupancy may contribute to the low incidence of extrapyramidal side effects in olanzapine-treated patients.

Neuropsychopharmacology (2005) 30, 2283–2289. doi:10.1038/sj.npp.1300836; published online 3 August 2005

Keywords: dopamine D₂/D₃ receptor; olanzapine; haloperidol; atypical antipsychotic drug; substantia nigra; PET

INTRODUCTION

Olanzapine is an atypical antipsychotic drug (APD), that is, therapeutically efficacious with a significantly lower incidence of extrapyramidal side effects (EPS) than typical APDs such as haloperidol at clinically effective doses (Beasley *et al*, 1996). Olanzapine is chemically closely related to clozapine (Bymaster *et al*, 1996; Schotte *et al*, 1996). A number of animal studies have suggested that the atypical antipsychotic profiles of olanzapine and clozapine are related to their selective effects on cortical and limbic

regions and lesser effects on the dorsolateral striatum. Both clozapine and olanzapine produce c-fos induction in the medial prefrontal cortex, which is not seen with haloperidol, and greater induction of c-fos in the nucleus accumbens and lateral septal nucleus than in dorsolateral striatum, whereas haloperidol produces similar levels in the dorsolateral striatum and nucleus accumbens (Robertson and Fibiger, 1992; Robertson *et al*, 1994; Robertson and Fibiger, 1996). Chronic administration of both clozapine and olanzapine produce greater upregulation of the medial frontal cortical than dorsolateral striatal DA D₂/D₃ receptors (Janowsky *et al*, 1992; Tarazi *et al*, 2001). Consistent with the above hypothesis, [¹²³I] epidepride SPECT (Bigliani *et al*, 2000) and [⁷⁶Br] FLB457 PET (Xiberas *et al*, 2000) studies have reported preferential occupancy of temporal cortical DA D₂/D₃ receptors by olanzapine. This was not confirmed by a recent PET study using [¹¹C] raclopride and [¹¹C] FLB457, which failed to find preferential occupancy of extrastriatal DA D₂/D₃ receptors by olanzapine (Tauscher *et al*, 2002). In addition, some PET (Kapur *et al*, 1998;

*Correspondence: Dr RM Kessler, Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center, 1161 21st Avenue South, Suite CCC-1121 MCN, Nashville, TN 37232-2675, USA, Tel: +1 615 343 3938, Fax: +1 615 343 6531, E-mail: robert.kessler@vanderbilt.edu

Received 8 November 2004; revised 25 March 2005; accepted 9 June 2005

Online publication: 21 June 2005 at <http://www.acnp.org/citations/Npp062105050011/default.pdf>

Nordstrom *et al*, 1998) and SPECT (Raedler *et al*, 1999; Tauscher *et al*, 1999) studies have reported that therapeutic doses of olanzapine occupy levels of striatal DA D₂/D₃ receptors similar to that seen with typical APDs, that is, 65–75%, while other PET (Xiberas *et al*, 2000) and SPECT (Bigliani *et al*, 2000; Pilowsky *et al*, 1996) studies report that therapeutic doses of olanzapine occupy less than 50% of striatal DA D₂/D₃ receptors. Given these discordant results, we undertook a study of the occupancy of striatal and extrastriatal DA D₂/D₃ receptors in olanzapine and haloperidol-treated schizophrenic subjects using [¹⁸F] fallypride PET studies (Kessler *et al*, 2000; Mukherjee *et al*, 2002). As atypical APDs have been distinguished from typical APDs by a high 5-HT_{2A}/DA D₂/D₃ receptor affinity ratio (Meltzer *et al*, 1989, 2003) and olanzapine has a high 5-HT_{2A}/DA D₂/D₃ affinity ratio (Bymaster *et al*, 1996; Schotte *et al*, 1996), we also performed [¹⁸F] setoperone (Trichard *et al*, 1998; Kessler *et al*, 2002) PET studies of cortical 5-HT_{2A} receptor occupancy by olanzapine. Preliminary results from this study have been previously presented (Kessler *et al*, 2002).

METHODS

This study was conducted with the approval of the Vanderbilt University School of Medicine Intramural Research Board. All subjects were evaluated to assure that subjects met DSM IV criteria for schizophrenia, had a medical history, and a physical examination was performed. A CBC with differential, plasma electrolytes, BUN, Cr, plasma glucose, cholesterol, triglycerides, total protein, bilirubin, urine analysis, and urinary drug screen were obtained. Subjects were free of current substance dependence—other than tobacco and caffeine, major medical illness, history of major trauma, or other psychiatric or neurological disorders.

PET [¹⁸F] fallypride studies were performed in six subjects receiving olanzapine monotherapy (5 M, 1 F, mean age = 29.8 years, age range of 20–45 years) and six subjects receiving haloperidol monotherapy (4 M, 2 F, mean age of 34.3 years, age range of 21–45 years). Four of the six olanzapine-treated subjects were studied off-medication and following 3–6 weeks of olanzapine therapy (doses of 10 mg, *N* = 1; or 20 mg, *N* = 3). Regional receptor occupancies for these subjects were computed using their individual regional off-medication values. Two of the olanzapine-treated subjects were studied only while on medication (doses of 15 mg). Haloperidol-treated subjects received either depot haloperidol every 4 weeks (*N* = 4; doses of 75 mg, *N* = 1; 100 mg, *N* = 1; or 150 mg, *N* = 2), oral haloperidol (*N* = 1, daily dose of 10 mg), or a combination of depot and oral haloperidol (*N* = 1, 50 mg depot and alternating oral daily doses of 2 and 5 mg). Subjects receiving depot haloperidol were studied 3.5 weeks following their last dose. For the subjects studied only on medication, regional receptor occupancies were calculated using mean regional values of 10 off-medication schizophrenic subjects (5 M, 5 F, mean age of 33.0 years, age range of 20–45 years). Off-medication subjects were either previously unmedicated (*N* = 4) or self-withdrawn from all medications for at least 3 weeks. Off-medication subjects received no medications during their off-medication period

except for one subject who received 1 mg lorazepam p.o. during the later portion of the PET study.

PET [¹⁸F] setoperone studies (Petit-Taboue *et al*, 1996) were performed in seven olanzapine-treated schizophrenic subjects (7 M, mean age of 33.1, age range 19–47 years); all subjects were studied off medication initially and following 3–6 weeks of olanzapine therapy (doses of 5 mg, *N* = 2; 10 mg, *N* = 3; 20 mg, *N* = 2).

For studies of dopamine D₂/D₃ receptor occupancy, PET [¹⁸F] fallypride scans were performed using a GE Advance PET Scanner (resolution of 5–6 mm) in the 3D mode. [¹⁸F] fallypride (3.8–5.2 mCi, specific activity ≥ 3500 Ci/mmol) was administered as a slow bolus over a 20 s period and serial PET scans of the brain were obtained over a period of from 180 to 250 min as previous modeling studies have shown demonstrated stable fits using the reference region method (Lammertsma *et al*, 1996) in all brain regions from 180 min onward (Kessler *et al*, 2000). MRI scans of the brain were performed using a GE Signa LXi echospeed 1.5 T MRI scanner using sagittal and coronal T1-weighted thin section gradient echo acquisitions (IR SPGR, TE = 3.6, TR = 19, TR = 400, slice thickness of 1.2–1.5 mm), an axial spin density weighted FSE acquisition (TE = 19, TR = 5000, slice thickness of 3 mm), and an axial T2-weighted FSE acquisition (TE = 10.6, TR = 5000, slice thickness of 3 mm). PET scans were coregistered to each other and to the MRI scans using a mutual information rigid body algorithm (Maes *et al*, 1997; Wells *et al*, 1996). Regions of interest, that is, the putamen, caudate, ventral striatum, medial thalamus, amygdala, temporal cortex, and ventral midbrain/region of substantia nigra, were manually drawn by a neuroradiologist experienced in PET data analysis (RMK) on the MRI studies and transferred to the corresponding PET images. The ventral striatum was delineated using the criteria of Mawlawi *et al* (2001). Regional binding potentials were estimated using the reference region method (Lammertsma *et al*, 1996).

For studies of 5-HT_{2A} receptor occupancy, PET [¹⁸F] setoperone (Kapur *et al*, 1997; Petit-Taboue *et al*, 1996) studies were performed using an ECAT 933/08/16 PET scanner following the injection of 2.6–7.55 mCi (specific activity greater than 2000 Ci/mmol) of [¹⁸F] setoperone. Serial PET scans were obtained for up to 70 min following injection. Cortical regions of interest were manually delineated for planes above the level of the lateral ventricles, corresponding to levels 28–40 mm above the ACPC line in the Talairach atlas (Talairach and Tournoux, 1988). Binding potentials were calculated using the reference region method (Lammertsma *et al*, 1996; Bonab *et al*, 1998). Cortical 5-HT_{2A} receptor occupancies were calculated for each subject using their individual off-medication binding potential.

Regional 5-HT_{2A} or D₂/D₃ receptor occupancies were calculated using the following formula:

Percent occupancy in region i

$$= 1 - \left[\frac{Ri(\text{medicated})}{Ri(\text{off-medication})} \right] \times 100$$

where *Ri* refers to the binding potential in region '*i*' in the indicated state.

The differences in DA D₂/D₃ receptor occupancy in brain regions during treatment with olanzapine or haloperidol were evaluated using analysis of variance with drug and region as factors.

RESULTS

Regional occupancies of DA D₂/D₃ receptors for haloperidol and olanzapine are shown in Table 1 and Figure 1. Representative time-activity curves for an olanzapine-treated subject studied off-medication and following olanzapine therapy (20 mg/day) are shown in Figure 2. Analysis of variance for regional dopamine DA D₂/D₃ receptor occupancies with region and drug as factors, corrected for multiple comparisons, demonstrates no

significant differences between the olanzapine- and haloperidol-treated subjects in the putamen, ventral striatum, medial thalamus, amygdala, or temporal cortex. For olanzapine-treated subjects, comparison of occupancies in the putamen, ventral striatum, medial thalamus, amygdala, and temporal cortex demonstrated no significant differences in indicating that olanzapine did not produce preferential occupancy in extrastriatal regions at clinically therapeutic doses, that is, 10–20 mg. Similarly for haloperidol-treated subjects, no preferential occupancy of the ventral striatal, thalamic, amygdalar, or temporal cortical receptors was observed. The occupancy in the substantia

Table 1 Occupancy of Striatal and Extrastriatal DA D₂/D₃ Receptors Haloperidol and Olanzapine

Region	Percent occupancy	
	Haloperidol (n = 6)	Olanzapine (n = 6)
Putamen	76.5 ± 8.2	69.2 ± 10.2
Ventral striatum	75.5 ± 8.2	70.9 ± 6.9
Medial thalamus	78.2 ± 6.2	71.0 ± 9.0
Amygdala	75.6 ± 8.1	72.4 ± 6.7
Temporal cortex	70.9 ± 5.5	67.5 ± 7.1
Substantia nigra/VTA	59.3 ± 9.2	40.2 ± 12.2*

*Significantly different from regional occupancy for haloperidol, $p = 0.0014$ corrected for multiple comparisons using ANOVA with drug and region as factors.

Figure 1 Plots of regional DA D₂/D₃ receptor occupancy for haloperidol (left hand column) and olanzapine (right hand column)-treated subjects in the putamen, ventral striatum, substantia nigra, medial thalamus, amygdala, and temporal cortex.

Figure 2 Time-activity curves of regional [¹⁸F] fallypride uptake in a schizophrenic subject studied off-medication, that is, never medicated, and following 6 weeks of olanzapine therapy (20 mg/day). (a) and (b) show the putamenal, ventral striatal, and cerebellar time-activity curves prior to and following olanzapine treatment. (c) and (d) show time-activity curves for the thalamus, substantia nigra, temporal cortex, and cerebellum prior to and following olanzapine therapy. The time-activity curve of the substantia nigra shows a lesser decrement in uptake after olanzapine treatment than other extrastriatal regions.

Figure 3 Summed parametric images of dopamine D_2/D_3 receptor occupancy for olanzapine-treated subjects studied off-medication and after 3–6 weeks of olanzapine therapy ($N = 4$) which demonstrate different, that is, lower occupancy—40%, in the substantia nigra/VTA in comparison to the putamen and other brain regions—about 70%. The color scale has been adjusted to show lower occupancies in white and red while higher occupancies are shown in yellow and green.

nigra/VTA for olanzapine-treated subjects was significantly lower than in all other brain regions sampled for both olanzapine ($p < 0.001$ corrected for multiple comparisons) and for haloperidol treated subjects ($p \geq 0.0014$, corrected for multiple comparisons) including the substantia nigra/VTA in haloperidol-treated subjects (see Table 1 and Figure 1). Only one haloperidol-treated subject had a DA D_2/D_3 receptor occupancy in the substantia nigra, which fell in the range seen in olanzapine-treated subjects. In contrast, the olanzapine subjects with the highest putamenal occupancy, 86%, had a nigral/VTA occupancy of 55%, lower than all but one haloperidol-treated subject. Olanzapine appears to spare DA D_2/D_3 receptor occupancy in the substantia nigra (see Figure 3).

$[^{18}\text{F}]$ setoperone PET studies of cortical 5-HT $_{2A}$ occupancy indicate very high levels of cortical 5-HT $_{2A}$ receptor occupancy at all doses studied, that is, a 5 mg dose of olanzapine produced a mean occupancy of 85%, a 10 mg dose 88%, and a 20 mg dose 93%.

DISCUSSION

The main findings of this study are: (1) similar levels of occupancy of DA D_2/D_3 receptors in the putamen, ventral striatum, medial thalamus, amygdala, and temporal cortex with therapeutic doses of haloperidol and olanzapine in patients with schizophrenia; (2) significantly lower levels of DA D_2/D_3 receptor occupancy in the ventral midbrain/substantia nigra in olanzapine than haloperidol-treated subjects; and (3) over 80% occupancy of 5-HT $_{2A}$ receptors in the cortex with olanzapine at doses from 5 to 20 mg/day.

This study is consistent with a number of previous $[^{11}\text{C}]$ raclopride PET and $[^{123}\text{I}]$ IBZM SPECT studies in showing similar levels of striatal DA D_2/D_3 receptor occupancy by haloperidol and olanzapine at typical therapeutic doses, that is, 68–84% at doses of 10–20 mg of olanzapine (Kapur *et al*, 1998; Nordstrom *et al*, 1998; Raedler *et al*, 1999; Tauscher *et al*, 1999). Tauscher *et al* (2002) using $[^{11}\text{C}]$ FLB457 has recently reported similar levels of DA D_2/D_3 receptor occupancy in the thalamus and striatum in normal subjects receiving a single 15 mg dose of olanzapine, that is, peak occupancies of 70 and 81% occupancies, respectively. In contrast, both Bigliani *et al* (2000) and Xiberas *et al* (2000)

have reported preferential occupancy of thalamic and temporal cortical receptors by olanzapine using $[^{123}\text{I}]$ epidepride SPECT and $[^{76}\text{Br}]$ FLB457 PET studies. For doses of 10–20 mg the mean DA D_2/D_3 receptor occupancies reported in these studies were 41.3% (Bigliani *et al*, 2000) and 44.0% (Xiberas *et al*, 2000) in the striatum, 66.4% (Xiberas *et al*, 2000) in the thalamus, and 82.8% (Bigliani *et al*, 2000) and 81.4% (Xiberas *et al*, 2000) in the temporal cortex. The results in the present study are consistent with those of Tauscher *et al* (2002), who calculated regional receptor levels in the thalamus using the simplified reference region method (Gunn *et al*, 1997). Both Bigliani *et al* (2000) and Xiberas *et al* (2000) utilized ratio methods to estimate available regional DA D_2/D_3 receptor levels. The lower striatal DA D_2/D_3 receptor occupancy reported by Bigliani and Xiberas for olanzapine compared to $[^{11}\text{C}]$ raclopride studies (Kapur *et al*, 1998; Nordstrom *et al*, 1998) and the current study suggests that these ratios were calculated at a time prior to the attainment of a transient equilibrium in the striatum, resulting in an underestimation of striatal DA D_2/D_3 receptor occupancy (Olsson and Farde, 2001; Erlandsson *et al*, 2003). Previous modeling studies of $[^{123}\text{I}]$ epidepride have produced conflicting findings regarding whether temporal cortical: cerebellar ratios obtained at 3 or 4 h, the time used in the $[^{123}\text{I}]$ epidepride SPECT study of olanzapine (Bigliani *et al*, 2000), are significantly correlated with modeled estimates of the striatal and temporal cortical receptor density (Fujita *et al*, 1999; Erlandsson *et al*, 2003). The results of the current study do not support preferential occupancy of DA D_2/D_3 receptors in the ventral striatum, thalamus, amygdala, or temporal cortex by olanzapine. The level of DA D_2/D_3 receptor occupancies in the putamen, thalamus, amygdala, and temporal cortex were not significantly different for olanzapine- and haloperidol-treated subjects.

In olanzapine-treated subjects, only the ventral midbrain/substantia nigra demonstrated significantly different, that is, lower, DA D_2/D_3 receptor occupancy than that seen in haloperidol-treated subjects. Some of the difference between DA D_2/D_3 receptor occupancy in the ventral midbrain/substantia nigra and other brain regions may be due to the small size of this brain region, with a loss of quantitation due to the limited resolution of the scanner utilized (5–6 mm) and a resulting lowering of apparent occupancy

(Kessler *et al*, 1984). While a partial volume correction would be desirable for the substantia nigra, its exact borders are difficult to trace on high-resolution T1-weighted scans, making definition of a partial volume correction difficult. Partial voluming would affect ventral midbrain/substantia nigra occupancies in haloperidol and olanzapine-treated subjects in an identical fashion. The results of this study suggest that olanzapine produces lower occupancy of ventral midbrain/substantia nigra DA D₂/D₃ receptors than is seen with haloperidol.

There are a number of observations that suggest that DA D₂/D₃ receptor-mediated neurotransmission within the substantia nigra and/or VTA may be involved in mediating the atypical profile of olanzapine. These include the ability of olanzapine (Stockton and Rasmussen, 1996), like clozapine (Chiodo and Bunney, 1983), but unlike haloperidol (Chiodo and Bunney, 1983), to produce a selective depolarization of DA neurons in the VTA, but not in the substantia nigra, and the role of dopaminergic neurotransmission in the substantia nigra in motor behaviors (Gerhardt *et al*, 1999, 2002; Robertson and Robertson, 1989). Studies in animals have demonstrated that a number of motor behaviors require normal nigral as well as normal striatal dopaminergic function. Motor dysfunction in middle-aged and elderly monkeys compared to young monkeys was most closely correlated with declines in DA release in the substantia nigra as opposed to putamen (Gerhardt *et al*, 2002). In unilaterally MPTP-lesioned rhesus monkeys, a single intraventricular injection of GDNF produced improvement in motor function, which was correlated with DOPAC and HVA levels in the lesioned nigra but not in the striatum (Gerhardt *et al*, 1999). In unilaterally 6-OH-dopamine-lesioned rats, L-DOPA-induced rotational behavior was temporally correlated to the duration of increased DA levels in the substantia nigra rather than the striatum (Robertson and Robertson, 1989). In rats, neuroleptic-induced increases in hind limb muscle tone, a measure of rigidity, requires blockade of nigral DA D₂/D₃ receptors (Double and Crocker, 1995). It has been suggested that EPS requires blockade of both striatal and nigral DA D₂/D₃ receptors (Crocker and Hemsley, 2001; Hemsley and Crocker, 1999). It is of interest that amisulpride, a substituted benzamide-type atypical APD produced a wide range of occupancies of caudate and putamenal D₂/D₃ receptors including occupancy levels greater than 80% in 2/9 schizophrenia subjects (Vernaleken *et al*, 2004). No linear relationship could be found between D₂/D₃ receptor occupancy and EPS, suggesting that factors other than striatal D₂ occupancy are important with regard to EPS. The lower incidence of motor side effects seen with olanzapine may relate to the sparing of ventral midbrain/nigral DA D₂/D₃ receptor occupancy, which may confer a conditional freedom from EPS. In order for EPS to occur, higher levels of striatal DA D₂/D₃ receptor occupancy may be needed with olanzapine compared to haloperidol. If sparing of DA D₂/D₃ receptor occupancy occurs in the VTA as well as in the substantia nigra, this may have significant effect on cortical DA release, which may be important to the improvement in cognition produced by olanzapine and other related atypical APDs (Meltzer and McGurk, 1999).

The current study utilized off-medication schizophrenic subjects to compute occupancies for subjects studied only

on medication. A potential confound of using off-medication schizophrenic subjects is residual medication effects. While a comparison of schizophrenic and normal subjects will be published in a separate communication, it is noteworthy that putamenal binding potentials were nearly identical for off-medication schizophrenic subjects and normal controls, that is, 36.08 ± 4.53 for the 10 schizophrenic subjects and 37.34 ± 2.53 for the 10 matched normal control subjects, indicating no residual medication effects. Unlike the striatum where most studies indicate no changes in DA D₂/D₃ receptor levels in schizophrenics, there are now studies indicating lower levels of DA D₂/D₃ levels in the thalamus (Talvik *et al*, 2003; Yasuno *et al*, 2004) and temporal cortex (Tuppurainen *et al*, 2003; Buchsbaum *et al*, 2004). Using normal subjects may not be appropriate in studies of DA D₂/D₃ occupancy in these regions. The use of the reference tissue method with [¹⁸F] fallypride may be criticized due to the presence of low levels of DA D₂/D₃ receptors in the cerebellum. Comparison of regional binding potentials calculated using Logan plots with a metabolite corrected plasma input function with those calculated using the reference region method demonstrated a correlation coefficient greater than 0.99 with a slope of 1.0 (Kessler *et al*, 2000). To examine the fraction of cerebellar uptake due to binding to DA D₂/D₃ receptors, ratios of the uptake in the cerebellar reference region to white matter were calculated for the four subjects studied off-medication and following olanzapine therapy; post-mortem studies have shown no specific binding in white matter (unpublished data, R.M. Kessler). Ratios at 90 min after injection were 1.128 ± 0.046 in the off-medication state and 1.103 ± 0.042 in the medicated state. These observations indicate that for [¹⁸F] fallypride, the reference region method is an appropriate method for estimating regional DA D₂/D₃ receptor levels.

The high occupancy of 5-HT_{2A} receptors seen in this study, 88% or greater at therapeutic doses, 10 or 20 mg of olanzapine, is consistent with the results from previous studies (Kapur *et al*, 1998, 1999). Our results suggest that this occupancy is dose-dependent but more subjects need to be studied to establish this. A number of findings support the role of 5-HT_{2A} receptor blockade in producing an atypical APD profile. These include the ability of 5-HT_{2A} receptor blockade to decrease the effect of haloperidol to increase DA efflux in the ventral striatum (Liegeois *et al*, 2002), to reverse haloperidol induced catalepsy in animals (Lucas *et al*, 1997), to decrease the incidence of EPS effects in subjects receiving typical APDs (Bersani *et al*, 1990), and studies of binding profiles of APDs demonstrating that a high ratio of 5-HT_{2A}/D₂ affinities distinguishes atypical from typical APDs (Meltzer *et al*, 1989; Roth *et al*, 1995). 5-HT_{2A} receptor antagonists can ameliorate haloperidol induced catalepsy even without affecting striatal DA release (Lucas *et al*, 1997) and modulate nigral dopaminergic neurotransmission (Sorenson *et al*, 1993; Bruggeman *et al*, 2000; Timmerman *et al*, 1999; Andersson *et al*, 1995). This modulation of nigral dopaminergic function may be a factor in producing an atypical profile.

In conclusion, the principal findings in this study are the lack of preferential occupancy of cortical and other extrastriatal regions by olanzapine and the relative sparing of ventral midbrain/substantia nigra DA D₂/D₃ receptor

occupancy in olanzapine-treated subjects, and the high 5HT_{2A} receptor blockade seen with therapeutic doses of olanzapine. The lower incidence of EPS seen with olanzapine may be related to sparing of ventral midbrain/substantia nigra DA D₂/D₃ receptor occupancy. Further studies are needed to confirm these conclusions and to elucidate the role of the substantia nigra and VTA and 5-HT_{2A} receptor blockade in mediating an atypical antipsychotic profile.

ACKNOWLEDGEMENTS

This work was supported by Eli Lilly, Indianapolis, a grant—NIMH R01MH660890, The William T Warren Foundation, The Ritter Foundation, and a gift of Mr Donald Test.

REFERENCES

- Andersson JL, Nomikos GG, Marcus M, Hertel P, Mathe JM, Svensson TH *et al* (1995). Risperidone potentiates the stimulatory effects of raclopride on neuronal activity and dopamine release selectively in the mesolimbic dopaminergic system. *Naunyn-Schmiedeberg's Arch Pharmacol* 352: 374–385.
- Beasley CM, Tollefson G, Tran P, Saterlee W, Sanger T, Hamilton S *et al* (1996). Olanzapine vs placebo and haloperidol acute phase results of the North American double-blind olanzapine trial. *Neuropsychopharmacology* 14: 111–123.
- Bersani G, Grispi A, Marini S, Pasini A, Valducci M, Ciani N (1990). 5-HT_{2A} antagonist risperidone in neuroleptic-induced parkinsonism: a double-blind comparison with orphenadrine and placebo. *Clin Neuropharmacol* 13: 500–506.
- Bigliani V, Mulligan RS, Acton PD, Ohlsen RI, Pike VW, Ell PJ *et al* (2000). Striatal and temporal cortical D₂/D₃ receptor occupancy by olanzapine and sertindole *in vivo*: a [¹²³I] epidepride single photon emission tomography (SPET) study. *Psychopharmacology* 150: 132–140.
- Bonab AA, Fischman AJ, Alpert NM (1998). Estimation of binding potential for the 5HT₂ receptor ligand [¹⁸F]setoperone by a noninvasive reference region graphical method. In: Carson R, Herscovitch P, Daube-Witenspoon M (eds). *Quantitative Functional Brain Imaging with Positron Emission Tomography*. Academic Press: New York. pp 421–426.
- Bruggeman RM, Heeringa M, Westerink BHC, Timmerman W (2000). Combined 5-HT_{2A}/D₂/D₃ receptor blockade inhibits the firing rate of SNR neurons in the rat brain. *Prog Neuro-Psychopharmacol Biol Psychiatr* 24: 579–593.
- Buchsbaum MS, Christian BT, Lehrer DS, Mukherjee J, Shi B, Moutil J (2004). D₂/D₃/D₃ dopamine receptor binding in the thalamus of medication-naïve schizophrenics. *Int J Neuropsychopharm* 7: S430.
- Bymaster FP, Calligaro DO, Falcone JR, Marsh RD, Moore NA, Tye NC *et al* (1996). Radioreceptor binding profile of the atypical antipsychotic olanzapine. *Neuropsychopharmacology* 14: 87–96.
- Chiodo LA, Bunney BS (1983). Typical and atypical neuroleptics: differential effects of chronic administration on the activity of A9 and A10 midbrain dopaminergic neurons. *J Neurosci* 3: 1607–1619.
- Crocker AD, Hemsley KM (2001). An animal model of extrapyramidal side effects induced by antipsychotic drugs: relationship with D₂/D₃ dopamine receptor occupancy. *Prog Neuro-Psychopharmacol Biol Psychiatr* 25: 573–590.
- Double KL, Crocker AS (1995). Dopamine receptors in the substantia nigra are involved in the regulation of muscle tone. *Proc Natl Acad Sci USA* 92: 1669–1673.
- Erlandsson K, Bressan RA, Mulligan RS, Ell PJ, Cunningham VJ, Pilowsky LS (2003). Analysis of D₂/D₃ dopamine receptor occupancy with quantitative SPET using the high-affinity ligand [¹²³I] epidepride: resolving conflicting findings. *Neuroimage* 19: 1205–1214.
- Fujita M, Seibyl JP, Verhoeff NP, Ichise M, Baldwin RM, Zoghbi SS *et al* (1999). Kinetic and equilibrium analyses of [¹²³I] epidepride binding to striatal and extrastriatal dopamine D₂/D₃ receptors. *Synapse* 34: 290–304.
- Gerhardt GA, Cass WA, Huettl P, Brock S, Zhang ZM, Gash DM (1999). GDNF improves dopamine function in the substantia nigra but not the putamen of unilateral MPTP-lesioned rhesus monkeys. *Brain Res Rev* 817: 163–171.
- Gerhardt GA, Cass WA, Yi A, Zhang Z, Gash DM (2002). Changes in somatodendritic but not terminal dopamine regulation in aged rhesus monkeys. *J Neurochem* 80: 168–177.
- Gunn RN, Lammertsma AA, Hume SP, Cunningham VJ (1997). Parametric imaging of ligand-receptor binding in PET using a simplified reference region model. *Neuroimage* 6: 279–287.
- Hemsley KM, Crocker AD (1999). Raclopride and chlorpromazine, but not clozapine, increase muscle rigidity in the rat: relationship with D₂/D₃ dopamine receptor occupancy. *Neuropsychopharmacology* 21: 101–109.
- Janowsky A, Neve KA, Kinzie JM, Taylor B, dePaulis T, Belknap JK (1992). Extrastriatal dopamine D₂/D₃ receptors: distribution pharmacological characterization and region-specific regulation by clozapine. *J Pharmacol Exp Ther* 261: 1282–1290.
- Kapur S, Jones C, DaSilva J, Wilson A, Houle S (1997). Reliability of a simple non-invasive method for the evaluation of 5-HT_{2A} receptors using [¹⁸F]-setoperone PET imaging. *Nucl Med Com* 18: 395–399.
- Kapur S, Zipursky RB, Remington G, Jones C, DaSilva J, Wilson AA *et al* (1998). 5-HT_{2A} and D₂/D₃ receptor occupancy of olanzapine in schizophrenia: a PET investigation. *AM J Psychiatry* 155: 921–928.
- Kapur S, Zipursky RB, Remington G (1999). Clinical and theoretical implications of 5-HT_{2A} and D₂/D₃ receptor occupancy of clozapine, risperidone, and olanzapine in schizophrenia. *Am J Psychiatry* 156: 286–293.
- Kessler R, Ansari AS, Li R, Lee M, Schmidt D, Dawant B *et al* (2002). Occupancy of cortical and substantia nigra DA D₂/D₃ receptors by typical and atypical antipsychotic drugs. *Neuroimage* 16: 59.
- Kessler RM, Ellis JR, Eden M (1984). Analysis of emission tomographic scan data: limitations imposed by resolution and background. *J Comput Assist Tomogr* 8: 514–522.
- Kessler RM, Mason NS, Jones C, Ansari MS, Manning RF, Price RR (2000). [¹⁸F] N-allyl-5-fluoropropyl epidepride (fallypride): radiation dosimetry, quantification of striatal and extrastriatal dopamine receptors in man. *Neuroimage* 11: S32.
- Lammertsma AA, Bench CJ, Hume SP, Osman S, Gunn K, Brooks DJ *et al* (1996). Comparison of methods of analysis of clinical [¹¹C] raclopride studies. *J Cerebral Blood Flow Metab* 16: 42–52.
- Liegeois J-F, Ichikawa J, Meltzer HY (2002). 5HT_{2A} receptor antagonism potentiates haloperidol-induced dopamine release in rat medial prefrontal cortex and inhibits that in the nucleus accumbens in a dose-dependent manner. *Brain Res* 947: 157–165.
- Lucas G, Bonhomme N, DeDeurwaerdere P, Moal ML, Spampinato U (1997). 8-OH-DPAT, a 5-HT_{1A} agonist and risperidone, a 5-HT_{2A/C} agonist, reverse haloperidol-induced catalepsy in rats independently of striatal dopamine release. *Psychopharmacology* 131: 57–63.
- Maes F, Collignon A, Vandermuele D, Marchal G, Suetens P (1997). Multimodality image registration by maximization of mutual information. *IEEE Trans Med Imag* 16: 187–198.
- Mawlawi O, Martinez D, Slifstein M, Broft A, Chatterjee R, Hwang DR *et al* (2001). Imaging human mesolimbic dopamine

- transmission with positron emission tomography: I. Accuracy and precision of D(2) receptors parameter measurements in ventral striatum. *J Cereb Blood Flow Metab* 21: 1034–1057.
- Meltzer HY, Li Z, Kaneda Y, Ichikawa J (2003). Serotonin receptors: their key role in drugs to treat schizophrenia. *Progr Neuropsychopharmacol Biol Psychiatr* 27: 1159–1172.
- Meltzer HY, Matsubara S, Li JC (1989). Classification of typical and atypical antipsychotic drugs on the basis of dopamine D-1, C-2 and serotonin 2 pK_i values. *J Pharmacol Exp Ther* 251: 238–251.
- Meltzer HY, McGurk SR (1999). The effect of clozapine, risperidone and olanzapine on cognitive function in schizophrenia. *Schizophrenia Bull* 25: 233–255.
- Mukherjee J, Christian BT, Dunigan KA, Shi B, Narayanan TK, Satter M et al (2002). Brain images of ¹⁸F-fallypride in normal volunteers: blood analysis, distribution, test–retest studies, and preliminary assessment of sensitivity to aging effects on dopamine D-2/D-3 receptors. *Synapse* 46: 170–188.
- Nordstrom AL, Nyberg S, Olsson H, Farde L (1998). Positron emission tomography finding of a high striatal D2/D3 receptor occupancy in olanzapine-treated patients. *Arch Gen Psychiatry* 55: 283–284.
- Olsson H, Farde L (2001). Potentials and pitfalls using high affinity radioligands in PET and SPET determinations on regional drug induced D2/D3 receptor occupancy—a simulation study based on experimental data. *NeuroImage* 14: 936–945.
- Petit-Taboue MC, Landeau B, Osmont A, Tillet I, Barre L, Baron JC (1996). Estimation of neocortical serotonin-2 receptor binding potential by single-dose fluorine-18-Setoperone kinetic PET data analysis. *J Nucl Med* 37: 95–104.
- Pilowsky LS, Busatto GF, Taylor M, Costa DC, Sharma T, Sigmundsson T et al (1996). Dopamine D2/D3 receptor occupancy *in vivo* by the novel atypical antipsychotic olanzapine—a ¹²³I IBZM single photon emission tomography (SPET) study. *Psychopharmacology* 124: 148–153.
- Raedler TJ, Knable MB, Lafargue T, Urbina RA, Egan MF, Pickar D et al (1999). *In vivo* determination of striatal dopamine D2/D3 receptor occupancy in patients treated with olanzapine. *Psychiatr Res: Neuroimaging Section* 90: 81–90.
- Robertson GS, Robertson HA (1989). Evidence that L-Dopa-induced rotational behavior is dependent on both striatal and nigral mechanisms. *J Neurosci* 9: 3326–3331.
- Robertson GS, Fibiger HC (1992). Neuroleptics increase c-fos expression in the forebrain: contrasting effects of haloperidol and clozapine. *Neuroscience* 46: 315–328.
- Robertson GS, Matsumura H, Fibiger HC (1994). Induction patterns of Fos-like immunoreactivity in the forebrain as predictors of atypical antipsychotic activity. *J PET* 271: 1058–1066.
- Robertson GS, Fibiger HC (1996). Effects of olanzapine on regional C-FOS expression in rat forebrain. *Neuropsychopharmacology* 14: 105–110.
- Roth BL, Tandra S, Burgess LH, Sibley DR, Meltzer HY (1995). D4 dopamine receptor binding affinity does not distinguish between typical and atypical antipsychotic drugs. *Psychopharmacology* 120: 365–368.
- Schotte A, Janssen PFM, Gommeren W, Luyten WH, VanGompel P, Lesage AS et al (1996). Risperidone compared with new and reference antipsychotic drugs: *in vitro* and *in vivo* receptor binding. *Psychopharmacology* 124: 57–73.
- Sorenson SM, Kehne JH, Fadaye GM, Humphreys TM, Ketteler HJ, Sullivan CK et al (1993). Characterization of the 5-HT_{2A} receptor antagonist MLD 100907 as a putative atypical antipsychotic: behavioral, electrophysiological and neurochemical studies. *J Pharmacol Exp Ther* 266: 684–691.
- Stockton ME, Rasmussen K (1996). Electrophysiological effects of olanzapine, a novel atypical antipsychotic, on A9 and A10 dopamine neurons. *Neuropharmacology* 14: 97–105.
- Talairach J, Tournoux P (1988). *Co-Planar Stereotaxic Atlas of the Human Brain*. Thieme: New York.
- Talvik M, Nordstrom AL, Olsson H, Hallidin C, Farde L (2003). Decreased thalamic D2/D3/D3 receptor binding in drug-naïve patients with schizophrenia: a PET study with [¹¹C]FLB457. *Int J Neuropsychopharmacol* 6: 361–370.
- Tarazi FI, Zhang K, Baldessarini RJ (2001). Long-term effects of olanzapine, risperidone, and quetiapine on dopamine receptor types in regions of rat brain: implications for antipsychotic drug treatment. *J Pharmacol Exp Ther* 297: 711–717.
- Tauscher J, Küfferle B, Asenbaum S, Fischer P, Pezawas L, Barnas C et al (1999). *In vivo* ¹²³I IBZM SPECT imaging of striatal dopamine-2 receptor occupancy I schizophrenic patients treated with olanzapine in comparison to clozapine and haloperidol. *Psychopharmacology* 141: 175–181.
- Tauscher J, Jones C, Remington G, Zipursky RB, Kapur S (2002). Significant dissociation of brain and plasma kinetics with antipsychotics. *Mol Psychiatr Res* 7: 317–321.
- Timmerman W, Heijmen M, Westerink BH, Bruggeman R, denBoer JA (1999). Effects of acute and chronic administration of olanzapine in comparison to clozapine and haloperidol on extracellular recordings of substantia nigra reticulata neurons in the rat brain. *Psychopharmacologia* 144: 286–294.
- Trichard C, Paillere-Martinot ML, Attar-Levy D, Recassens C, Monnet F, Martinot JL (1998). Binding of antipsychotic drugs to cortical 5-HT_{2A} receptors: a PET study of chlorpromazine, clozapine, and amisulpride in schizophrenic patients. *Am J Psychiatry* 155: 505–508.
- Tuppurainen H, Kuikka J, Viinamäki H, Husso-Saastamoinen M, Bergström K, Tiihonen J (2003). Extrastriatal dopamine D2/D3/3 receptor density and distribution in drug-naïve schizophrenic patients. *Mol Psychiatry* 8: 453–455.
- Vernaleken I, Siessmeier T, Buchholz HG, Hartter S, Hiemke C, Stoeter P et al (2004). High striatal occupancy of D2-like dopamine receptors by amisulpride in the brain of patients with schizophrenia. *Int J Neuropsychopharmacol* 7: 421–430.
- Wells WM, Viola P, Atsumi H, Nakajima S, Kikinis R (1996). Multimodal volume registration by maximization of mutual information. *Med Imag Anal* 1: 35–51.
- Xiberas X, Marinot JL, Mallet L, Artiges E, Loc'h C, Mazière B et al (2000). Extrastriatal and striatal D2/D3 dopamine receptor blockade with haloperidol or new antipsychotic drugs in patients with schizophrenia. *Br J Psychiatr* 179: 503–508.
- Yasuno F, Suhara T, Okubo Y, Sudo Y, Inoue M, Ichimiya T et al (2004). Low dopamine d(2) receptor binding in subregions of the thalamus in schizophrenia. *Am J Psychiatry* 161: 1016–1022.