

Reply to Ebenbichler *et al*

# Reply: ‘Second Generation Antipsychotic Drugs: Is There a Common Mechanism in the Development of Obesity?’

Helliner S Vestri<sup>\*1</sup>, Lidia Maianu<sup>1</sup>, Douglas R Moellering<sup>1</sup> and W Timothy Garvey<sup>1,2</sup>

<sup>1</sup>Department of Nutrition Sciences, University of Alabama at Birmingham, Birmingham, AL, USA; <sup>2</sup>Birmingham VA Medical Center, Birmingham, AL, USA

Neuropsychopharmacology (2007) 32, 2433–2434; doi:10.1038/sj.npp.1301367

Sir

We would like to thank Dr Ebenbichler and co-workers for their comments and express common interest in a multi-disciplinary hypothesis regarding the effects of second generation antipsychotic (SGA) drugs. Even though a unifying mechanism and primary sites of action have not yet been defined, their observations open up an excellent opportunity for a more global perspective. Given the complexity of the pathogenesis of type II diabetes, it is challenging to understand the role that SGAs play in the progression to insulin resistance and diabetes (O’Neill, 2005). This topic has been intensely debated in the literature and considerable evidence points to a variety of adverse physiological effects, including dyslipidemia (Meyer and Koro, 2004), hyperglycemia (Mackin *et al*, 2005), and increased leptin levels (Zhang *et al*, 2004). The mechanisms underlying the association of SGA medication with increased prevalence of weight gain and diabetes are multifactorial (Nasrallah, 2003). Certainly, the studies carried out by our laboratory showing a direct and differential effect of antipsychotic agents in adipocytes help to elucidate how cell enlargement may occur and ultimately play a role in body fat distribution and obesity that is seen with the use of SGAs. In addition, other studies focusing on different organs have reported evidence of adverse effects of the drugs (Schwenkreis and Assion, 2004; Horacek *et al*, 2006; Richards *et al*, 2006). Moreover, second-generation antipsychotic agents vary significantly in their risk for diabetes and their effects on glucose and lipid metabolism (Newcomer, 2005). More importantly yet, this risk may be correlated with drug concentration in blood (or plasma, or serum), and levels reached at the site of therapeutic and/or toxic effect (s) (Medori *et al*, 2006).

The comments of Dr Ebenbichler and co-workers are constructive and support the body of evidence behind the incidence of diabetes in schizophrenic patients on SGAs. Furthermore, the reports of Ebenbichler and co-workers on insulin signaling pathways in muscle (Engl *et al*, 2005) may create additional hypothesis-driven research directed toward understanding molecular mechanisms that promote poor cellular response in insulin target tissues. Conversely, attention should also be given to neuroendocrine aspects of schizophrenia when it comes to carbohydrate abnormalities. Neuroendocrine dysfunction has been documented in those subjects (Tandon and Halbreich, 2003), particularly hypercortisolemia, which increases the risk for comorbid diabetes even before treatment with SGAs is initiated. This might be a condition where plasma concentrations of SGAs would be critical to activate malfunctioning of insulin-target tissues. To make matters even more complex, genetic polymorphism has been implicated in the SGA-induced weight gain (Reynolds *et al*, 2005). Therefore, the association between these polymorphisms and the symptom responses to antipsychotic treatment and side effects of such treatment, notably weight gain, should be further investigated as well, and related to the presumed functionality of these polymorphisms. Such studies clearly demonstrate the potential of pharmacogenetics in optimizing treatment for the individual patient.

## DISCLOSURE/CONFLICT OF INTEREST

We disclose that there are no conflicts of interest for any of the authors relating to this submission.

## REFERENCES

- Engl J, Laimer M, Niederwanger A, Kranebitter M, Starzinger M, Pedrini MT *et al* (2005). Olanzapine impairs glycogen synthesis and insulin signaling in L6 skeletal muscle cells. *Mol Psychiatry* 10: 1089–1096.
- Horacek J, Bubenikova-Valesova V, Kopecek M, Palenicek T, Dockery C, Mohr P *et al* (2006). Mechanism of action of atypical

\*Correspondence: Dr HS Vestri, Department of Nutrition Sciences (Webb 244), University of Alabama at Birmingham, 1675 University Boulevard, Birmingham, AL-35294-3360, USA, Tel: +1 205 996 4014, Fax: +1 205 934 7049, E-mail: helliner@uab.edu  
Received 21 December 2006; accepted 4 January 2007

- antipsychotic drugs and the neurobiology of schizophrenia. *CNS Drugs* **20**: 389–409.
- Mackin P, Watkinson HM, Young AH (2005). Prevalence of obesity, glucose homeostasis disorders and metabolic syndrome in psychiatric patients taking typical or atypical antipsychotic drugs: a cross-sectional study. *Diabetologia* **48**: 215–221.
- Medori R, Mannaert E, Grunder G (2006). Plasma antipsychotic concentration and receptor occupancy, with special focus on risperidone long-acting injectable. *Eur Neuropsychopharmacol* **16**: 233–240.
- Meyer JM, Koro CE (2004). The effects of antipsychotic therapy on serum lipids: a comprehensive review. *Schizophr Res* **70**: 1–17.
- Nasrallah H (2003). A review of the effect of atypical antipsychotics on weight. *Psychoneuroendocrinology* **28**: 83–96.
- Newcomer JW (2005). Second-generation (atypical) antipsychotics and metabolic effects: a comprehensive literature review. *CNS Drugs* **19**: 1–93.
- O'Neill MF (2005). Study focuses in on potential cause of antipsychotic-induced diabetes. *Drug News Perspect* **18**: 481–482.
- Reynolds GP, Templeman LA, Zhang ZJ (2005). The role of 5-HT<sub>2C</sub> receptor polymorphisms in the pharmacogenetics of antipsychotic drug treatment. *Prog Neuropsychopharmacol Biol Psychiatry* **29**: 1021–1028.
- Richards AA, Hickman IJ, Wang AY, Jones AL, Newell F, Mowry BJ *et al* (2006). Olanzapine treatment is associated with reduced high molecular weight adiponectin in serum: a potential mechanism for olanzapine-induced insulin resistance in patients with schizophrenia. *J Clin Psychopharmacol* **26**: 232–237.
- Schwenkreis P, Assion HJ (2004). Atypical antipsychotics and diabetes mellitus. *World J Biol Psychiatry* **5**: 73–82.
- Tandon R, Halbreich U (2003). The second-generation 'atypical' antipsychotics: similar improved efficacy but different neuroendocrine side effects. *Psychoneuroendocrinology* **28**: 1–7.
- Zhang ZJ, Yao ZJ, Liu W, Fang Q, Reynolds GP (2004). Effects of antipsychotic on fat deposition and changes in leptin and insulin levels. Magnetic resonance imaging study of previously untreated people with schizophrenia. *Br J Psychiatry* **184**: 58–62.